

## ORIGINAL ARTICLES

- J. LORDAN, S. ALEGRE, G. ALINS, M. J. SARASÚA, A. MORTON, F. GARCÍA DEL PINO**  
Compatibility between *Forficula auricularia* and entomopathogenic nematodes to be used in pome fruit pest management **635–643**
- N. BERKVEN, J. VAN VAERENBERGH, M. MAES, T. BELIËN, N. VIAENE**  
Entomopathogenic nematodes fail to parasitize the woolly apple aphid *Eriosoma lanigerum* as their symbiotic bacteria are suppressed **644–655**
- N. TOFANGSAZI, R. H. CHERRY, S. P. ARTHURS**  
Efficacy of commercial formulations of entomopathogenic nematodes against tropical sod webworm, *Herpetogramma phaeopteralis* (Lepidoptera: Crambidae) **656–661**
- H. WU, Y. ZHANG, P. LIU, J. XIE, Y. HE, C. DENG, P. DE CLERCQ, H. PANG**  
*Cryptolaemus montrouzieri* as a predator of the striped mealybug, *Ferrisia virgata*, reared on two hosts **662–669**
- T. HYNES, R. J. Mc DONNELL, M. J. GORMALLY**  
Oviposition, adult longevity and temperature effects on the eggs of *Tetanocera elata* (Fab.) (Diptera: Sciomyzidae): a potential biocontrol agent for slugs **670–676**
- N. S. DIAS-PINI, C. S. B. DASILVA, M. F. G. V. PEÑAFLO, J. R. P. PARRA**  
Does host determine short-range flight capacity of trichogrammatids? **677–682**
- J. S. TERBLANCHE, Z. DE JAGER, L. BOARDMAN, P. ADDISON**  
Physiological traits suggest limited diapause response in false codling moth, *Thaumatotibia leucotreta* (Lepidoptera: Tortricidae) **683–691**
- Y.-Q. NIU, C. NANSEN, X.-W. LI, T.-X. LIU**  
Geographical variation of *Plutella xylostella* (Lepidoptera: Plutellidae) populations revealed by mitochondrial COI gene in China **692–700**
- M. MAIN, J. P. McCAFFREY, M. J. MORRA**  
Insecticidal activity of *Brassica juncea* seed meal to the fungus gnat *Bradysia impatiens* Johannsen (Diptera: Sciaridae) **701–707**
- MINI-REVIEW**
- H. CHEN, M. F. CHAUDHURY, A. SAGEL, P. L. PHILLIPS, S. R. SKODA**  
Artificial diets used in mass production of the New World screwworm, *Cochliomyia hominivorax* **708–714**