

CONTENTS

Freshwater ecosystem service footprint model: A model to evaluate regional freshwater sustainable development—A case study in Beijing-Tianjin-Hebei, China Y. Gao, Z. Feng, Y. Li and S. Li (Beijing, China)	1
Multicriteria development of synthetic indicators of the environmental profile of the Spanish regions J.M. Cabello, E. Navarro (Malaga, Spain), F. Prieto (Madrid, Spain), B. Rodríguez and F. Ruiz (Malaga, Spain)	10
A quantitative framework for assessing spatial flows of ecosystem services H.M. Serna-Chavez, C.J.E. Schulp, P.M. van Bodegom, W. Bouten, P.H. Verburg and M.D. Davidson (Amsterdam, The Netherlands).....	24
Recreation potential assessment at large spatial scales: A method based in the ecosystem services approach and landscape metrics F. Weyland and P. Larterra (Balcarce, Argentina)	34
Effects of nicosulfuron on the abundance and diversity of arbuscular mycorrhizal fungi used as indicators of pesticide soil microbial toxicity D.G. Karpouzas (Larisa, Greece), E. Papadopoulou (Larisa, Greece and Thessaloniki, Greece), I. Ipsilantis (Thessaloniki, Greece), I. Friedel (Dijon, France), I. Petric, N. Udikovic-Kolic (Zagreb, Croatia), S. Djuric (Sr-Novi-Sad, Serbia), E. Kandeler (Stuttgart, Germany), U. Menkissoglu-Spiroudi (Thessaloniki, Greece) and F. Martin-Laurent (Dijon, France)	44
A hyperspectral indicator system for rangeland degradation on the Tibetan Plateau: A case study towards spaceborne monitoring L.W. Lehnert, H. Meyer (Marburg, Germany), N. Meyer (Bonn, Germany), C. Reudenbach and J. Bendix (Marburg, Germany)	54
Chlorophyll-related indicators are linked to visible ozone symptoms: Evidence from a field study on native <i>Viburnum lantana</i> L. plants in northern Italy E. Gottardini, A. Cristofori, F. Cristofolini (San Michele all'Adige, Italy), C. Nali, E. Pellegrini (Pisa, Italy), F. Bussotti (Firenze, Italy) and M. Ferretti (Grosseto, Italy)	65
Exploring the self-assessment of sustainability indicators by different stakeholders A. Mascarenhas, L.M. Nunes and T.B. Ramos (Portugal)	75
Quantile-based grading improves the effectiveness of a multimetric index as a tool for communicating estuarine condition C.S. Hallett (Murdoch, Australia)	84
Quantitative assessment of soil functioning across a representative range of Dutch soils A.V. Semenov (Groningen, The Netherlands and Enkhuizen, The Netherlands), M.C. Pereira e Silva, J.F. Salles (Groningen, The Netherlands), H. Schmitt (Utrecht, The Netherlands) and J.D. van Elsas (Groningen, The Netherlands).....	88
Evaluating the efficiency of environmental monitoring programs C.R. Levine, R.D. Yanai (Syracuse, NY, USA), G.G. Lampman (Albany, NY, USA), D.A. Burns (Troy, NY, USA), C.T. Driscoll (Syracuse, NY, USA), G.B. Lawrence (Troy, NY, USA), J.A. Lynch (Washington, DC, USA) and N. Schoch (Gorham, ME, USA)	94
Silage corn production in conventional and conservation tillage systems. Part I: Sustainability analysis using combination of GIS/AHP and multi-fuzzy modeling E. Houshyar, M.J. SheikhDavoodi, M. Almassi, H. Bahrami (Ahvaz, Iran), H. Azadi (Belgium), M. Omid (Tehran, Iran), G. Sayyad (Ahvaz, Iran) and F. Witlox (Belgium)	102

Nitrogen deposition and the reduction of butterfly biodiversity quality in the Netherlands A. Feest (Bristol, UK and Bath, UK), C. van Swaay (Wageningen, Netherlands) and A. van Hinsberg (Bilthoven, Netherlands)	115
Mapping soil organic matter in low-relief areas based on land surface diurnal temperature difference and a vegetation index M.-S. Zhao (Nanjing, China, Beijing, China and Huainan, China), D.G. Rossiter (Nanjing, China and AE Enschede, The Netherlands), D.-C. Li, Y.-G. Zhao, F. Liu (Nanjing, China) and G.-L. Zhang (Nanjing, China and Beijing, China)	120
Physiological indices as indicators of ecosystem status in shellfish aquaculture sites R. Filgueira, T. Guyondet, L.A. Comeau (Moncton, Canada) and J. Grant (Halifax, Canada)	134
The influence of contrasting ambient SO ₂ concentrations in the Czech Republic in 1995 and in 2010 on the characteristics of spruce bark, used as an air quality indicator I. Suchara, J. Sucharová and M. Holá (Pruhonice, Czech Republic)	144
Development of an ecological security evaluation method based on the ecological footprint and application to a typical steppe region in China X. Li, M. Tian, H. Wang, H. Wang and J. Yu (Beijing, China)	153
Eco-footprint-based life-cycle eco-efficiency assessment of building projects J. Teng and X. Wu (Wuhan, China)	160
The effect of sampling intensity on patterns of rarity and community assessment metrics in stream fish samples J.J. Pritt and E.A. Frimpong (Blacksburg, VA, USA)	169
How do low magnitudes of hydrologic alteration impact riverine fish populations and assemblage characteristics? R.J. Rolls and A.H. Arthington (Nathan, Australia)	179
Constructing the FEEM sustainability index: A Choquet integral application M. Pinar (Ormskirk, UK), C. Cruciani, S. Giove (Venice, Italy) and M. Sostero (Venice, Italy and Pisa, Italy)	189
The contribution of stranding data to monitoring and conservation strategies for cetaceans: Developing spatially explicit mortality indicators for common dolphins (<i>Delphinus delphis</i>) in the eastern North-Atlantic H. Peltier (La Rochelle, France), P.D. Jepson (London, UK), W. Dabin (La Rochelle, France), R. Deaville (London, UK), P. Daniel (Toulouse, France), O. Van Canneyt and V. Ridoux (La Rochelle, France)	203