

CONTENTS

| | |
|---|-----|
| Trophic niche differentiation and microhabitat utilization revealed by stable isotope analyses in a dry-forest bat assemblage at Ankarana, northern Madagascar Melanie Dammhahn & Steven M. Goodman | 97 |
| Diversity of Malaysian insectivorous bat assemblages revisited Lee-Sim Lim, Adura Mohd-Adnan, Akbar Zubaid, Matthew J. Struebig & Stephen J. Rossiter | 111 |
| Spatiotemporal variation in phyllostomid bat assemblages over a successional gradient in a tropical dry forest in southeastern Brazil Luiz Alberto Dolabela Falcão, Mário Marcos do Espírito-Santo, Lemuel Olívio Leite, Raphael Neiva Souza Lima Garro, Luis Daniel Avila-Cabadilla & Kathryn Elizabeth Stoner | 123 |
| Fire resistance in a Caribbean dry forest: inferences from the allometry of bark thickness Brett T. Wolfe, Gabriel E. Saldaña Diaz & Skip J. Van Bloem | 133 |
| First comparison of quantitative estimates of termite biomass and abundance reveals strong intercontinental differences Cecilia A.L. Dahlsjö, Catherine L. Parr, Yadvinder Malhi, Homathevi Rahman, Patrick Meir, David T. Jones & Paul Eggleton | 143 |
| Reciprocal transplant experiment suggests host specificity of the mistletoe <i>Agelanthus natalitius</i> in South Africa Desale Y. Okubamichael, Megan E. Griffiths & David Ward | 153 |
| <i>Short Communications</i> | |
| Importance of interaction frequency in analysis of ant-plant networks in tropical environments Wesley Dáttilo, Ingrid Sánchez-Galván, Denise Lange, Kleber Del-Claro & Víctor Rico-Gray | 165 |
| Activity pattern of the neotropical marsupial <i>Didelphis aurita</i> in south-eastern Brazilian Atlantic Forest Mariana Silva Ferreira & Marcus Vinícius Vieira | 169 |
| Microhabitat and spatial complexity predict group size of the whip spider <i>Heterophrynus batesii</i> in Amazonian Ecuador Kenneth James Chapin | 173 |