

- Serra-Llinares RM, Bjørn PA, Finstad B, Nilsen R, Harbitz A, Berg M, Asplin L**
FEATURE ARTICLE: Salmon lice infection on wild salmonids in marine protected areas: an evaluation of the Norwegian 'National Salmon Fjords' 1–16
- Mirto S, Arigò C, Genovese L, Pusceddu A, Gambi C, Danovaro R**
Nematode assemblage response to fish-farm impact in vegetated (*Posidonia oceanica*) and non-vegetated habitats 17–28
- Bannister RJ, Valdemarsen T, Hansen PK, Holmer M, Ervik A**
Changes in benthic sediment conditions under an Atlantic salmon farm at a deep, well-flushed coastal site 29–47
- Pert CC, Fryer RJ, Cook P, Kilburn R, McBeath S, McBeath A, Matejusova I, Urquhart K, Weir SJ, McCarthy U, Collins C, Amundrud T, Bricknell IR**
Using sentinel cages to estimate infestation pressure on salmonids from sea lice in Loch Shieldaig, Scotland 49–59
- Beirão J, Purchase CF, Wringe BF, Fleming IA**
Wild Atlantic cod sperm motility is negatively affected by ovarian fluid of farmed females 61–70
- Brigolin D, Meccia VL, Venier C, Tomassetti P, Porrello S, Pastres R**
Modelling biogeochemical fluxes across a Mediterranean fish cage farm 71–88
- Bruno DW, Noguera PA, Black J, Murray W, Macqueen DJ, Matejusova I**
Identification of a wild reservoir of salmonid alphavirus in common dab *Limanda limanda*, with emphasis on virus culture and sequencing 89–98