

**Pernet F, Lagarde F, Le Gall P,  
Roque D'Orbcastel E**

Associations between farming practices and disease mortality of Pacific oyster *Crassostrea gigas* in a Mediterranean lagoon 99–106

**Petersen MA, Alam MA, Rahman MM,  
Ali ML, Mahmud S, Schlüter L,  
Jørgensen NOG**

Geosmin off-flavour in pond-raised fish in southern Bangladesh and occurrence of potential off-flavour producing organisms 107–116

**Murray AG, Hall M**

Treatment rates for sea lice of Scottish inshore marine salmon farms depend on local (sea loch) farmed salmon biomass and oceanography 117–125

**Johnsen IA, Fiksen Ø, Sandvik AD, Asplin L**

Vertical salmon lice behaviour as a response to environmental conditions and its influence on regional dispersion in a fjord system 127–141

**Warren-Myers F, Dempster T, Fjellidal PG,  
Hansen T, Jensen AJ, Swearer SE**

Stable isotope marking of otoliths during vaccination: a novel method for mass-marking fish 143–154

**Fjellidal PG, Wennevik V, Fleming IA,  
Hansen T, Glover KA**

Triploid (sterile) farmed Atlantic salmon males attempt to spawn with wild females 155–162

**Nikolaou M, Neofitou N, Skordas K,  
Castritsi-Catharios I, Tziantziou L**

Fish farming and anti-fouling paints: a potential source of Cu and Zn in farmed fish 163–171

**Noble TH, Smith-Keune C, Jerry DR**

Genetic investigation of the large-scale escape of a tropical fish, barramundi *Lates calcarifer*, from a sea-cage facility in northern Australia 173–183