

**Gatune WC, Vanreusel A, Ruwa R,
Bossier P, De Troch M**

FEATURE ARTICLE: Fatty acid profiling reveals a trophic link between mangrove leaf litter biofilms and the post-larvae of giant tiger shrimp *Penaeus monodon*

1–10

**Olsen LM, Hernández KL, Ardelan MV,
Iriarte JL, Sánchez N, González HE,
Tokle N, Olsen Y**

Responses in the microbial food web to increased rates of nutrient supply in a southern Chilean fjord: possible implications of cage aquaculture

11–27

**Law BA, Hill PS, Maier I, Milligan TG,
Page F**

Size, settling velocity and density of small suspended particles at an active salmon aquaculture site

29–42

**Šegvić-Bubić T, Talijančić I, Grubišić L,
Izquierdo-Gomez D, Katavić I**

Morphological and molecular differentiation of wild and farmed gilthead sea bream *Sparus aurata*: implications for management

43–54

**Ruiz Y, Suárez P, Alonso A, Longo E,
San Juan F**

Chemicals used for maintenance of wood rafts in mussel farms: evaluation of their potential toxic risk to mussel culture

55–66

Bengil F, Bizsel KC

Assessing the impact of aquaculture farms using remote sensing: an empirical neural network algorithm for Ildırı Bay, Turkey

67–79

**Samsing F, Oppedal F, Johansson D, Bui S,
Dempster T**

High host densities dilute sea lice *Lepeophtheirus salmonis* loads on individual Atlantic salmon, but do not reduce lice infection success

81–89

**Uglem I, Karlsen Ø, Sanchez-Jerez P,
Sæther BS**

REVIEW: Impacts of wild fishes attracted to open-cage salmonid farms in Norway

91–103

**Sarker P, Pilote A, Auffret M, Proulx É,
Villemur R, Deschamps MH, Vandenberg G**

Reducing geosmin off-flavor compounds and waste outputs through dietary phosphorus management in rainbow trout aquaculture

105–117