

Aquatic Ecosystem Health & Management.

Jul-Sep2014, Vol. 17 Issue 3

Deriving inherent optical property for highly turbid productive inland water from MERIS data by semi-analytical model: A case study in Taihu Lake, China.

By: Huang, C.; Chen, X.; Li, Y.; Yang, H.; Sun, D.; Le, C.; Xu, L. p252-260

The analysis of phytoplankton blooms off the Yangtze River Estuary in the spring of 2007.

By: Fu, Dongyang; Pan, Delu; Wang, Difeng; Zhang, Ying; Ding, Youzhan; Liu, Dazhao. p221-232.

Mapping the distribution of suspended particulate matter concentrations influenced by storm surge in the Yellow River Estuary using FY-3A MERSI 250-m data.

By: Zhao, W.; Sun, L.; Cui, T.; Cao, W.; Ma, Y.; Hao, Y. p290-298

Temporal variability of chromophoric dissolved organic matter in the Pearl River Estuary, China from 2003 to 2009.

By: Liu, Fenfen; Tang, Shilin; Chen, Chuqun p261-270

Flow prediction using ENVISAT RA-2 sea surface height validated model: A case study for the effect of Hong Kong-Zhuhai-Macau Bridge in the Pearl River Estuary, China.

By: Lu, Jianzhong; Wai, Onyx W. H.; Chen, Xiaoling; Zhang, Peng. p305-315

Remote sensing assessment of sediment variation in the Pearl River Estuary induced by Typhoon Vicente.

By: Ye, Haibin; Chen, Chuqun; Tang, Shilin; Tian, Liqiao; Sun, Zhaohua; Yang, Chaoyu; Liu, Fenfen. p271-279

Preliminary remote sensing observation of sea surface temperature increase during *Ulva prolifera* blooms.

By: Wang, Sufen; Tang, Danling. p299-304