

## CONTENTS

- Root growth of maize as studied with minirhizotrons and monolith methods  
*Rongwei Liao, Yueming Bai, Hong Liang, Shunqing An, Sanxue Ren, Yujing Cao, Zhangyan Le, Jianli Lu and Jingmiao Liu* 1343
- Non-destructive estimation of sunflower leaf area and leaf area index under different water regime managements  
*Ali Ghadami Firouzabadi, Mahmoud Raeini-Sarjaz, Ali Shahnazari and Hamid Zareabyaneh* 1357
- Availability of nutrients, productive properties, and quality of sunflower (*Helianthus annuus* L.) seeds in acid soils amended with different molar ratios between Ca and Mg  
*João Paulo Gonsiorkiewicz Rigon, Silvia Capuani, Rubiana Falopa Rossi, Leonardo Theodoro Büll and Dirceu Maximino Fernandes* 1369
- Effect of magnesium on crop yields within maize–green manure–T. aman rice cropping pattern on acid soil  
*Shamsun Noor, Sohela Akhter, Md. Monirul Islam, Nirmal Chandra Shil and Md. Abu Kawochar* 1381
- Potassium forms and quantity–intensity relationships in some wetland soils of Abeokuta, Southwestern Nigeria  
*Anjorin G. Ajiboye, Jamiu O. Azeez and Akinwande J. Omotunde* 1393
- Comparison of nitrogen fertilizer demand for wheat production between humid and semi-arid portions of the Argentinean Pampas using a mass balance method  
*Nicolás F. Romano, Roberto Alvarez, Alfredo A. Bono and Haydee S. Steinbach* 1409
- Amelioration of salinity on photosynthesis and some characteristics of three rapeseed (*Brassica napus*) cultivars under increased concentrations of carbon dioxide  
*Abbas Dehshiri, Seyed Ali Mohammad Modarres-Sanavy and Batoool Mahdavi* 1423
- Evaluation of water schemes for peanut, using CSM-CROPGRO-Peanut model  
*E. Amiri, A. Abdzad Gohari and A. Mianabadi* 1439
- Evapotranspiration model selection for estimation of actual evaporation from bare soil, as required in annual potential groundwater recharge studies of a semi-arid foothill region  
*Seyed Adib Banimahd, Davar Khalili, Ali Akbar Kamgar-Haghighi and Shahrokh Zand-Parsa* 1455
- Particle size fractions and pedogenesis of coastal plain sands  
*Jude C. Obi* 1473
- Iron requirement for soybean [*Glycine max* (L.) Merrill] inoculated with selected exotic and native isolates of *Bradyrhizobium* sp. under irrigated conditions  
*Anteneh Argaw* 1491

---

**CONTENTS****Review Article**

- Perspective on crop modelling in the management of intercropping systems  
*Vimbayi Grace Petrova Chimonyo, Albert Thembinkosi Modi and Tafadzwanashe Mabhaudhi* 1511

**Articles**

- Crop rotation effects on yield of oilseed rape, wheat and barley and residual effects on the subsequent wheat  
*Klaus Sieling and Olaf Christen* 1531
- Effect of integrated management on Fusarium wilt progression and grain yield of chickpea in Syria  
*Seid Ahmed Kemal, Mathew Abang, Muhammad Imtiaz and Asaad Nader* 1551
- Diverse responses of tolerant and sensitive lines of Chickpea to drought stress  
*Masoumeh Pouresmael, Ramazan Ali Khavari-Nejad, Javad Mozafari, Farzaneh Najafi and Foad Moradi* 1561
- Effect of soil water management and different sowing dates on maize yield and water use efficiency under drip irrigation system  
*Mohammad Taghi Feyzbakhsh, Behnam Kamkar, Hassan Mokhtarpour and Mohammad Esmail Asadi* 1581
- Grain oil and fatty acids composition of soybean affected by nano-iron chelate, chemical fertilizers and farmyard manure  
*Khosro Mohammadi* 1593
- Elemental composition and release characteristics of some South African fly ashes and their potential for land application  
*Hupenyu Allan Mupambwa and Pearson Nyari Stephano Mkeni* 1601
- Distribution of carbon, nitrogen, carbohydrate and calcium carbonate in aggregates of histosols in Shahrekord, Iran  
*Asiye Abasian, Ahmad Golchin, Mohsen Sheklabadi and Ali Beheshti Alagha* 1625
- Molecular characterization of salinity tolerance in wheat (*Triticum aestivum* L.)  
*Waqas Manzoor Bhutta and Muhammad Amjad* 1641
- Short Communication**
- Assessment of critical limit of boron for cowpea in piedmont soils of Arunachal Pradesh  
*Prankanu Debnath, Saroj Kumar Pattanaaik, Dinesh Sah and Ajai Kumar Pandey* 1649



## CONTENTS

- Foliar application of ascorbate enhances the physiological and biochemical attributes of maize (*Zea mays* L.) cultivars under drought stress  
*Ali Noman, Shafaqat Ali, Fomia Naheed, Qasim Ali, Mujahid Farid, Muhammad Rizwan and Muhammad Kashif Irshad* 1659
- Roles of glycine betaine in mitigating deleterious effect of salt stress on lettuce (*Lactuca sativa* L.)  
*Ertan Yildirim, Melek Ekinci, Metin Turan, Atilla Dursun, Raziye Kul and Fazilet Parlakova* 1673
- Halotolerant rhizobacteria: beneficial plant metabolites and growth enhancement of *Triticum aestivum* L. in salt-amended soils  
*Asif Raheem and Basharat Ali* 1691
- Cu-tolerant *Sinorhizobium meliloti* strain is beneficial for growth, Cu accumulation, and mineral uptake of alfalfa plants grown in Cu excess  
*Issam Nouairi, Moncef Mrabet, Mokded Rabhi, Haythem Mhadhbi and Kais Zribi* 1707
- Role of carrier-based biofertilizer in reclamation of saline soil and wheat growth  
*Tamoor Ul Hassan and Asghari Bano* 1719
- Metal micronutrients relationships in crop, soil, and common weeds of two maize (*Zea mays* L.) fields  
*Reza Ghasemi-Fasaei and Yalda Mansoorpoor* 1733
- The yield of wheat genotypes associated with yield components under irrigated and drought stress after anthesis  
*Sareh Sadat Sayyah, Mokhtar Ghobadi, Sirous Mansoorifar and Ali Reza Zebarjadi* 1743