

Contents

DISEASES

Occurrence of grey mould on <i>Bucida buceras</i> , a tree of subtropical origin, under indoor conditions – A disease note Á. Richweisz, M. Z. Németh, L. Kiss and T. Érsek	1
Variation of some morphological and molecular characteristics of Hungarian <i>Crivellia</i> and <i>Brachycladlum</i> isolates from opium poppy (<i>Papaver somniferum</i> L.) E. Kiss, L. Palkovics, E. Szathmáry and G. Nagy	5
Emergence of new virulence in <i>Plasmopara halstedii</i> (Sunflower downy mildew) N. Sakr	17
Defense related enzyme induction in coconut by endophytic bacteria (EPC 5) L. Rajendran, R. Akila, G. Karthikeyan, T. Raguchander and R. Samiyappan	29
Efficacy of wild medicinal plant extracts against predominant seed-borne fungi of broad bean cultivars Z. A. M. Baka	45

INSECT PESTS

<i>Brachycaudus</i> species on herbaceous plants along highways in Hungary Zs. Basky	67
New data to the mite fauna of Hungarian bamboo plantations J. Kentschán, A. Ács, G. Q. Wang and A. Neményi	77
Monitoring and testing different doses of disparlure for Indian gypsy moth, <i>Lymantria obfuscata</i> , in a temperate region of India (Kashmir Valley) B. Hussain, A. R. War, S. A. Ganie and S. Bilal	85
Economic impact of dissemination of management strategies for sucking insect pests on transgenic cotton in Punjab, India R. S. Chandi, V. Kumar, A. K. Dhawan and S. Saini	93
Prey suitability of <i>Tuta absoluta</i> Larvae (Lepidoptera: Gelechiidae) for three predatory phytoseiid mites (Acari: Phytoseiidae) under laboratory conditions A. M. Metwally, F. M. Momen, A. K. Nasr, A. A. Abdallah, I. M. Ebadah and Kh. M. Saleh.....	105
Virulence determination of <i>Beauveria bassiana</i> isolates on a predatory hemipteran, <i>Andrallus spinidens</i> Fabricius (Hemiptera: Pentatomidae) H. Firouzbakht, A. Zibaee, H. Hoda and M. M. Sohani.....	115

Evaluation of petroleum based horticulture oil for the management of the tea red spider
mite, *Oligonychus coffeae* Neitner (Acarina: Tetranychidae)

S. Roy, A. Rahman, A. Barua, F. R. Bora, G. Handique and D. Pujari 127

Influence of *Melissa officinalis* essential oil and its formulation on *Typhlodromips swirskii*
and *Neoseiulus barkeri* (Acari: Phytoseiidae)

M. M. Abdel Kader, F. M. Momen, E. A. Sammour, S. M. Aly and S. F. Fahim 139