

CONTENTS

| | |
|--|-----|
| Assessing the ecological integrity of endorheic wetlands, with focus on Mediterranean temporary ponds M. Van den Broeck (Leuven, Belgium and Rabat, Morocco), A. Waterkeyn (Leuven, Belgium and Arles, France), L. Rhazi (Rabat, Morocco), P. Grillas (Arles, France) and L. Brendonck (Leuven, Belgium) | 1 |
| Bryophytes as indicators of ancient woodlands in Schleswig-Holstein (Northern Germany) A. Mölder, M. Schmidt, F. Engel, E. Schönfelder (Göttingen, Germany) and F. Schulz (Reinbek, Germany)..... | 12 |
| Glutathione-dependent enzyme activities and concentrations of glutathione, vitamin E and sulfhydryl groups in barbel (<i>Barbus barbus</i>) and its intestinal parasite <i>Pomphorhynchus laevis</i> (Acanthocephala) T.B. Radovanović, M.D. Prokić, J.P. Gavrić, S.G. Despotović, B.R. Gavrilović, S.S. Borković-Mitić, S.Z. Pavlović and Z.S. Saičić (Belgrade, Serbia) | 31 |
| Impact of sewage pollution on two species of sea urchins in the Mediterranean Sea (Cortiou, France): Radial asymmetry as a bioindicator of stress Y. Savriama, L.C. Stige (Dijon, France), S. Gerber (Cambridge, UK), T. Pérez (Marseille, France), P. Alibert and B. David (Dijon, France)..... | 39 |
| Comparative ecology of vascular plant, bryophyte and testate amoeba communities in four <i>Sphagnum</i> peatlands along an altitudinal gradient in Switzerland I. Koenig (Neuchâtel, Switzerland), E. Feldmeyer-Christe (Birmensdorf, Switzerland) and E.A.D. Mitchell (Neuchâtel, Switzerland) | 48 |
| Urban expansion and its consumption of high-quality farmland in Beijing, China W. Song (Beijing, People's Republic of China), B.C. Pijanowski (West Lafayette, USA) and A. Tayyebi (West Lafayette, USA and Madison, WI, USA) | 60 |
| Assessing land clearing potential in the Canadian agriculture-forestry interface with a multi-attribute frontier approach D. Yemshanov (Sault Ste. Marie, Canada), F.H. Koch, K.H. Riitters (Research Triangle Park, NC, USA), B. McConkey (Swift Current, Canada), T. Huffman and S. Smith (Ottawa, Canada) | 71 |
| Developing an improved biomonitoring tool for fine sediment: Combining expert knowledge and empirical data M.D. Turley, G.S. Bilotta (East Sussex, UK), T. Krueger (Berlin, Germany), R.E. Brazier (Exeter, UK) and C.A. Extence (Spalding, UK)..... | 82 |
| Using range filling rather than prevalence of invasive plant species for management prioritisation: the case of <i>Spathodea campanulata</i> in the Society Islands (South Pacific) R. Pouteau (Koohnê (Koné), New Caledonia), J.-Y. Meyer (Papeete, French Polynesia) and S. Larrue (Clermont-Ferrand, France)..... | 87 |
| Best hyperspectral indices for tracing leaf water status as determined from leaf dehydration experiments Z. Cao, Q. Wang (Shizuoka, Japan) and C. Zheng (Shizuoka, Japan and Beijing, China) | 96 |
| Impact of nutrients and heavy metals capture by weeds on the growth and production of rice (<i>Oryza sativa</i> L.) irrigated with different water sources T.M. Galal (Cairo, Egypt) and H.S. Shehata (Zagazig, Egypt) | 108 |
| Developing and applying novel spectral feature parameters for classifying soil salt types in arid land P. Jin (Hangzhou, China), P. Li (Urumqi, China), Q. Wang (Shizuoka, Japan) and Z. Pu (Urumqi, China) | 116 |

| | |
|---|-----|
| Structural equation modeling as a tool to develop conservation strategies using environmental indicators: The case of the forests of the Magdalena river basin in Mexico City G. Santibáñez-Andrade, S. Castillo-Argüero (Mexico City, Mexico), E.V. Vega-Peña, R. Lindig-Cisneros (Morelia, Mexico) and J.A. Zavala-Hurtado (Mexico City, Mexico) | 124 |
| An ecosystem-based approach to evaluate the ecological quality of Mediterranean undersea caves P.-A. Rastorgueff (Marseille, France and South Africa), D. Bellan-Santini (Marseille, France), C.N. Bianchi (Genova, Italy), S. Bussotti (Nice, France), P. Chevaldonné (Marseille, France), P. Guidetti (Nice, France), J.-G. Harmelin (Marseille, France), M. Montefalcone, C. Morri (Genova, Italy), T. Perez, S. Ruitton, J. Vacelet and S. Personnic (Marseille, France) | 137 |
| Comparative evaluation of genotoxicity induced by nitrofurazone in two ciliated protozoa by detecting DNA strand breaks and DNA–protein crosslinks Y. Hong (Guangzhou, China and Xiamen, China), X. Lin, X. Cui, L. Zhou (Guangzhou, China), K.A.S. Al-Rasheid (Riyadh, Saudi Arabia) and J. Li (Guangzhou, China) | 153 |
| Detecting water quality improvement along the Catalan coast (Spain) using stress-specific biochemical seagrass indicators G. Roca (Barcelona, Spain and Blanes, Spain), T. Alcoverro (Blanes, Spain and Mysore, India), M. de Torres, M. Manzanera (Barcelona, Spain), B. Martínez-Crego (Faro, Portugal), S. Bennett (Perth, Australia), S. Farina (Blanes, Spain), M. Pérez and J. Romero (Barcelona, Spain) | 161 |
| Measuring CO ₂ emission linkages with the hypothetical extraction method (HEM) Y. Ali (Topi, Pakistan and Macerata, Marche) | 171 |
| Estuarine intertidal meiofauna and nematode communities as indicator of ecosystem’s recovery following mitigation measures A.S. Alves, A. Caetano (Coimbra, Portugal and Lisboa, Portugal), J.L. Costa, M.J. Costa (Lisboa, Portugal) and J.C. Marques (Coimbra, Portugal) | 184 |
| Integrating neglected ecological impacts of road transport into corporate management J. Friedrich (Göttingen, Germany) | 197 |
| A multiple-site dissimilarity measure for species presence/absence data and its relationship with nestedness and turnover C. Ricotta (Rome, Italy) and S. Pavoine (Paris, France and Oxford, UK) | 203 |
| Helping farmers to reduce herbicide environmental impacts F. Le Bellec, A. Vélú, P. Fournier, S. Le Squin, T. Michels, A. Tendero (Saint-Pierre, France) and C. Bockstaller (Colmar, France) | 207 |
| Towards the automated detection and occupancy estimation of primates using passive acoustic monitoring A.K. Kalan, R. Mundry, O.J.J. Wagner, S. Heinicke, C. Boesch and H.S. Köhl (Leipzig, Germany) | 217 |
| Key issues for the development and application of the species sensitivity distribution (SSD) model for ecological risk assessment F.-L. Xu, Y.-L. Li, Y. Wang, W. He, X.-Z. Kong, N. Qin, W.-X. Liu, W.-J. Wu (Beijing, China) and S.E. Jorgensen (Copenhagen Ø, Denmark) | 227 |
| An interval fuzzy model for magnetic biomonitoring using the specie <i>Tillandsia recurvata</i> L M.A.E. Chaparro A.E. (Tandil, Argentina and Mar del Plata, Argentina), M.A.E. Chaparro (Tandil, Argentina), A.G. Castañeda Miranda, H.N. Böhnelt (Juriquilla, Mexico) and A.M. Sinito (Tandil, Argentina) | 238 |
| Corrigendum to “Monetary valuation of road noise. Residential property prices as an indicator of the acoustic climate quality” [Ecol. Indic. 52 (2015) 472–479] D. Łowicki (Poznań, Poland) and S. Piotrowska (Oslo, Norway) | 246 |