

# CONTENTS

How to value biodiversity in environmental management? M. Laurila-Pant (Helsinki, Finland), A. Lehtikoinen (Kotka, Finland), L. Uusitalo (Helsinki, Finland) and R. Venesjärvi (Kotka, Finland) .....	1
Evolution of a multidimensional architectural landscape under urban regeneration: A case study of Jinan, China C. Xue, X. Zheng (Beijing, China), B. Zhang (Shandong, China) and Z. Yuan (Beijing, China) .....	12
Stable isotope-based statistical tools as ecological indicator of pollution sources in Mediterranean transitional water ecosystems G. Jona-Lasinio, M.L. Costantini, E. Calizza (Rome, Italy), A. Pollice (Bari, Italy), F. Bentivoglio, L. Orlandi, G. Careddu and L. Rossi (Rome, Italy) .....	23
Land-use changes assessed by overlay or mosaic methods: Which method is best for management planning? L.S. Bertolo, R.F. dos Santos (São Paulo, Brazil), P.M. de Agar and C.T.L. de Pablo (Madrid, Spain) .....	32
A method to evaluate the combined effect of tree species composition and woodland structure on indicator birds O. Dondina, V. Orioli (Milano, Italy), D. Massimino (Thetford, UK), G. Pinoli and L. Bani (Milano, Italy) .....	44
Presence of Cuckoo reliably indicates high bird diversity: A case study in a farmland area P. Tryjanowski (Poznań, Poland) and F. Morelli (Prague 6, Czech Republic) .....	52
Monitoring with lichens - Conductivity methods assess salt and heavy metal damage more efficiently than chlorophyll fluorescence O. Yemets, Y. Gauslaa and K.A. Solhaug (Ås, Norway) .....	59
Dung beetle persistence in human-modified landscapes: Combining indicator species with anthropogenic land use and fragmentation-related effects B.K.C. Filgueiras, M. Tabarelli, I.R. Leal (Recife, Brazil), F.Z. Vaz-de-Mello (Cuiabá, Brazil) and L. Iannuzzi (Recife, Brazil) .....	65
Do fish larvae have advantages over adults and other components for assessing estuarine ecological quality? S. Ramos (Porto, Portugal), H. Cabral (Lisboa, Portugal) and M. Elliott (Hull, UK) .....	74
Mercury, cadmium, zinc and copper concentrations and stable isotope ratios of carbon and nitrogen in tiger sharks ( <i>Galeocerdo cuvier</i> ) culled off Ishigaki Island, Japan T. Endo, O. Kimura, H. Ogasawara (Hokkaido, Japan), C. Ohta, N. Koga (Fukuoka, Japan), Y. Kato (Kagawa, Japan) and K. Haraguchi (Fukuoka, Japan) .....	86
A benchmarking and assessment framework to operationalise ecological indicators based on time series analysis W.N. Probst and V. Stelzenmüller (Hamburg, Germany) .....	94
Water productivity evaluation for grain crops in irrigated regions of China X. Cao, Y. Wang, P. Wu and X. Zhao (Yangling, China) .....	107
<i>Synechococcus</i> as an indicator of trophic status in the Cochin backwaters, west coast of India K.M. Rajaneesh, S. Mitbavkar, A.C. Anil and S.S. Sawant (India) .....	118

*Contents continued from outside back cover*

Forecasting sagebrush ecosystem components and greater sage-grouse habitat for 2050: Learning from past climate patterns and Landsat imagery to predict the future  
C.G. Homer (Sioux Falls, SD, USA), G. Xian (USA), C.L. Aldridge (Fort Collins, CO, USA), D.K. Meyer (USA), T.R. Loveland (Sioux Falls, SD, USA) and M.S. O'Donnell (Fort Collins, CO, USA). . . . . 131

Mismatches between ecosystem services supply and demand in urban areas: A quantitative assessment in five European cities  
F. Baró (Barcelona, Spain), D. Haase (Leipzig, Germany and Berlin, Germany), E. Gómez-Baggethun (Barcelona, Spain and Oslo, Norway) and N. Frantzeskaki (Rotterdam, The Netherlands) . . . . . 146

Mapping ecosystem services demand: A review of current research and future perspectives  
S. Wolff, C.J.E. Schulp and P.H. Verburg (Amsterdam, The Netherlands) . . . . . 159

Monitoring the Convention on Biological Diversity (CBD) framework using evaluation of effectiveness methods. The Italian case  
D. Marino, A. Marucci, M. Palmieri and P. Gaglioppa (Italy). . . . . 172