

CONTENTS

ORIGINAL ARTICLES

---

- 97 Influences of past application rates of nitrogen and a catch crop on soil microbial communities between an intensive rotation  
*Weishou Shen, Nan Gao, Ju Min, Weiming Shi, Xinhua He and Xiangui Lin*
- 107 Analysis of downy mildew data on field pea: an empirical comparison of two logistic models  
*Ewa Bakinowska, Wieslaw Pilarczyk and Bogna Zawieja*
- 117 Response of soil microbial community and diversity to increasing water salinity and nitrogen fertilization rate in an arid soil  
*Wei Min, Huijuan Guo, Wen Zhang, Guangwei Zhou, Lijuan Ma, Jun Ye, Yongchao Liang and Zhenan Hou*
- 127 Grain legumes differ in nitrogen accumulation and remobilisation during seed filling  
*Silvia Pampana, Alessandro Masoni and Iduna Arduini*
- 133 Evaluation of physiological traits of summer maize under drought stress  
*Weiming Yan, Yangquanwei Zhong and Zhouping Shangguan*
- 141 Saline water irrigation stimulate N<sub>2</sub>O emission from a drip-irrigated cotton field  
*Wen Zhang, Guangwei Zhou, Qi Li, Na Liao, Huijuan Guo, Wei Min, Lijuan Ma, Jun Ye and Zhenan Hou*
- 153 Functional traits associated with nitrogen use efficiency in wheat  
*Linnéa Asplund, Göran Bergkvist and Martin Weih*

SHORT COMMUNICATIONS

---

- 170 Photosynthate transport rather than photosynthesis rate is critical for low potassium adaptation of two cotton genotypes  
*Yanshu Hao, Jing Lei, Xiuwen Wu, Lishu Wu and Cuncang Jiang*
- 178 Targeted management of organic resources for sustainably increasing soil organic carbon: Observations and perspectives for resource use and climate adaptations in northern Ghana  
*William K. Heve, Jørgen E. Olesen, Ngonidzashe Chirinda and Samuel G.K. Adiku*
-