

CONTENTS

Research Reports

- A blend of benzoic acid and essential oil compounds as an alternative to antibiotic growth promoters in broiler diets**
P. C. Aristimunha, A. P. Rosa, L. S. Boemo, D. C. Garcez, D. P. Rosa, A. Londero, A. Scher, and J. Forgiarini 455
- Lysine requirements of Korean native ducklings for three weeks after hatch**
S. S. Wickramasuriya, Y. J. Yi, J. Yoo, J. C. Kim, K. N. Heo, and J. M. Heo 464
- Effects of enzymatic treatment and shell content of palm kernel expeller meal on performance, nutrient digestibility, and ileal bacterial population in broiler chickens**
B. Navidshad, J. B. Liang, M. F. Jahromi, A. Akhlaghi, and N. Abdullah 474
- Digestible lysine requirement for growing Japanese quails**
H. J. D. Lima, S. L. T. Barreto, J. L. Donzele, G. S. Souza, R. L. Almeida, I. F. F. Tinoco, and L. F. T. Albino 483
- Evaluation of whelks as an alternative source of calcium in laying hens**
J. L. Maclsaac, D. M. Anderson, and A. Safamehr 492
- Effect of dietary digestible lysine concentration on performance, egg quality, and blood metabolites in laying hens**
R. Akbari Moghaddam Kakhki, A. Golian, and H. Zarghi 506
- Comparison of flow behavior and physicochemical characteristics of low-cholesterol mayonnaises produced with cholesterol-reduced egg yolk**
M. Yüceer, H. İlyasoğlu, and B. Özçelik 518
- Supplemental defatted microalgae affects egg and tissue fatty acid composition differently in laying hens fed diets containing corn and flaxseed oil**
Jonggun Kim, Matthew Barcus, Andrew Magnuson, Ling Tao, and Xin Gen Lei 528
- Prevalence and antimicrobial resistance of *Salmonella* isolates in Moroccan laying hens farms**
N. Ziyate, B. Karraouan, A. Kadiri, S. Darkaoui, A. Soulaymani, and B. Bouchrif 539
- Does age of hot-blade trimming impact the performance and welfare of 2 strains of White Leghorn hens?**
K. Schwean-Lardner, C. B. Annett-Christianson, J. Rajendram, and H. L. Classen 547
- Effects of feeding dried broccoli floret residues on performance, ileal and total digestive tract nutrient digestibility, and selected microbial populations in broiler chickens**
Arif F. Mustafa and Bushansingh Baurhoo 561
- Growth performance and carcass characteristics of male broilers fed diets varying in digestible threonine to lysine ratio from 21 to 49 days of age**
W. A. Dozier, III, P. B. Tillman, and Z. Jiang 571

Effect of provitamin A biofortified maize inclusion on quality of meat from indigenous chickens

F. A. Odunitan-Wayas, U. Kolanisi, M. Chimonyo, and M. Siwela

581

Field Report

Routes of transmission of *Salmonella* and *Campylobacter* in breeder turkeys

M. D. Crespo, S. Kathariou, J. L. Grimes, N. A. Cox, R. J. Buhr, J. G. Frye,
W. G. Miller, C. R. Jackson, and D. P. Smith

591

Corrigenda

610