

CONTENTS

(Full texts are incorporated in CJELSEVIER, a file in the Chemical Journals Online database available on STN International; Abstracted, indexed in: Aluminium Abstracts; Anal. Abstr.; Biol. Abstr.; BIOSIS; Chem. Abstr.; Curr. Contents Phys. Chem. Earth Sci.; Engineered Materials Abstracts; Excerpta Medica; Index Med.; Life Sci.; Mass Spectrum. Bull.; Material Business Alerts; Metals Abstracts; Sci. Citation Index)

Growing degree days – Ecosystem indicator for changing diurnal temperatures and their impact on corn growth stages in Kansas A. Anandhi (Manhattan, KS, USA and Tallahassee, FL, USA)	149
Multiple-stressor effects on stream invertebrates: DNA barcoding reveals contrasting responses of cryptic mayfly species J.N. Macher (Bochum, Germany and Essen, Germany), R.K. Salis, K.S. Blakemore (Dunedin, New Zealand), R. Tollrian (Bochum, Germany), C.D. Matthaei (Dunedin, New Zealand) and F. Leese (Bochum, Germany and Essen, Germany)	159
Uncertainty and sensitivity analysis of West Java Water Sustainability Index – A case study on Citarum catchment in Indonesia I. Juwana (Bandung, Indonesia), N. Muttill and B.J.C. Perera (Melbourne, Australia).....	170
Structure and spatial patterns of macrobenthic community in Tai Lake, a large shallow lake, China D. Li (Nanjing, China), R.A. Erickson (La Crosse, WI, USA), S. Tang (Saskatoon, Canada), Y. Zhang, Z. Niu, H. Liu and H. Yu (Nanjing, China)	179
Towards an indicator of urban centrality? Exploring changes in present and resident population (1991–2011) in Greece L. Salvati, A. Sateriano (Rome, Italy) and K. Rontos (Mytilini, Greece)	188
CO ₂ emissions and urbanization correlation in China based on threshold analysis C. Zi, W. Jie and C. Hong-Bo (Beijing, China).....	193
A Relevance Index for the habitat areas of Natura 2000 Network based on their Rarity and Representativeness I. Duarte, F.C. Rego (Lisboa, Portugal), J.P. Casquilho (Lisboa, Portugal and Dili, Timor-Leste) and P. Arsénio (Lisboa, Portugal)	202
Toxicity evaluation of parboiled rice effluent using sperm quality of zebrafish as bioindicator M.D. Gerber (Pelotas, Brazil), A.S. Varela Junior, J.S. Caldas (Rio Grande, Brazil), C.D. Corcini (Rio Grande, Brazil and Pelotas, Brazil), T. Lucia Jr., L.B. Corrêa and É.K. Corrêa (Pelotas, Brazil)	214
Typology of Riverbed Structures and Habitats (TRiSHa) – A new method for a high resolution characterization of the spatial distribution and temporal dynamic of riverbed substrates and microhabitats M. Groll, A. Thomas, L. Jungermann and K. Schäfer (Germany)	219
Odonates as indicators of the ecological integrity of the river corridor: Development and application of the Odonate River Index (ORI) in northern Italy B. Golfieri (Padova, Italy), S. Hardersen (Marmirolo (MN), Italy), B. Maiolini (San Michele all'Adige (TN), Italy) and N. Surian (Padova, Italy).....	234
Ecological classification of lakes: Uncertainty and the influence of year-to-year variability M. Søndergaard (Silkeborg, Denmark and Beijing, China), S.E. Larsen, L.S. Johansson (Silkeborg, Denmark), T.L. Lauridsen and E. Jeppesen (Silkeborg, Denmark and Beijing, China)	248
The threshold responses of phytoplankton community to nutrient gradient in a shallow eutrophic Chinese lake X. Cao (Beijing, PR China and Shenzhen, PR China), J. Wang (Beijing, PR China), J. Liao (Ithaca, NY, USA), J. Sun and Y. Huang (Beijing, PR China).....	258
Characteristics of a landscape water with high salinity in a coastal city of China and measures for eutrophication control J. Xiong, X.C. Wang, Q. Zhang, R. Duan and N. Wang (Xi'an, China)	268

Development and implementation of a high nature value (HNV) farming indicator for Denmark A.K. Brunbjerg (Rønne, Denmark and Birmingham, UK), J. Bladt (Rønne, Denmark), M. Brink (Copenhagen, Denmark), J. Fredshavn (Denmark), P. Mikkelsen, J.E. Moeslund, B. Nygaard, F. Skov and R. Ejrnæs (Rønne, Denmark)	274
Synthesising the trait information of European Chironomidae (Insecta: Diptera): Towards a new database S.R.Q. Serra (Coimbra, Portugal), F. Cobo (Santiago de Compostela A Coruña, Spain and Vilagarcía de Arousa, Spain), M.A.S. Graça (Coimbra, Portugal), S. Dolédec (Villeurbanne, France) and M.J. Feio (Coimbra, Portugal)	282
GIS-based integrated evaluation of environmentally sensitive areas (ESAs) for land use planning in Langkawi, Malaysia N. Leman (Serdang, Malaysia and Bangi, Malaysia), M.F. Ramli (Serdang, Malaysia) and R.P.K. Khirotdin (Bangi, Malaysia).	293
Assessing water quality of five typical reservoirs in lower reaches of Yellow River, China: Using a water quality index method W. Hou, S. Sun, M. Wang, X. Li, N. Zhang, X. Xin, L. Sun, W. Li and R. Jia (Shandong Province, China)	309
A new view on EU agricultural landscapes: Quantifying patchiness to assess farmland heterogeneity C.J. Weissteiner, C. García-Feced and M.L. Paracchini (Ispra, Italy)	317
The effects of population density changes on ecosystem services value: A case study in Western Jilin, China L. Fei, Z. Shuwen, Y. Jiuchun, B. Kun, W. Qing, T. Junmei and C. Liping (Jilin Province, China)	328
A framework for quantifying environmental sustainability Z. Somogyi (Budapest, Hungary).	338
Impact of land use intensity on sandy desertification: An evidence from Horqin Sandy Land, China X. Ge, K. Dong (Wuhan, PR China), A.E. Luloff (University Park, PA, USA), L. Wang and J. Xiao (Wuhan, PR China)	346
Herbivory-induced stress: Leaf developmental instability is caused by herbivore damage in early stages of leaf development E. Alves-Silva and K. Del-Claro (Uberlândia, Brazil)	359
Spatial interpolation of N concentrations and $\delta^{15}\text{N}$ values in the moss <i>Hypnum cupressiforme</i> collected in the forests of Slovenia M. Skudnik, Z. Jeran, F. Batič and D. Kastelec (Ljubljana, Slovenia)	366
The performance of trait-based indices in an estuarine environment P. van der Linden (Coimbra, Portugal), A. Marchini (Pavia, Italy), M. Dolbeth (Aveiro, Portugal and Coimbra, Portugal), J. Patrício (Ispra, Italy), H. Veríssimo and J.C. Marques (Coimbra, Portugal)	378
Ecological Footprint: Refining the carbon Footprint calculation M.S. Mancini (Siena, Italy and Geneva, Switzerland), A. Galli (Geneva, Switzerland), V. Niccolucci (Siena, Italy), D. Lin (Oakland, CA, USA), S. Bastianoni (Siena, Italy), M. Wackernagel (Oakland, CA, USA) and N. Marchettini (Siena, Italy)	390
Multiple afforestation programs accelerate the greenness in the 'Three North' region of China from 1982 to 2013 Y. Zhang (Yangling, China and Norman, OK, USA), C. Peng (Yangling, China and Montreal, Canada), W. Li (Yangling, China), L. Tian (Yangling, China and TX, USA), Q. Zhu (Yangling, China), H. Chen (Chengdu, China), X. Fang (Nanjing, China), G. Zhang (Norman, OK, USA), G. Liu, X. Mu, Z. Li, S. Li, Y. Yang (Yangling, China), J. Wang (Norman, OK, USA) and X. Xiao (Norman, OK, USA and Shanghai, China)	404
Partitioning multiple-site tree-like beta diversity into turnover and nestedness components without pairwise comparisons Y. Chen	413
An interactive method to select a set of sustainable urban development indicators L. Tran (Knoxville, TN, USA)	418
Inter-regional linkage analysis of industrial CO ₂ emissions in China: An application of a hypothetical extraction method Y. Zhao, Y. Liu, S. Wang, Z. Zhang and J. Li (Beijing, China)	428
Do human disturbance variables influence more on fish community structure and function than natural variables in the Wei River basin, China? W. Wu (Jinan, China and Beijing, China), Z. Xu (Beijing, China), M.J. Kennard (Nathan, Australia), X. Yin (Dalian, China) and D. Zuo (Beijing, China)	438
Rating species sensitivity throughout gradient systems – a consistent approach for the Baltic Sea K.S. Schiele, A. Darr, M.L. Zettler (Warnemünde, Germany), T. Berg (Schönkirchen, Germany), M. Blomqvist (Stenhamra, Sweden), D. Daunys (Klaipeda, Lithuania), V. Jermakovs (Riga, Latvia), S. Korpinen (Helsinki, Finland), J. Kotta (Tallinn, Estonia), H. Nygård (Helsinki, Finland), M. von Weber (Güstrow, Germany), J. Voss (Flintbek, Germany) and J. Warzocha (Gdynia, Poland)	447
The soil quality concept as a framework to assess management practices in vulnerable agroecosystems: A case study in Mediterranean vineyards C. Salomé (Montpellier, France), P. Coll (Montpellier, France and Perpignan, France), E. Lardo (Matera, Italy), A. Metay (Montpellier, France), C. Villenave (Congénies, France and Montpellier, France), C. Marsden, E. Blanchart, P. Hinsinger and E. Le Cadre (Montpellier, France)	456

Performance evaluation of lake basin water governance using composite index P.E. Cookey, R. Darnsawasdi and C. Ratanachai (Songkhla, Thailand).....	466
Mapping technological and biophysical capacities of watersheds to regulate floods B. Mogollón (Bogotá, Colombia), A.M. Villamagna (Plymouth, NH, USA), E.A. Frimpong and P.L. Angermeier (Blacksburg, VA, USA).....	483
Ecological suitability evaluation for mountainous area development based on conceptual model of landscape structure, function, and dynamics J. Peng (Beijing, China and Shenzhen, China), J. Ma, Y. Du (Beijing, China), L. Zhang (Shenzhen, China) and X. Hu (Beijing, China).....	500
Developing a systematic simulation-based approach for selecting indicators in strategic cumulative effects assessments with multiple environmental valued components G.D. Sutherland (Victoria, Canada), F.L. Waterhouse (Nanaimo, Canada), J. Smith (Victoria, Canada), S.C. Saunders (Nanaimo, Canada), K. Paige (Victoria, Canada) and J. Malt (Surrey, Canada).....	512
Microzooplankton as a bioindicator of environmental degradation in the Amazon B.N.S. Costa, S.C.C. Pinheiro, L.L. Amado and M. de Oliveira Lima (Brazil).....	526
Source apportionment of non-storm water entries into storm drains using marker species: Modeling approach and verification Z. Xu, L. Wang, H. Yin, H. Li (Shanghai, China) and B.R. Schweigler (Stanford, CA, USA).....	546
Potential application of macroinvertebrates indices in bioassessment of Mexican streams E.C. Serrano Balderas (Montpellier, France and México, Mexico), C. Grac (Strasbourg, France), L. Berti-Equille (Montpellier, France and Doha, Qatar) and Ma.A. Armienta Hernandez (México, Mexico).....	558
Plant ecological traits highlight the effects of landscape on riparian plant communities along an urban-rural gradient E. Schwoertzig (Strasbourg, France), N. Poulin (Strasbourg, France), L. Hardion and M. Trémolières (Strasbourg, France).....	568
Assessing the sustainability of water companies: A synthetic indicator approach M. Molinos-Senante (Santiago, Chile), R.C. Marques (Lisbon, Portugal), F. Pérez, T. Gómez (Málaga, Spain), R. Sala-Garrido (Valencia, Spain) and R. Caballero (Málaga, Spain).....	577
New trophic indicators and target values for an ecosystem-based management of fisheries P. Bourdaud (Boulogne-sur-Mer, France), D. Gascuel (Rennes, France), A. Bentorcha (Plouzané, France) and A. Brind'Amour (Nantes, France).....	588
Different roles of environmental variables and spatial factors in structuring stream benthic diatom and macroinvertebrate in Yangtze River Delta, China S. Liu (Chongqing, China and Nanjing, China), G. Xie (Chongqing, China), L. Wang (Detroit, MI, USA), K. Cottenie (Guelph, Canada), D. Liu (Nanjing, China) and B. Wang (Nanjing, China).....	602
Composite indicator for measuring corporate sustainability M.P. Dočekalová and A. Kocmanová (Brno, Czech Republic).....	612
Evaluation of integrating topographic wetness index with backscattering coefficient of TerraSAR-X image for soil moisture estimation in a mountainous region S. Lei, H. Chen, Z. Bian and Z. Liu (Xuzhou, China).....	624
Features and influencing factors of carbon emissions indicators in the perspective of residential consumption: Evidence from Beijing, China Z. Wang and Y. Yang (Beijing, China).....	634
Identifying indicators and quantifying large-scale effects of dams on fishes A.R. Cooper, D.M. Infante (East Lansing, MI, USA), K.E. Wehrly (Ann Arbor, MI, USA), L. Wang (Detroit, MI, USA) and T.O. Brenden (East Lansing, MI, USA).....	646
The Enhanced Vegetation Index (EVI) as a proxy for diet quality and composition in a mountain ungulate M. Villamuelas (Barcelona, Spain and Legnaro, Italy), N. Fernández (Seville, Spain and Leipzig, Germany), E. Albanell, A. Gálvez-Cerón, J. Bartolomé (Bellaterra, Spain), G. Mentaberre, J.R. López-Olvera, X. Fernández-Aguilar, A. Colom-Cadena (Barcelona, Spain), J.M. López-Martín (Barcelona, Spain and Generalitat de Catalunya, Spain), J. Pérez-Barbería (Aberdeen, UK, Sevilla, Spain and Córdoba, Spain), M. Garel (Juvignac, France), I. Marco (Barcelona, Spain) and E. Serrano (Barcelona, Spain and Aveiro, Portugal).....	658
Distribution, bioavailability, and potential risk assessment of the metals in tributary sediments of Three Gorges Reservoir: The impact of water impoundment X. Wei, L. Han, B. Gao, H. Zhou, J. Lu and X. Wan (Beijing, China).....	667

The use of a functional approach as surrogate of Collembola species richness in European perennial crops and forests F. Reis, F. Carvalho, P. Martins da Silva, S. Mendes (Coimbra, Portugal), S.A.P. Santos (Bragança, Portugal) and J.P. Sousa (Coimbra, Portugal).....	676
Flexible risk metrics for identifying and monitoring conservation-priority species J.C. Stanton (La Crosse, WI, USA), B.X. Semmens (La Jolla, WI, USA), P.C. McKann (La Crosse, WI, USA), T. Will (Bloomington, CA, USA) and W.E. Thogmartin (La Crosse, MN, USA).....	683
Relating landscape to stream nitrate-N levels in a coastal eastern-Atlantic watershed (Portugal) Z. Teixeira and J.C. Marques (Coimbra, Portugal)	693
Does income growth relocate ecological footprint? A.A. Aşıcı and S. Acar (Istanbul, Turkey).....	707
The ecosystem service assessment challenge: Reflections from Flanders-REA S. Jacobs, T. Spanhove, L. De Smet, T. Van Daele, W. Van Reeth, P. Van Gossum, M. Stevens, A. Schneiders, J. Panis, H. Demolder, H. Michels, M. Thoonen, I. Simoens and J. Peymen (Belgium)	715
Quantile regression analysis as a predictive tool for lake macroinvertebrate biodiversity R. Fornaroli, R. Cabrini (Milano, Italy), S. Zaupa (Verbania Pallanza, Italy), R. Bettinetti (Como, Italy), M. Ciampittiello and A. Boggero (Verbania Pallanza, Italy)	728
The inorganic component of green roof substrates impacts the growth of Mediterranean plant species as well as the C and N sequestration potential S. Ondoño (Espinardo, Spain), J.J. Martínez-Sánchez (Cartagena, Spain) and J.L. Moreno (Espinardo, Spain)	739
Energy budgeting and emergy synthesis of rainfed maize-wheat rotation system with different soil amendment applications R.J. Singh, B.N. Ghosh, N.K. Sharma, S. Patra, K.S. Dadhwal and P.K. Mishra (Dehradun, India).....	753
Vertical concentration gradients of heavy metals in <i>Cladonia</i> lichens across different parts of thalli P. Osyczka, K. Rola and K. Jankowska (Kraków, Poland).....	766
Artificial neural networks as an indicator search engine: The visualization of natural and man-caused taxa variability D. Milošević (Nis, Serbia), D. Čerba (Osijek, Croatia), J. Szekeres, B. Csányi (Göd, Hungary), B. Tubić (Belgrade, Serbia), V. Simić (Kragujevac, Serbia) and M. Paunović (Belgrade, Serbia).....	777
Evaluating the effectiveness of overstorey cover as a surrogate for bird community diversity and population trends J.C. Pierson, A. Mortelliti, P.S. Barton, P.W. Lane and D.B. Lindenmayer (Canberra, Australia)	790
Applying indicators of disturbance from plant ecology to vertebrates: The hemeroby of bird species C. Battisti and G. Fanelli (Rome, Italy)	799
The effects of a 9-year nitrogen and water addition on soil aggregate phosphorus and sulfur availability in a semi-arid grassland R. Wang (Shenyang, China), C.A. Creamer (Glen Osmond, Australia), X. Wang, P. He (Shenyang, China and Beijing, China), Z. Xu and Y. Jiang (Shenyang, China).....	806
Estimation of big sagebrush leaf area index with terrestrial laser scanning P.J. Olsoy (Boise, ID, USA), J.J. Mitchell (Boone, NC, USA), D.F. Levia (Newark, DE, USA), P.E. Clark and N.F. Glenn (Boise, ID, USA).....	815
Spatial heterogeneity of ecosystem health and its sensitivity to pressure in the waters of nearshore archipelago C. Shen, H. Shi, W. Zheng and D. Ding (Qingdao City, China)	822
Evaluating ecosystem-level anthropogenic impacts in a stressed transitional environment: The case of the Seine estuary S. Tecchio (Caen, France), A. Chaalali (Caen, France and Nantes, France), A. Raoux (Caen, France), A. Tous Rius, J. Lequesne, V. Girardin (Caen, France), G. Lassalle (Cestas Cedex, France), M. Cachera (Plouzané, France), P. Riou (Port-en-Bessin, France), J. Lobry (Cestas, France), J.-C. Dauvin (Caen, France) and N. Niquil (Caen, France).....	833
Genetic diversity of riverine reed stands indicating the water regime of the habitat A.I. Engloner and D. Szegő (Budapest, Hungary).....	846
Response of macroalgae and macroinvertebrates to anthropogenic disturbance gradients in rocky shores P.A. Vinagre, A.J. Pais-Costa, R. Gaspar (Portugal), Á. Borja (Pasaia, Spain), J.C. Marques and J.M. Neto (Portugal).....	850
An approach to determining homogeneity of body-size spectrum of biofilm-dwelling ciliates for colonization surveys Z. Wang, G. Xu (Qingdao, China), Z. Yang (Qingdao, China and Xiamen, China) and H. Xu (Qingdao, China).....	865
Integrated assessment results depend on aggregation method and framework structure – A case study within the European Marine Strategy Framework Directive W.N. Probst (Hamburg, Germany) and C.P. Lynam (Suffolk, UK)	871

Assessing functional redundancy in chronically trawled benthic communities A. Muntadas (Barcelona, Spain), S. de Juan (Santiago, Chile) and M. Demestre (Barcelona, Spain)	882
Reduction and reallocation of water use of products in Beijing X. Zhao (Beijing, China), M. Tillotson (Leeds, UK), Z. Yang (Beijing, China), H. Yang (Duebendorf, Switzerland) and J. Liu (Beijing, China).....	893
Analysis of over-consumption of natural resources and the ecological trade deficit in China based on ecological footprints J. Gao and M. Tian (Nanjing, China)	899
Biological indicators track differential responses of pelagic and littoral areas to nutrient load reductions in German lakes F. Eigemann, U. Mischke, M. Hupfer (Berlin, Germany), J. Schaumburg (Wielenbach, Germany) and S. Hilt (Berlin, Germany)	905
Corporate sustainability for architecture engineering and construction (AEC) organizations: Framework, transition and implication strategies Y. Lu (Singapore, Singapore) and X. Zhang (Hong Kong)	911
Positioning of remotely sensed spectral heterogeneity in the framework of life cycle impact assessment on biodiversity B. Rugani and D. Rocchini (Belvaux, Luxembourg and Italy)	923
Benchmarking sustainability using indicators: An Indian case study S. Kwatra (New Delhi, India and Delhi, India), A. Kumar (Delhi, India), P. Sharma, S. Sharma and S. Singhal (New Delhi, India).....	928
Fingerprinting environmental conditions and related stress using stable isotopic composition of rice (<i>Oryza sativa</i> L.) grain organic matter R. Kaushal, P. Ghosh (Bangalore, India) and H. Geilmann (Jena, Germany).....	941
Toward a practical use of Neotropical odonates as bioindicators: Testing congruence across taxonomic resolution and life stages F. Valente-Neto, F. de Oliveira Roque, M.E. Rodrigues (Campo Grande, Brazil), L. Juen (Belém, Brazil) and C.M. Swan (Baltimore, MD, USA)	952
Forest carbon storage along the north-south transect of eastern China: Spatial patterns, allocation, and influencing factors D. Wen and N. He (Beijing, China).....	960
Are red wood ants (<i>Formica rufa</i> -group) tectonic indicators? A statistical approach G. Berberich (Essen, Germany), A. Grumpe (Dortmund, Germany), M. Berberich (Erfstadt, Germany), D. Klimetzek (Freiburg, Germany) and C. Wöhler (Dortmund, Germany).....	968
Understanding agricultural virtual water flows in the world from an economic perspective: A long term study R. Duarte, V. Pinilla (Zaragoza, Spain) and A. Serrano (Girona, Spain).....	980
Environmental determinants of chironomid communities in remote northern lakes across the treeline – Implications for climate change assessments T.P. Luoto (Helsinki, Finland and Jyväskylä, Finland), M.V. Rantala, A. Galkin (Helsinki, Finland), M. Rautio (Québec, Canada) and L. Nevalainen (Jyväskylä, Finland)	991
A hydrologically sensitive invertebrate community index for New Zealand rivers M.J. Greenwood, D.J. Booker (Christchurch, New Zealand), B.J. Smith (Hamilton, New Zealand) and M.J. Winterbourn (Christchurch, New Zealand)	1000
Modelling the spatial distribution of the seagrass <i>Posidonia oceanica</i> along the North African coast: Implications for the assessment of Good Environmental Status M. Zucchetta, C. Venier (Italy), M.A. Taji (Morocco and Casablanca, Morocco), A. Mangin (Sophia-Antipolis, France) and R. Pastres (Italy)	1011
The spatial correlation and interaction between manufacturing agglomeration and environmental pollution Z. Cheng (Nanjing, China and Taian, China)	1024
Taxonomic distinctness along nutrient gradients: More diverse, less diverse or not different from random? A. Vilmi, S.M. Karjalainen, M. Kuoppala, K.T. Tolonen and J. Heino (Oulu, Finland).....	1033
Mapping biodiversity in three-dimensions challenges marine conservation strategies: The example of coralligenous assemblages in North-Western Mediterranean Sea A. Doxa (Montpellier France), F. Holon, J. Deter (Montpellier France and Carnon, France), S. Villéger (Montpellier, France), P. Boissery (Marseille, France) and N. Mouquet (Montpellier France).....	1042
Corrigendum to “Application of fuzzy VIKOR for evaluation of green supply chain management practices” [Ecol. Indic. 49 (2015) 188–203] R. Rostamzadeh (Urmia, Iran), K. Govindan (Odense, Denmark), A. Esmaili (Tehran, Iran) and M. Sabaghi (Quebec, Canada).....	1055