BRITISH POULTRY ABSTRACTS

September 2015 CONTENTS Volume 11, Number 1

2015 SPRING MEETING OF THE WPSA UK BRANCH

Oral communications, invited talks and posters presented at the WPSA UK Branch Meeting held at the University of Chester (14–15 April 2015). These summaries have been edited for clarity and style by the WPSA UK Programme Committee but have not been fully peer-reviewed.

| M. Kasprzak, G. White, S. Kightley, J. Houdijk, O. Olukosi, P. Carre and J. Wiseman. Amino acid digestibility in rapeseed co-products varies between processing conditions and varieties. | 1 |
|--|----|
| J. Abdulla, V. Pirgozliev, S. P. Rose and A. M. MacKenzie. Feeding quality of field beans for broiler chickens. | 2 |
| V. Pirgozliev, W. Mirza and S. P. Rose. Hard wheat instead of soft wheat in pelleted diets results in improved feed efficiency in broiler chickens. | 3 |
| G. Parkes, M. W. Mirza, S. P. Rose and V. Pirgozliev. The influence of feed form and steam pelleting of diets on the growth performance and digestive tract development of broilers. | 5 |
| M. A. Mitchell. Metabolic disease in poultry – shared mechanisms of cardiovascular and muscle pathologies? | 6 |
| C. D. Parker. Metabolic disease in poultry – prevalence and industry concerns. | 7 |
| B. Duggan, D. Clements and P. M. Hocking. Skeletal leg morphology – adaptation to high growth rate in chickens and ducks. | 8 |
| D. N. R. G. Kapell, AM. Neeteson, J. McAdam, M. S. Swalander, A. N. M. Koerhuis and S. Avendaño. The genetic basis of walking ability in purebred broiler lines. | 9 |
| A. M. A. Reid, P. M. Hocking, P. W. Wilson, G. W. Robertson and I. C. Dunn. Understanding genetic selection for growth: the case of satiety in meat chickens. | 10 |
| J. E. Martin, D. E. F. McKeegan, S. Brocklehurst and V. Sandilands. Novel analysis of electroencephalography (EEG) data of on-farm-killed poultry. | 12 |
| K. Morrissey, T. Widowski, L. Baker and V. Sandilands. Effects of dietary protein source, fibre level and extra environmental enrichments on feather condition and behaviour. | 13 |
| F. Shariatmadari, H. Shirzadi, M. A. Karimi Torshizi and S. Rahimi. Effect of phenolic Prosopis farcta and Rhus coriaria L. extracts on broiler chicken performance. | 14 |
| E. Chatzidimitriou, S. Stergiadis, M. D. Eyre, M. Bestman, T. van Niekerk, and G. Butler. Effect of management practices on fatty acid composition of chicken eggs: a farm survey in The Netherlands. | 16 |
| O. O. Adeleye, S. S. Abiola, M. O. Sogunle, L. T. Egbeyale, D. A. Ekunseitan, M. A. Adeyemi, O. Oduwole and O. O. Oduguwa. Growth performance, Eimeria oocyst and worm count (Ascaris galli) of broiler chickens fed Lagenaria breviflora. | 17 |
| W. Mirza, V. Pirgozliev, P. M. Rose and M. Bedford. Interaction of exogenous phytase and xylanase on nutrient availability for broilers. | 18 |
| C. O. Sanni, E. J. Burton and J. Boyd. Bone mineralisation in broilers: evaluating the methodology for bone ash determination. | 19 |

| N. Samat and H. M. Miller. Effects of anticoagulants on the measurement of circulating myo-inositol (MYO), calcium (Ca) and phosphorus (P) in broiler chickens fed a phytase-supplemented wheat-maize-soya-based diet. | 21 |
|--|----|
| H. D. Chapman and T. K. Jeffers. Vaccination of broiler chickens against coccidiosis may ameliorate drug resistance in Eimeria and lead to long-term sustainable control of this disease in poultry production. | 23 |
| A. Sinclair, C. Weber Wyneken, T. Veldkamp, L. J. Vinco and P. M. Hocking. Behavioural assessment of pain in turkeys with foot pad dermatitis. | 24 |
| N. Corcionivoschi, L. Moran, C. Kelly, P. Scates and R. Madden. Isolation and virulence characteristics of Campylobacter coli and Campylobacter jejuni hcp ⁺ isolates from retail chicken produced in Northern Ireland. | 25 |
| A. L. Smith. Animal science, politics and policy - a New Zealand perspective. | 27 |
| M. Howell. Methods to control the level of Campylobacter during poultry processing. | 27 |
| I. Connerton and P. L. Connerton. Bacteriophage control of Campylobacter in broiler chickens. | 28 |
| I. M. Whiting, V. Pirgozliev, S. P. Rose, A. M. Mackenzie and A. M. Amerah. The effect of xylanase supplementation on digestible energy and nutrient digestibility of diets containing wheat DDGS when fed to laying hens. | 29 |
| S. C. Pace, M. R. Bedford and H. M. Miller. The effect of betaine and phytase on egg quality over a 4-week storage period. | 30 |
| J. L. Harmse, J. C. Engelbrecht and J. L. Bekker. Can occupational health practices in poultry abattoirs impact the quality of poultry meat? A South African perspective. | 32 |
| P. M. Hocking, T. Veldkamp, L. J. Vinco and P. Woodward. High water intake is associated with soya compared with non-soya protein sources and may be associated with foot pad dermatitis in growing turkeys. | 33 |
| C. Combemorel, O. A. Olukosi, S. Kightley, J. Wiseman and J. G. M. Houdijk. True ileal phosphorus digestibility of oilseed rape meal for broiler determined by the regression method. | 34 |
| A. Tosti, K. Morrissey and V. Sandilands. Effects of diets containing animal protein and extra fibre on eggshell quality in laying hens. | 36 |
| C. Wang, V. Sandilands, H. Yang, Q. Zhou, G. Cheng, T. Miao, D. Fu and C. Han. The effects of different low-temperature exposure methods on the behaviour of commercial broilers. | 37 |
| T. Goossens, M. Marien, A. De Cesare and G. Manfreda. Precision-delivery-coated butyrate to control Campylobacter in poultry. | 38 |
| I. Al-Homidan, M. M. Fathi and A. Al-Shumaymiri. The effect of litter type on cell-mediated response and serum IgM concentration of broiler chicks. | 39 |
| I. M. Whiting, V. R. Pirgozliev, S. P. Rose, A. M. Mackenzie and A. M. Amerah. Mineral availability of wheat distillers dried grains with solubles (DDGS) for laying hens. | 40 |
| I. M. Whiting, V. R. Pirgozliev, S. P. Rose, A. M. Mackenzie and A. M. Amerah. Exogenous xylanase improves dietary nutrient availability in diets containing wheat DDGS for laying hens. | 42 |
| W. Mirza, V. Pirgozliev, S. P. Rose, H. Graham, H. Masey O'Neill, M. Bedford, E. Ball, J. George and J. Sloss. The effect of quality characteristics of wheat on litter quality and hock burn when fed to broiler chickens. | 43 |
| C. Lückstädt, L. Lorenz and S. Mellor. Effect of dietary sodium diformate on the growth performance of broilers till 35 d post-hatch. | 44 |

F. Khattak, M. A. Hossain, Z. Hayat, K. Morrissey and V. Sandilands. Effect of storage time, temperature and conditions on egg quality and composition of commercial non-beak-trimmed laying hens fed diets differing in protein and fibre.

45

R. Plimmer, N. Morgan, D. Scholey and E. Burton. Effect of three different iron supplement forms on breast and liver iron levels and tibia strength.

47