

ORIGINAL CONTRIBUTIONS**J. A. SANCHEZ, F. M. DEL AMOR, P. FLORES, E. LÓPEZ-GALLEGO**Nutritional variations at *Nesidiocoris tenuis* feeding sites and reciprocal interactions between the mirid and tomato plants

161–173

J. A. OTIENO, P. PALLMANN, H.-M. POEHLING

The combined effect of soil-applied azadirachtin with entomopathogens for integrated management of western flower thrips

174–186

K. B. NEETHU, P. PRIJI, K. N. UNNI, S. SAJITH, S. SREEDEVI, N. RAMANI, K. ANITHA, B. ROSANA, M. B. GIRISH, S. BENJAMINNew *Bacillus thuringiensis* strain isolated from the gut of Malabari goat is effective against *Tetranychus macfarlanei*

187–198

A. FERRER, A. F. G. DIXON, J.-L. HEMPTINNE

Habitat or prey specialization in predatory ladybird beetles: a case study of two closely related species

199–208

G. MOLLOT, N. BOROWIEC, P.-F. DUYCK, S. GLÉNAC, S. QUILICILife-history traits of *Encarsia guadeloupa*, a natural enemy of the invasive spiralling whitefly *Aleurodicus dispersus*

209–217

M. ST-ONGE, D. CORMIER, S. TODOROVA, É. LUCASConservation of *Ephestia kuehniella* eggs as hosts for *Trichogramma ostrinae*

218–222

SHORT COMMUNICATIONS**U. AZIZOGLU, A. AYVAZ, S. YILMAZ, R. TEMIZGUL**

The synergic and antagonistic activity of Cry1Ab and Cry2Aa proteins against lepidopteran pests

223–227

S. K. JACOBSEN, I. ALEXAKIS, L. SIGSGAARDAntipredator responses in *Tetranychus urticae* differ with predator specialization

228–231

J. M. MALYSH, I. A. KAZARTSEV, A. N. FROLOV, À. A. ZVEREV, Y. S. TOKAREVMolecular detection of *Cotesia vestalis* (Hymenoptera: Braconidae) in the beet webworm *Loxostege sticticalis* L. (Lepidoptera: Crambidae)

232–235

A. BAYRAM, A. TONĀFirst report of *Chilo partellus* in Turkey, a new invasive maize pest for Europe

236–240