

CONTENTS

Articles

VANG, L.E., OPPERMAN, C.H., SCHWARZ, M.R. & DAVIS, E.L.— Spirotetramat causes an arrest of nematode juvenile development	121-131
KANZAKI, N., AIKAWA, T., MAEHARA, N. & THU, P.Q.— <i>Bursaphelenchus kesiyae</i> n. sp. (Nematoda: Aphelenchoididae), isolated from dead wood of <i>Pinus kesiya</i> Royle ex Gordon (Pinaceae) from Vietnam, with proposal of new subgroups in the <i>B. fungivorus</i> group	133-146
SAPKOTA, R., SKANTAR, A.M. & NICOLAISEN, M.— A TaqMan real-time PCR assay for detection of <i>Meloidogyne hapla</i> in root galls and in soil	147-154
QING, X., SÁNCHEZ-MONGE, A., JANSSEN, T., COUVREUR, M. & BERT, W.— Description of <i>Malenchus sexlineatus</i> n. sp., new records of three known species of <i>Malenchus</i> Andrassy, 1968 (Nematoda: Tylenchidae) and notes on amphidial aperture development	155-174
TANDINGAN DE LEY, I., HOLOVACHOV, O., MC DONNELL, R.J., BERT, W., PAINE, T.D. & DE LEY, P.— Description of <i>Phasmarhabditis californica</i> n. sp. and first report of <i>P. papillosa</i> (Nematoda: Rhabditidae) from invasive slugs in the USA	175-193
KANZAKI, N., KAJIMURA, H. & OKABE, K.— <i>Rhabditoides aegus</i> n. sp., an associate of stag beetles and sap flow in Japan	195-208
KANZAKI, N., MAEHARA, N., AKIBA, M., TANAKA, S.E. & IDE, T.— Morphological characters of dauer juveniles of three species of <i>Bursaphelenchus</i> Fuchs, 1937	209-220
KANZAKI, N. & IDE, T.— <i>Diplogasteroides luxuriosae</i> n. sp. associated with <i>Acalolepta luxuriosa</i> (Coleoptera: Cerambycidae) from Japan	221-233
ABATE, B.A., MALAN, A.P., TIEDT, L.R., WINGFIELD, M.J., SLIPPERS, B. & HURLEY, B.P.— <i>Steinernema fabii</i> n. sp. (Rhabditida: Steinernematidae), a new entomopathogenic nematode from South Africa	235-255