

ORIGINAL CONTRIBUTIONS

- J. C. DO PRADO, M. F. G. V. PEÑAFLOR, E. CIA, S. S. VIEIRA, K. I. SILVA, L. A. CARLINI-GARCIA, A. L. LOURENÇÃO**
Resistance of cotton genotypes with different leaf colour and trichome density to *Bemisia tabaci* biotype B 405–413
- N. A. TINSLEY, P. D. MITCHELL, R. J. WRIGHT, L. J. MEINKE, R. E. ESTES, M. E. GRAY**
Estimation of efficacy functions for products used to manage corn rootworm larval injury 414–425
- P. A. UMINA, M. SCHIFFER, P. PARKER, A. A. HOFFMANN**
Distribution and influence of grazing on wheat curl mites (*Aceria tosichella* Keifer) within a wheat field 426–433
- G. BOZSIK, A. TRÖGER, W. FRANCKE, G. SZŐCS**
Thuja occidentalis: identification of volatiles and electroantennographic response by the invasive cedar bark beetle, *Phloeosinus aubei* 434–443
- P. R. C. DODDALA, M. A. MINOR, Q. WANG, D. J. ROGERS, E. M. KOOT, S. A. TREWICK**
Role of olfaction in host plant selection and local adaptation of a polyphagous herbivore, *Eucolaspis* Sharp 444–452
- S. ONDIKA, L. MIGIRO, M. RUR, G. BIRGERSSON, M. PORCEL, B. RÄMERT, M. TASIN**
Sunflower as a trap crop for the European tarnished plant bug (*Lygus rugulipennis*) 453–461
- S. KRUEGER, L. A. MOUND, G. B. MORITZ**
Offspring sex ratio and development are determined by copulation activity in *Echinothrips americanus* MORGAN 1913 (Thysanoptera: Thripidae) 462–473
- SHORT COMMUNICATIONS**
- A. N. COSTA, E. H. M. VIEIRA-NETO**
Species turnover regulates leaf-cutter ant densities in environmental gradients across the Brazilian Cerrado 474–478
- Y.-D. VARENNES, M. GONZALEZ CHANG, S. BOYER, S. D. WRATTEN**
Nectar feeding increases exploratory behaviour in the aphid parasitoid *Diaeretiella rapae* (McIntosh) 479–483