

ORIGINAL CONTRIBUTIONS**V. H. P. BUENO, A. M. CALIXTO, F. C. MONTES, J. C. VAN LENTEREN**Reproduction and population parameters of the Nearctic predator *Geocoris punctipes* at constant and varying temperature regimes 323–333**S. SAMRA, M. GHANIM, A. PROTASOV, Z. MENDEL**Comparative study of development parameters of four *Ooencyrtus* spp. (Hymenoptera: Encyrtidae) on natural and factitious hosts 334–345**X.-S. WANG, Q.-Z. CHEN, S.-Z. ZHANG, T.-X. LIU**Parasitism, host feeding and immature development of *Encarsia formosa* reared from *Trialeurodes vaporariorum* and *Bemisia tabaci* on *Trialeurodes ricini* 346–352**L. H. WU, A. A. HOFFMANN, L. J. THOMSON***Trichogramma* parasitoids for control of Lepidopteran borers in Taiwan: species, life-history traits and *Wolbachia* infections 353–363**S. MUNIR, L. M. DOSDALL, J. T. O'DONOVAN, A. KEDDIE***Diadegma insulare* development is altered by *Plutella xylostella* reared on water-stressed host plants 364–375**M. GHANINIA, M. AMOOGHLI TABARI**

Olfactory cues explain differential attraction of the striped rice stem borer to different varieties of rice plant 376–385

A. R. W. ELBERS, R. MEISWINKELLimited attractant range of the black-light suction trap for the capture of *Culicoides* biting midges (Diptera: Ceratopogonidae) 386–394**Š. GRUCMANOVÁ, J. HOLUŠA, V. ČERMÁK, J. NERMUŤ**Nematodes associated with *Ips cembrae* (Coleoptera: Curculionidae): comparison of generations, sexes and sampling methods 395–403