

## CONTENTS

**Volume 28 Number 2 2016**

- Survival of juvenile northern quahogs during seasonal temperature decline likely a function of diet and NMI-fatty acid synthesis  
*Sixto E. Portilla and Brett F. Branco* 53
- Tubifex production using agro-industrial wastes and raw cattle dung  
*R. N. Mandal, S. Kar, D. N. Chattopadhyay, J. Maity, B. N. Paul, P. P. Chakrabarti, and P. Jayasankar* 70
- Feed technology and production performance of Thai koi (*Anabas testudeneus*) in Bangladesh  
*Md. Al Amin Sarker, Fatimah Mohamed Arshad, Md. Ferdous Alam, Zainal Abidin Mohamed, and Md. Akhtaruzzaman Khan* 76
- Preclinical toxicity and safety evaluation of *Althaea officinalis* L. extract as naturopathic medicine in common carp (*Cyprinus carpio* L.): Hematological and biochemical study  
*Mahdi Banaee, Fahimeh Fallahpour, Vahid Soleimany, and Behzad Nematdoost Haghi* 92
- Growth pattern and biometric relationship of pearl oyster *Pinctada maxima* cultured in Beibu Bay, China  
*Feilong Liang, Shaohe Xie, Shao Fu, Junhui Li, and Yuewen Deng* 110
- Effects of omega-3 fatty acid supplementation on aquaponics system-raised tilapia (*Oreochromis niloticus* X *Oreochromis aureus*) growth, physiology, and immunology  
*Tiffany Hough, Christina Glaze, Elliott Blumenthal, and Ahmed Mustafa* 119