

- 54 **Calculation of threshold Olsen P values for fertilizer response from soil properties**  
R. Recena · I. Díaz · M.C. del Campillo · J. Torrent · A. Delgado
- 55 **Alternative use of tobacco as a sustainable crop for seed oil, biofuel, and biomass**  
S. Grisan · R. Polizzotto · P. Raiola · S. Cristiani · F. Ventura · F. di Lucia · M. Zuin · S. Tommasini · R. Morbidelli · F. Damiani · F. Pupilli · M. Bellucci
- 56 **Soiless system on peat reduce trace metals in urban-grown food: unexpected evidence for a soil origin of plant contamination**  
G. Pennisi · F. Orsini · D. Gasperi · S. Mancarella · R. Sanoubar · L. Vittori Antisari · G. Vianello · G. Gianquinto
- 57 **Life cycle assessment of edible insects for food protein: a review**  
A. Halloran · N. Roos · J. Eilenberg · A. Cerutti · S. Bruun
- 58 **Belowground nitrogen transfer from legumes to non-legumes under managed herbaceous cropping systems. A review**  
M.S. Thilakarathna · M.S. McElroy · T. Chapagain · Y.A. Papadopoulos · M.N. Raizada
- 59 **Viruses carried to soil by irrigation can be aerosolized later during windy spells**  
G. Girardin · P. Renault · F. Bon · Ł. Capowicz · J. Chadoeuf · C. Krawczyk · D. Courault
- 60 **Hyper-proteolytic mutant of *Beauveria bassiana*, a new biological control agent against the tomato borer**  
I. Borgi · J.-W. Dupuy · I. Blibech · D. Lapaillerie · A.-M. Lomenech · A. Rebai · M. Ksantini · M. Bonneau · A. Gargouri
- 61 **Weedy fallow as an alternative strategy for reducing nitrogen loss from annual cropping systems**  
S.E. Wortman
- 62 **Chestnut wood extract in boar diet reduces intestinal skatole production, a boar taint compound**  
D. Bilić-Šobot · G. Zamaratskaia · M.K. Rasmussen · M. Čandek-Potokar · M. Škrlep · M.P. Povše · D. Škorjanc
- 63 **Unexpectedly, intense livestock grazing in arid rangelands strengthens the seedling vigor of consumed plants**  
M. Tadey · C.P. Souto
- 64 **Processes of adaptation in farm decision-making models. A review**  
M. Robert · A. Thomas · J.-E. Bergez
- 65 **Erratum to: Belowground nitrogen transfer from legumes to non-legumes under managed herbaceous cropping systems. A review**  
M.S. Thilakarathna · M.S. McElroy · T. Chapagain · Y.A. Papadopoulos · M.N. Raizada
- 66 **Breeding for efficiency in the broiler chicken: A review**  
C.W. Tallentire · I. Leinonen · I. Kyriazakis
- 67 **Object-based early monitoring of a grass weed in a grass crop using high resolution UAV imagery**  
F. López-Granados · J. Torres-Sánchez · A.-I. De Castro · A. Serrano-Pérez · F.-J. Mesas-Carrascosa · J. Peña
- 68 **A resistant pepper used as a trap cover crop in vegetable production strongly decreases root-knot nematode infestation in soil**  
M. Navarrete · C. Djian-Caporalino · T. Mateille · A. Palloix · A.-M. Sage-Palloix · A. Lefèvre · A. Fazari · N. Marteu · J. Tavoillot · A. Dufils · C. Furnion · L. Pares · I. Forest
- 69 **Farming tactics to reduce the carbon footprint of crop cultivation in semiarid areas. A review**  
C. Liu · H. Cutforth · Q. Chai · Y. Gan
- 70 **Cacao agroforestry systems have higher return on labor compared to full-sun monocultures**  
L. Armengot · P. Barbieri · C. Andres · J. Milz · M. Schneider