

INDUSTRIAL CROPS AND PRODUCTS

VOLUME 109

15 December 2017

Contents

Abstracted/indexed in: Agricola; CAB Abstracts; CABS; Chemical Abstracts; Current Contents; AB & ES; EMBASE; Embiology Elsevier BIOBASE/Current Awareness in Biological Sciences; Elsevier/Geobase; Environmental Abstracts. Also covered in the abstract and citation database SCOPUS®. Full text available on ScienceDirect®

Fats and oils

- Change in some biochemical and bioactive properties and essential oil composition of coriander seed (*Coriandrum sativum* L.) varieties from Turkey
E. Beyzi, K. Karaman, A. Gunes and S. Buyukkilic Beyzi (Turkey) 74
- Deterpenation of *Origanum majorana* L. essential oil by reduced pressure steam distillation
G. Ben Salha (Tunisia, Spain), R. Herrera Díaz, J. Labidi (Spain) and M. Abderrabba (Tunisia) 116
- Characterization of *Xanthoceras sorbifolium* Bunge seeds: Lipids, proteins and saponins content
M. Venegas-Calación, M.V. Ruíz-Méndez, E. Martínez-Force, R. Garcés and J.J. Salas (Spain) 192
- Preparation and optimization nanoemulsion of Tarragon (*Artemisia dracunculus*) essential oil as effective herbal larvicide against *Anopheles stephensi*
M. Osanloo, A. Amani, H. Sereshti, M.R. Abai, F. Esmaeili and M.M. Sedaghat (Iran) 214
- Optimized nitrogen fertilizer application improves yield, water and nitrogen use efficiencies of winter rapeseed cultivated under continuous ridges with film mulching
X.-B. Gu, Y.-N. Li and Y.-D. Du (China) 233
- Towards improving the agronomic performance, chlorophyll fluorescence parameters and pigments in fenugreek using zeolite and vermicompost under deficit water stress
A. Baghbani-Arani, S.A.M. Modarres-Sanavy, M. Mashhadi-Akbar-Boojar and A. Mokhtassi-Bidgoli (Iran) 346
- Dodartia orientalis* L. essential oil exerts antibacterial activity by mechanisms of disrupting cell structure and resisting biofilm
F. Wang, F. Wei, C. Song, B. Jiang, S. Tian, J. Yi, C. Yu, Z. Song, L. Sun, Y. Bao, Y. Wu, Y. Huang and Y. Li (PR China) 358
- Preparation of fully biobased epoxy resins from soybean oil based amine hardeners
C.F. Frias, A.C. Serra, A. Ramalho, J.F.J. Coelho and A.C. Fonseca (Portugal) 434
- Antimicrobial and antioxidant activity of Juniper galbuli essential oil constituents eluted at different times
V.D. Zheljzakov (USA), I.B. Semerdjieva, I. Dincheva (Bulgaria), M. Kacaniova (Slovak Republic, Poland), T. Astatkie (Canada), T. Radoukova (Bulgaria) and V. Schlegel (USA) 529

(Contents continued on inside back cover)

- Dry-season deficit irrigation increases agricultural water use efficiency at the expense of yield and agronomic nutrient use efficiency of Sacha Inchi plants in a tropical humid monsoon area
Y.J. Geng, L. Chen, C. Yang, D.Y. Jiao, Y.H. Zhang and Z.Q. Cai (China) 570
- Foliar boron fertilization as factor affecting the essential oil content and yield of oil components from flower heads of *Arnica montana* L. and *Arnica chamissonis* Less. cultivated for industry
D. Sugier, P. Sugier, R. Kowalski, B. Kołodziej and K. Olesińska (Poland) 587
- Morphological and biochemical intraspecific characterization of *Ocimum basilicum* L.
F. Varga, K. Carović-Stanko (Croatia), M. Ristić (Serbia), M. Grdiša, Z. Liber and Z. Šatović (Croatia) 611
- Characterization of *Jatropha curcas* accessions based in plant growth traits and oil quality
N. Albuquerque, R.C. García-Almodóvar, J.M. Valverde, L. Burgos and D. Martínez-Romero (Spain) 693
- Characterization of *Perilla frutescens* (Linn.) Britt based on morphological, biochemical and STMS markers
S.K. Singh, P.C. Kole, A.K. Misra, S. Roy, L. Arya, M. Verma, R. Bhardwaj, P. Suneja, M.R. Verma, K.V. Bhat and R. Singh (India) 773
- Biomass production and essential oil in a new bred cultivar of peppermint (*Mentha × piperita* L.)
J. Fejér, D. Gruřová (Slovak Republic) and V. De Feo (Italy) 812
- Non-food bioactive products**
- Sophora flavescens* Seed as a promising high potential by-product: Phytochemical characterization and bioactivity evaluation
Z. Weng, S. Guo, D. Qian, Z. Zhu, S. Zhang, A. Li, Z. Lei and J. Duan (China) 19
- Agronomic and phytochemical evaluation of lavandin and lavender cultivars cultivated in the Tyrrhenian area of Tuscany (Italy)
L. Pistelli, B. Najar, S. Giovanelli, L. Lorenzini, S. Tavarini and L.G. Angelini (Italy) 37
- Essential oil content and composition of *Vitex pseudo-negundo* in Iran varies with ecotype and plant organ
T. Movahhed Haghighi, M.J. Saharkhiz, A.R. Khosravi, F. Raouf Fard and M. Moein (Iran) 53
- Composition and bioactivity assessment of essential oils of *Curcuma longa* L. collected in China
L. Zhang, Z. Yang, F. Chen, P. Su, D. Chen, W. Pan, Y. Fang (China), C. Dong (France), X. Zheng (USA) and Z. Du (China) 60
- In vitro* antimicrobial, antioxidant and antiviral activities of the essential oil and various extracts of wild (*Daucus virgatus* (Poir.) Maire) from Tunisia
A. Snene, R. El Mokni, H. Jmii, I. Jlassi, H. Jaidane (Tunisia), D. Falconieri, A. Piras (Italy), H. Dhaouadi (Tunisia), S. Porcedda (Italy) and S. Hammami (Tunisia) 109
- Dynamics of flower, achene and trichome development governs the accumulation of pyrethrins in pyrethrum (*Tanacetum cinerariifolium*) under irrigated and dryland conditions
D.D. Suraweera, T. Groom, P.W.J. Taylor, C.S. Jayasinghe and M.E. Nicolas (Australia) 123
- Marigold extract, azelaic acid and black caraway oil into lipid nanocarriers provides a strong anti-inflammatory effect *in vivo*
I. Lacatusu, G. Badea, M. Popescu, N. Bordei, D. Istrati, L. Moldovan, A.M. Seciu, M.I. Panteli, I. Rasit and N. Badea (Romania) 141
- Potential use of Negramina (*Siparuna guianensis* Aubl.) essential oil to control wax moths and its selectivity in relation to honey bees
T.P. Ferreira, E.E. Oliveira, P.H. Tschoeke, R.G. Pinheiro, A.M.S. Maia and R.W.S. Aguiar (Brazil) 151
- Isobutyrylshikonin and isovalerylshikonin from the roots of *Onosma visianii* inhibit larval growth of the tobacco cutworm *Spodoptera littoralis*
S. Sut (Italy), R. Pavela (Czech Republic), V. Kolarčik (Slovakia), G. Lupidi, F. Maggi, S. Dall'Acqua and G. Benelli (Italy) 266
- Antioxidant, antityrosinase and antibiofilm activities of synthesized peptides derived from *Vicia faba* protein hydrolysate: A powerful agents in cosmetic application
I. Karkouch, O. Tabbene, D. Gharbi (Tunisia), M.A. Ben Mlouka (France), S. Elkahoui (Tunisia), C. Rihouey, L. Coquet, P. Cosette, T. Jouenne (France) and F. Limam (Tunisia) 310
- Corymbia* spp. and *Eucalyptus* spp. essential oils have insecticidal activity against *Plutella xylostella*
C.A. Filomeno, L.C.A. Barbosa, R.R. Teixeira, A.L. Pinheiro, E. de Sá Farias, E.M. de Paula Silva and M.C. Picanço (Brazil) 374

- Effect of drying methods on the physico-chemical properties and adhesion performance of water-washed cottonseed meal
N. Li, S. Prodyawong, Z. He, X.S. Sun and D. Wang (USA) 281
- Volatiles of roots of wild-growing and cultivated *Armoracia macrocarpa* and their antimicrobial activity, in comparison to horseradish, *A. rusticana*
S. Petrović, M. Drobac, L. Ušjak, V. Filipović, M. Milenković and M. Niketić (Serbia) 398
- Lignin as a green reductant and morphology directing agent in the fabrication of 3D graphene-based composites for high-performance supercapacitors
W. Ye, X. Li, J. Luo, X. Wang and R. Sun (China) 410
- Highly efficient liquefaction of wheat straw for the production of bio-polyols and bio-based polyurethane foams
H. Li (China, Canada), S. Feng, Z. Yuan (Canada), Q. Wei (China) and C.C. Xu (Canada) 426
- Influence of degree of substitution on thermal dynamic mechanical and physicochemical properties of cellulose acetate
R.R.M. de Freitas, A.M. Senna and V.R. Botaro (Brazil) 452
- Chemical characterization of proanthocyanidins in purple, blue, and red maize coproducts from different milling processes and their anti-inflammatory properties
C. Chen, P. Somavat, V. Singh and E. Gonzalez de Mejia (United States) 464
- Organic foliar fertilization increases polyphenol content of *Calendula officinalis* L.
V. Onofrej, G.-C. Teliban, M. Burducea, A. Lobiuc, C.B. Sandu, M. Tocai and T. Robu (Romania) 509
- Polyethylene composites made from below-ground miscanthus biomass
L. Chupin, S. Arnoult, M. Brancourt-Hulmel, C. Lapiere, E. Gineau and P. Navard (France) 523
- Immobilization of white rot fungi to carbohydrate-rich corn cob as a basis for tertiary treatment of secondarily treated pulp and paper mill wastewater
A. Zhang, G. Wang, G. Gong and J. Shen (China) 538
- Starch content affects physicochemical properties of corn and cassava starch-based films
C.L. Luchese, J.C. Spada and I.C. Tessaro (Brazil) 619
- Mechanical and barrier properties of spray dried carboxymethyl cellulose (CMC) film from bleached bagasse pulp
S. Kamthai and R. Magaraphan (Thailand) 753
- Fabricating 3D hierarchical porous TiO₂ and SiO₂ with high specific surface area by using nanofibril-interconnected cellulose aerogel as a new biotemplate
X. Zhang, S. Jing, Z. Chen, L. Zhong, Q. Liu, X. Peng and R. Sun (China) 790
- Active biodegradable cassava starch films incorporated lycopene nanocapsules
R.Q. Assis, S.M. Lopes, T.M.H. Costa, S.H. Flôres and A.d.O. Rios (Brazil) 818
- All natural cork composites with suberin-based polyester and lignocellulosic residue
H. De Oliveira, B. Yoon (South Korea), V. Michaud (Switzerland), J.-D. Nam and J. Suhr (South Korea) 843
- The isolation of cellulose nanocrystals from pistachio shells via acid hydrolysis
J. Marett, A. Aning and E.J. Foster (USA) 869
- Properties and structure of poly(3-hydroxybutyrate-co-4-hydroxybutyrate)/wood fiber biodegradable composites modified with maleic anhydride
S. An and X. Ma (China) 882

Potential for ethanol production from different sorghum cultivars <i>E. Castro (United States, Spain), I.U. Nieves, V. Rondón, W.J. Sagues, M.T. Fernández-Sandoval (United States), L.P. Yomano, S.W. York, J. Erickson (USA) and W. Vermerris (USA, United States)</i>	367	Tracking lipid profiles of <i>Jatropha curcas</i> L. seeds under different pruning types and water managements by low-field and HR-MAS NMR spectroscopy <i>O.N.A. Santos, M.V. Folegatti, L.M. Dutra, I.P.d.S. Andrade, E.D. Fanaya Jr., B.P. Lena, A. Barison and A.D.d.C. Santos (Brazil)</i>	918
Comparison of alkaline and acid pretreatments for enzymatic hydrolysis of soybean hull and soybean straw to produce fermentable sugars <i>Q. Qing, Q. Guo, L. Zhou, X. Gao, X. Lu and Y. Zhang (China)</i>	391	Bio-based Materials from Crops	
Potential antioxidants for biodiesel from a softwood lignin pyrolyzate <i>R.A. Larson, B.K. Sharma, K.A. Marley, B. Kunwar, D. Murali and J. Scott (United States)</i>	476	Methanolysis of epoxidized soybean oil in continuous flow conditions <i>V. Pantone, A.G. Laurenza, C. Annese, F. Fracassi, C. Fusco, A. Nacci, A. Russo and L. D'Accolti (Italy)</i>	1
Assessing multi-scale deconstruction of wood cell wall subjected to mechanical milling for enhancing enzymatic hydrolysis <i>J. Jiang, J. Wang, X. Zhang and M. Wolcott (USA)</i>	498	<i>Musa balbisiana</i> Colla peel as highly effective renewable heterogeneous base catalyst for biodiesel production <i>M. Gohain, A. Devi and D. Deka (India)</i>	8
A risk averse cross-efficiency data envelopment analysis model for sustainable switchgrass cultivation location optimization <i>H. Khanjarpanah, M.S. Pishvaei and S.M. Seyedhosseini (Iran)</i>	514	Self-reinforcing composites from commercial chemical pulps via partial dissolution with NaOH/urea <i>N.C. Hildebrandt (Germany, Finland), P. Piltonen (Finland), J.-P. Valkama (Germany) and M. Illikainen (Finland)</i>	79
Modelling the hydrophilic extraction of the bark of <i>Eucalyptus nitens</i> and <i>Eucalyptus globulus</i> : Adsorption isotherm and thermodynamic studies <i>M.S. Parada and K. Fernández (Chile)</i>	558	Preparation of biomass-based activated carbons and their evaluation for biogas upgrading purposes <i>P.C. Vilella, J.A. Lira, D.C.S. Azevedo, M. Bastos-Neto and R. Stefanutti (Brazil)</i>	134
Redefinition of sweet sorghum harvest time: New approach for sampling and decision-making in field <i>T.P.M. Teixeira, L.D. Pimentel, L.A.d.S. Dias, R.A.d.C. Parrella, M.Q. da Paixão and E.M. Biesdorf (Brazil)</i>	579	Ionic liquid pretreatment of bagasse improves mechanical property of bagasse/polypropylene composites <i>K. Ninomiya, M. Abe, T. Tsukegi, K. Kuroda, M. Omichi, K. Takada, M. Noguchi, Y. Tsuge, C. Ogino, K. Taki, T. Taima, K. Uzawa and K. Takahashi (Japan)</i>	158
Structural and compositional changes in eucalyptus wood chips subjected to dry torrefaction <i>C.M.S. da Silva, B.R. Vital, A.d.C.O. Carneiro, E.V.S. Costa, M.A. de Magalhães and P.F. Trugilho (Brazil)</i>	598	Radiation-induced modifications in natural fibres and their biocomposites: Opportunities for controlled physico-chemical modification pathways? <i>N. Le Moigne, R. Sonnier, R. El Hage and S. Rouif (France)</i>	199
Antioxidant and anticancer potential of bioactive compounds following UV-C light-induced plant cambium meristematic cell cultures <i>S.H. Moon, B. Mistry, D.H. Kim and M. Pandurangan (Republic of Korea)</i>	762	Effects of preparation methods on the morphology and properties of nanocellulose (NC) extracted from corn husk <i>X. Yang, F. Han, C. Xu, S. Jiang, L. Huang, L. Liu and Z. Xia (PR China)</i>	241
Industrial clarification of wheat-based distillers' solubles and thin stillage <i>K. Ratanapariyanuch (Canada), Y.Y. Shim (Canada, China), S. Emami (Canada) and M.J.T. Reaney (Canada, China)</i>	828	Boll sampling protocols and their impact on measurements of cotton fiber quality <i>N. Kothari, S. Hague, L. Hinze and J. Dever (United States)</i>	248
Bending properties of loblolly pine <i>O. Oyedeji, A. Young and O. Fasina (United States)</i>	905	The features of emulsion copolymerization for plant oil-based vinyl monomers and styrene <i>K. Kingsley (United States), O. Shevchuk (Ukraine), Z. Demchuk (United States), S. Voronov (Ukraine) and A. Voronov (United States)</i>	274

Biorefinery, valorization of byproducts		Bioenergy - crops and processing	
Integrated forest biorefineries: Recovery of acetic acid as a by-product from eucalyptus wood hemicellulosic hydrolysates by solvent extraction <i>F. Cebreiros, M.D. Guigou and M.N. Cabrera (Uruguay)</i>	101	Investigating the effects of grinding time and grinding load on content of terpenes in extract from fennel obtained by supercritical fluid extraction <i>T. Hatami (Brazil, Iran), J.C.F. Johner and M.A.A. Meireles (Brazil)</i>	85
Enhancing enzymatic hydrolysis of corn cob, corn stover and sorghum stalk by dilute aqueous ammonia combined with ultrasonic pretreatment <i>Q.-Q. Xu, M.-J. Zhao, Z.-Z. Yu, J.-Z. Yin, G.-M. Li, M.-Y. Zhen and Q.-Z. Zhang (China)</i>	220	A new lab scale corn dry milling protocol generating commercial sized flaking grits for quick estimation of coproduct yield and composition <i>P. Somavat, Q. Li, D. Kumar, E.G. de Mejia, W. Liu, K.D. Rausch, J.A. Juvik, M.E. Tumbleson and V. Singh (USA)</i>	92
Structural functionalization of industrial softwood kraft lignin for simple dip-coating of urea as highly efficient nitrogen fertilizer <i>J. Li, M. Wang (Sweden), D. She (Sweden, China) and Y. Zhao (Sweden)</i>	255	Effects of reduced severity ammonia pretreatment on pelleted corn stover <i>N. Nahar and S.W. Pryor (USA)</i>	163
Thermo-mechano-chemical extraction of hydroxycinnamic acids from industrial hemp by-products using a twin-screw extruder <i>L. Candy, S. Bassil, L. Rigal, V. Simon and C. Raynaud (France)</i>	335	Synergistic action of an <i>Aspergillus</i> (hemi-) cellulolytic consortium on sugarcane bagasse saccharification <i>P. de Oliveira Rodrigues, J. de Cássia Pereira, D.Q. dos Santos, L.V.A. Gurgel, D. Pasquini and M.A. Baffi (Brazil)</i>	173
Extraction of nanolignin from coconut fibers by controlled microbial hydrolysis <i>S.J. Juikar and N. Vigneshwaran (India)</i>	420	Study of the effects of extraction methods on Argan oil quality through its metal content <i>F.E. Ennoukh, R. Bchitou (Morocco), F. Mohammed (Morocco, Yemen), D. Guillaume (France), H. Harhar and A. Bouhaouss (Morocco)</i>	182
Enzymatic activity and protein expression of cellulase from rice straw produced by <i>Trichoderma reesei</i> in the presence of oxygen vectors <i>J. Xia, A. He, R. Li, Y. Zhang, J. Xu, X. Liu and J. Xu (China)</i>	654	Utilization of agricultural residues for enhancement of cellulolytic enzyme production and enzymatic saccharification by <i>Trichoderma harzianum</i> KUC1716 <i>H. Lee, Y.M. Lee, Y.M. Heo, J. Lee, J.S. Kim, K.Y. Kang and J.-J. Kim (Republic of Korea)</i>	185
Extraction of polyphenols from <i>Eucalyptus nitens</i> and <i>Eucalyptus globulus</i> : Experimental kinetics, modeling and evaluation of their antioxidant and antifungal activities <i>N. González, J. Elissetche, M. Pereira and K. Fernández (Chile)</i>	737	Chemical composition of five <i>Miscanthus sinensis</i> harvests and nitric-acid cellulose therefrom <i>Y.A. Gismatulina and V.V. Budaeva (Russia)</i>	227
Applicability of fruit blanching and intermittent microwave-convective belt drying to industrial peel waste of different mango cultivars for the recovery of functional coproducts <i>A. Nagel, S. Neidhart, S. Kuebler (née Wulfkuehler), P. Elstner, T. Anders, S. Korhummel, T. Sulzer, S. Kienzle, C. Winkler (Germany), S. Qadri (Thailand), C. Rentschler (Germany), N. Pholpipattanapong, J. Wuthisomboon (Thailand), H.-U. Endress (Germany), P. Sruamsiri (Thailand) and R. Carle (Germany, Saudi Arabia)</i>	923	Methane and hydrogen production from anaerobic digestion of soluble fraction obtained by sugarcane bagasse ozonation <i>O.F.H. Adarme, B.E.L. Baêta, D.R.S. Lima, L.V.A. Gurgel and S.F. de Aquino (Brazil)</i>	288
		Ratoon cold tolerance of <i>Pennisetum</i> , <i>Erianthus</i> , and <i>Saccharum</i> bioenergy feedstocks <i>D.M. Burner, A.L. Hale, R.P. Viator, D.P. Belesky, J.H. Houx III, A.J. Ashworth and F.B. Fritschi (United States)</i>	327

- Effect of post-pruning vine-shoots storage on the evolution of high-value compounds
C. Cebrián, R. Sánchez-Gómez, M.R. Salinas, G.L. Alonso and A. Zalacain (Spain) 730
- Chemical composition, antioxidant and evidence antimicrobial synergistic effects of *Periploca laevigata* essential oil with conventional antibiotics
L. Ait Dra, M. Ait Sidi Brahim, B. Boualy, A. Aghraz, M. Barakate, S. Oubaassine, M. Markouk and M. Larhsini (Morocco) 746
- A tetracosatetraene as larvicidal compound isolated from *Alpinia katsumadai*
K.S. Hwang, Y.K. Kim, Y.T. Kim, J. Lee and K.W. Park (Republic of Korea) 786
- Verification of the antioxidant activity of a subterranean part of *Suaeda japonica* Makino
K.-Y. Kang, Y.-H. Hwang, S.-J. Lee (Republic of Korea), J.-J. Kim (Singapore), S.-J. Nam and S.-T. Yee (Republic of Korea) 836
- Monoterpene chiral ratios: Chemotype diversity and interspecific commonality in *Melaleuca alternifolia* and *M. linariifolia*
I. Southwell, A. Dowell, S. Morrow, G. Allen, D. Savins and M. Shepherd (Australia) 850
- Antimicrobial kinetic activities of lignin from sugarcane bagasse for textile product
J. Sunthornvarabhas, S. Liengprayoon and T. Suwonsichon (Thailand) 857
- Chemical composition, antimicrobial and anti-quorum sensing activities of pummelo peel flavonoid extract
Z. Liu, Y. Pan, X. Li, J. Jie and M. Zeng (China) 862
- Summer savory extracts prepared by novel extraction methods resulted in enhanced biological activity
P. Mašković, V. Veličković, M. Mitić, S. Đurović, Z. Zeković, M. Radojković, A. Cvetanović, J. Švarc-Gajić and J. Vujić (Serbia) 875
- Phytochemical evaluation of major bioactive compounds in different cytotypes of five species of *Rumex* L.
S.M. Jeelani, U. Farooq, A.P. Gupta and S.K. Lattoo (India) 897
- The essential oil of *Melaleuca armillaris* (Sol. ex Gaertn.) Sm. leaves from Pakistan: A potential source of eugenol methyl ether
S. Siddique, Z. Parveen, Firdaus-e-Bareen, M.N. Chaudhary, S. Mazhar and S. Nawaz (Pakistan) 912
- Natural rubber and resins**
- A century of guayule: Comprehensive genetic characterization of the US national guayule (*Parthenium argentatum* A. Gray) germplasm collection
D.C. Ilut, P.L. Sanchez, T.A. Coffelt, J.M. Dyer, M.A. Jenks and M.A. Gore (USA) 300
- Mesostructure characterization by asymmetrical flow field-flow fractionation of natural rubber samples from different *Hevea brasiliensis* genotypes
S. Liengprayoon (Thailand), K. Chelbi, S. Dubascoux, C. Char, L. Vaysse, E. Dubreucq, J. Sainte Beuve (France), K. Sriroth (Thailand) and F. Bonfils (France) 936
- Fibres, proteins and carbohydrates**
- Evaluation of the incorporation of lignocellulose nanofibrils as sustainable adhesive replacement in medium density fiberboards
C.I.K. Diop, M. Tajvidi, Michael.A. Bilodeau, D.W. Bousfield and J.F. Hunt (USA) 27
- Fiber properties of axis and scale of eleven different coniferous cones
S.K. Gülsoy, H. Hafizoğlu, A. Kılıç Pekgözlü, İ. Tümen, İ.E. Dönmez and H. Sivrikaya (Turkey) 45
- Green and efficient PEG-based ultrasound-assisted extraction of polysaccharides from superfine ground lotus plumule to investigate their antioxidant activities
L. Zhang, Z. Cheng, Q. Zhao and M. Wang (PR China) 320
- Improving lignocellulose enzymatic saccharification in a bioreactor with an applied electric field
Y.-Z. Wang, L. Zhang, T. Xu and K. Ding (China) 404
- Facilitate hemicelluloses separation from chemical pulp in ionic liquid/water by xylanase pretreatment
X. Ma (China, Canada), Y. Long, C. Duan (Canada), X. Lin, S. Cao, L. Chen, L. Huang (China) and Y. Ni (China, Canada) 459
- How to reduce cotton fiber damage in the Xinjiang China
J. Tian, X. Zhang, Y. Yang, C. Yang, S. Xu, W. Zuo, W. Zhang, H. Dong, X. Jiu, Y. Yu and Z. Zhao (China) 803
- Preparation of 3D printable micro/nanocellulose-poly(lactic acid) (MNC/PLA) composite wire rods with high MNC constitution
Z. Wang, J. Xu, Y. Lu, L. Hu, Y. Fan, J. Ma and X. Zhou (China) 889

(Contents continued from inside back cover)

- Effects of thermo-vacuum treatment on secondary metabolite content and antioxidant activity of poplar (*Populus nigra* L.) wood extracts
L. Todaro, D. Russo, P. Cetera and L. Milella (Italy) 384
- Phenological changes in triterpenic and phenolic composition of *Thymus* L. species
L. Raudone, K. Zymone, R. Raudonis, R. Vainoriene, V. Motiekaityte and V. Janulis (Lithuania) 445
- X-ray structure of *O*-methyl-acrocol and anti-cancer, anti-parasitic, anti-bacterial and anti-Zika virus evaluations of the Brazilian palm tree *Acrocomia totai*
G.K. Souza, I.T. Albrecht Schuquel, V.M. Moura, A.C. Belloto, L.U. Rovigatti Chiavelli, A.L.T.G. Ruiz, L. Shiozawa, J.E. de Carvalho, F.P. Garcia, V. Kaplum, J.H. da Silva Rodrigues, D.B. Scariot, R. Delvecchio, E. Machado-Ferreira, R. Santana, C.A. Gomes Soares, C.V. Nakamura, S.M. de Oliveira Santin and A.M. Pomini (Brazil) 483
- Curcumin bioactive nanosizing: Increase of bioavailability
L.C. Massimino, H.A.M. Faria and S.A. Yoshioka (Brazil) 493
- Preconditions for industrial use of foliage as felling by-product of Scots pine for essential oil production
J. Labokas, K. Ložienė and R. Jurevičiūtė (Lithuania) 542
- Controlled enzymatic hydrolysis for improved exploitation of the antioxidant potential of wheat gluten
M.B. Elmalimadi, J.R. Jovanović, A.B. Stefanović, S.J. Tanasković, S.B. Djurović, B.M. Bugarski and Z.D. Knežević-Jugović (Serbia) 548
- Efficacy of sea fennel (*Crithmum maritimum* L., Apiaceae) essential oils against *Culex quinquefasciatus* Say and *Spodoptera littoralis* (Boisd.)
R. Pavela (Czech Republic), F. Maggi, G. Lupidi (Italy), K. Cianfaglione (France, Italy), X. Dauvergne (France), M. Bruno and G. Benelli (Italy) 603
- Growth, photosynthetic pigments, phenolic content and biological activities of *Foeniculum vulgare* Mill., *Anethum graveolens* L. and *Pimpinella anisum* L. (Apiaceae) in response to zinc
N. Majdoub (Tunisia, Portugal), S. el-Guendouz (Portugal), M. Rezgui (Tunisia), J. Carlier, C. Costa (Portugal), L.B.B. Kaab (Tunisia) and M.G. Miguel (Portugal) 627
- A combined process is used for efficient isolation and purification of xylobiose from xylanase-hydrolysed sugarcane bagasse xylan hydrolysate
S.-X. Jiang, S. Zhao, C.-Y. Lu, J.-L. Xue, C.-J. Duan and J.-X. Feng (People's Republic of China) 637
- Influence of water deficit on the main polysaccharides and the rheological properties of Aloe vera (*Aloe barbadensis* Miller) mucilage
R. Minjares-Fuentes (Spain, Mexico), L. Medina-Torres, R.F. González-Laredo, V.M. Rodríguez-González (Mexico), V. Eim and A. Femenia (Spain) 644
- The Apiaceae: Ethnomedicinal family as source for industrial uses
B. Sayed-Ahmad (France, Lebanon), T. Talou (France), Z. Saad, A. Hijazi (Lebanon) and O. Merah (France) 661
- Optimization of supercritical fluid extraction of steviol glycosides and total phenolic content from *Stevia rebaudiana* (Bertoni) leaves using response surface methodology and artificial neural network modeling
K. Ameer, B.-S. Chun and J.-H. Kwon (Republic of Korea) 672
- Potential botanical pesticides from Asteraceae essential oils for tomato production: Activity against whiteflies, plants and bees
M.L. Umpiérrez, J. Paullier (Uruguay), M. Porrini, M. Garrido (Argentina), E. Santos and C. Rossini (Uruguay) 686
- Cynara scolymus* L.: A promising Mediterranean extract for topical anti-aging prevention
P. Marques, J. Marto, L.M. Gonçalves, R. Pacheco, M. Fitas, P. Pinto, M.L.M. Serralheiro and H. Ribeiro (Portugal) 699
- Phytochemicals and anti-aging potentials of the extracts from *Lagerstroemia speciosa* and *Lagerstroemia floribunda*
P. Kolakul and B. Sripanidkulchai (Thailand) 707
- The bioaccessible phenolic profile and antioxidant potential of *Hypericum perforatum* L. after simulated human digestion
E. Celep, Y. İnan, S. Akyüz and E. Yesilada (Turkey) 717
- Trypanocide, antileishmania and cytotoxic activities of the essential oil from *Rosmarinus officinalis* L. *in vitro*
P.S. Pereira (Brazil), A.J. Maia, S.R. Tintino (Paraguay), C.D.d.M. Oliveira-Tintino (Brazil), I.S.d.S. Raulino, M.C. Vega, M. Rolón, C. Coronel (Paraguay), L.M. Barros, A.E. Duarte, I.R.A. de Menezes, H.D.M. Coutinho and T.G. da Silva (Brazil) 724