

## RESEARCH PAPERS

- Differential nitrogen supply causes large variability in photosynthetic traits in wheat germplasm  
*Szilvia Veres, Al Imran Malik, Zed Rengel* 703
- Elevated field atmospheric CO<sub>2</sub> concentrations affect the characteristics of winter wheat (cv. Bologna) grains  
*Francesca Verrillo, Franz-Werner Badeck, Valeria Terzi, Fulvia Rizza, Letizia Bernardo, Antimo Di Maro, Clara Fares, Alessandro Zaldei, Francesco Miglietta, Anna Moschella, Marcella Bracale, Candida Vannini* 713
- Alternate furrow irrigation affects yield and water-use efficiency of maize under deficit irrigation  
*Farid Golzardi, Amirsaleh Baghdadi, Reza Keshavarz Afshar* 726
- Biosolids differently affect seed yield, nodule growth, nodule-specific activity, and symbiotic nitrogen fixation of field bean  
*S. Pampana, A. Scartazza, R. Cardelli, A. Saviozzi, L. Guglielminetti, G. Vannacci, M. Mariotti, A. Masoni, I. Arduini* 735
- Can legumes provide greater benefits than millet as a spring cover crop in southern Queensland farming systems?  
*E. M. Wunsch, L. W. Bell, M. J. Bell* 746
- Allometric relationships for vegetative growth flush in pruned stems of macadamia (*Macadamia integrifolia*): effects of carbohydrate source and season  
*S. Karimaei, D. Doley, J. Hanan* 760
- An assessment of weed flora 14 years after the introduction of glyphosate-tolerant cotton in Australia  
*Sudheesh Manalil, Jeff Werth, Rod Jackson, Bhagirath Singh Chauhan, Christopher Preston* 773
- Persistence and productivity of phalaris (*Phalaris aquatica*) germplasm in dry marginal rainfall environments of south-eastern Australia  
*R. A. Culvenor, M. R. Norton, J. De Faveri* 781
- Biological nitrification inhibition by weeds: wild radish, brome grass, wild oats and annual ryegrass decrease nitrification rates in their rhizospheres  
*Cathryn A. O'Sullivan, Kelley Whisson, Karen Treble, Margaret M. Roper, Shayne F. Micin, Philip R. Ward* 798
- Corrigendum 805