Ecological Research

The Ecological Society of Japan

Volume 32 • Number 6 • November 2017

Special Feature on Filling the Gaps

■ Data Paper

Fukasawa K, Mishima Y, Yoshioka A, Kumada N, Totsu K Acoustic monitoring data of avian species inside and outside the evacuation zone of the Fukushima Daiichi power plant accident 769

■ Oshima Award

Han Q, Kabeya D

Recent developments in understanding mast seeding in relation to dynamics of carbon and nitrogen resources in temperate trees 771

■ Special Feature: Filling the Gaps Editorial

Nakamura M, Hsieh C, Miki T, Kondoh M

Establishment of an ecological research network involving Taiwan and Japan: developing a better understanding of ecological phenomena unique to East Asia 779

■ Special Feature: Data Paper

Iwayama A, Ogura H, Hirama Y, Chang C-W, Hsieh C, Kagami M Phytoplankton species abundance in Lake Inba (Japan) from 1986 to 2016 783

■ Special Feature: Biwako Prize for Ecology

Chang C-W, Ushio M, Hsieh C

Empirical dynamic modeling for beginners 785

■ Special Feature: Original Articles

Liu W-C, Lai S-M, Chen H-W

A topological similarity-based bootstrapping method for inferring food web parameters 797

Nakadai R

Species diversity of herbivorous insects: a brief review to bridge the gap between theories focusing on the generation and maintenance of diversity 811

Weng C-Y, Yang K-C, Hsieh C-F, Hsieh C, Su M-H

Local neighborhood communities in the understory play a critical role by affecting regeneration niches and subsequent community assembly in a montane cloud forest 821

Laplace S, Komatsu H, Tseng H, Kume T

Difference between the transpiration rates of Moso bamboo (*Phyllostachys pubescens*) and Japanese cedar (*Cryptomeria japonica*) forests in a subtropical climate in Taiwan 835

Lin M-Y, Hsieh I-F, Lin P-H, Laplace S, Ohashi M, Chen T-H, Kume T

Moso bamboo (*Phyllostachys pubescens*) forests as a significant carbon sink? A case study based on 4-year measurements in central Taiwan 845

Erratum 859

Itoh M, Kojima H, Ho P-C, Chang C-W, Chen T-Y, Hsiao SS-Y, Kobayashi Y, Fujibayashi M, Kao S-J, Hsieh C, Fukui M, Okuda N, Miki T, Shiah F-K

Integrating isotopic, microbial, and modeling approaches to understand methane dynamics in a frequently disturbed deep reservoir in Taiwan 861

■ Special Feature: Perspective Paper

Chen I-C, Hsieh C, Kondoh M, Lin H-J, Miki T, Nakamura M, Ohgushi T, Urabe J, Yoshida T

Filling the gaps in ecological studies of socioecological systems 873

■ Original Articles

Luo Y, Zhao X, Zuo X, Li Y, Wang T

Plant responses to warming and increased precipitation in three categories of dune stabilization in northeastern China 887

Aadrean, Usio N

Small-clawed otters (*Aonyx cinereus*) in Indonesian rice fields: latrine site characteristics and visitation frequency 899

N'Dri JK, Seka FA, Pokou PK, N'Da RAG, Lagerlöf J

Abundance and diversity of soil mite (Acari) communities after conversion of tropical secondary forest into rubber plantations in Grand-Lahou, Côte d'Ivoire 909

Contents continued on inside back cover

Stenert C, Wüsth R, Pires MM, Freiry RF, Nielsen D, Maltchik L

Composition of cladoceran dormant stages in intermittent ponds with different hydroperiod lengths 921

Haigh A, O'Riordan R, Butler F

Variations in aggression and activity levels amongst squirrels inhabiting low and high density areas 931

Hattori A

Aerial images can detect 3D small patch reefs that are potential habitats for anemonefish Amphiprion frenatus 943

Fantinato E, Del Vecchio S, Baltieri M, Fabris B, Buffa G

Are food-deceptive orchid species really functionally specialized for pollinators? 951

Yao B, Liu O, Gao Y, Cao Z

Characterizing vertical migration of *Microcystis aeruginosa* and conditions for algal bloom development based on a light-driven migration model 961

Nhiwatiwa T, Maseko Z, Dalu T

Fish communities in small subtropical reservoirs subject to extensive drawdowns, with focus on the biology of *Enteromius paludinosus* (Peters, 1852) and *Clarias gariepinus* (Burchell, 1822) 971

Zani D, Müller JV

Climatic control of seed longevity of Silene during the post-zygotic phase: do seeds from warm, dry climates possess higher maturity and desiccation tolerance than seeds from cold, wet climates? 983

Sanchooli N

Ecological variables determining the presence of lizards in the Sistan region, Eastern Iran 995

Wang Z, Han G, Hao X, Zhao M, Ding H, Li Z, Wang J, Hamilton A, Liu Y, Lata A, Hexige B

Effect of manipulating animal stocking rate on the carbon storage capacity in a degraded desert steppe 1001

Zhang M, Chen F, Chen S, Xie Z, Huang Y, Liu Y

The soil seed bank of a rehabilitated draw-down zone and its similarity to standing vegetation in the Three Gorges Reservoir Area 1011

Chikorowondo G, Muvengwi J, Mbiba M, Gandiwa E

Influence of abandoned cattle enclosures on plant assemblages and herbivory in a semi-arid savanna 1023

■ Acknowledgment

Acknowledgment 1035