

| | |
|--|---------|
| Monette, D., Ewe, S., Dinkins, J. M. & Markwith, S. H.: Interactions of exotic and native <i>Pomacea</i> with wetland vegetation structure in the Greater Everglades, Florida, USA | 291–299 |
| Ágoston-Szabó, E., Schöll, K., Kiss, A. & Dinka, M.: The effects of tree species richness and composition on leaf litter decomposition in a Danube oxbow lake (Gemenc, Hungary). | 301–314 |
| Gartzia de Bikuña, B., Arrate, J., Martínez, A., Agirre, A., Azpiroz, I. & Manzanos, A.: Development of a fish-based index to assess the ecological status of oceanic-temperate streams in the Northern Iberian Peninsula | 315–327 |
| Shuai, F., Li, X., Chen, F., Li, Y. & Lek, S.: Spatial patterns of fish assemblages in the Pearl River, China: environmental correlates | 329–340 |
| Sharifian, S., Kamrani, E., Safaie, M. & Sharifian, S.: Population structure and growth of freshwater crab <i>Sodhiana iranica</i> from the south of Iran | 341–349 |
| Yatigamma, S. & Cumming, B. F.: Detecting recent limnological changes in two ancient reservoirs from Sri Lanka’s Dry Zone using Cladocera | 351–363 |
| Erratum | 365 |