

DISEASES

Marker-free transgenic rice lines with a defensin gene are potentially active against phyto-pathogenic fungus <i>Sarocladium oryzae</i> <i>M. Pérez-Bernal, M. Delgado, A. Cruz, D. Abreu, O. Valdivia and R. Armas</i>	135
Effects of fly ash amendments, <i>Ralstonia solanacearum</i> , <i>Meloidogyne incognita</i> and <i>Phomopsis vexans</i> on the growth of <i>Solanum melongena</i> <i>M. Khan and Z. A. Siddiqui</i>	145
Isolation and characterization of bacterial agents associated of wetwood disease on elm trees in Iran <i>M. Alizadeh, R. Khakvar and A. Babai-Ahari</i>	157
Survey of the aflatoxin gene cluster in <i>Aspergilli</i> from Hungarian crops <i>Szilvia Kovács and Tünde Pusztahelyi</i>	169
Evaluation of antagonistic activity of <i>Trichoderma</i> spp. against <i>Meloidogyne incognita</i> <i>S. Sellami, N. Benttoui, S. Berrahia and H. Boureghda</i>	177
The role of silicon (Si) in increasing plant resistance against insect pests <i>N. Sakr</i>	185

Ladybirds (Coleoptera: Coccinellidae) of Iran: 45 New records from 10 provinces <i>R. Abdolahi, O. Nedvěd and J. Nozari</i>	205
Chemical control of the Lesser Date Moth, <i>Batrachedra amydraula</i> (Lepidoptera: Batrachedridae) <i>M. Javadzadeh and A. Hosseini-Gharalari</i>	215
A key to genera of Iranian lady beetles (Coleoptera: Coccinellidae) and species of subfamilies Chilocorinae, Coccinellinae, Epilachinae and Microweiseinae <i>R. Abdolahi, O. Nedvěd and J. Nozari</i>	219
Daily flight activity rhythms of the peach and mediterranean fruit flies using sexual and olfactory attractants <i>M. H. Bayoumy and M. M. El-Metwally</i>	227
Taxonomic study on the subfamily Curculioninae (Col., Curculionidae) in East Azarbaijan province with one species as new record for Iran <i>R. Rahati, J. Nozari and S. Khaghaninia</i>	245
Acute and persistence effects of oil of <i>Hippophae rhamnoides</i> and <i>Calendula officinalis</i> on <i>Sitophilus granarius</i> (Coleoptera: Curculionidae) in stored maize <i>S. Keszthelyi, R. Hoffmann, Zs. Pónya and F. Pál-Fám</i>	255
Effect of grain type on the insecticidal efficacy of Paya® against <i>Tribolium castaneum</i> (Herbst) (Coleoptera: Tenebrionidae) <i>M. G. Fazlabad, M. A. Badihi and M. Z. Khormizi</i>	265
New host records of parasitoids of scale insects and whitefly species in Egypt <i>S. Abd-Rabou and G. A. Evans</i>	275