

PERSPECTIVES PAPERS

Efficacy and risks of “biorationals”

F. Feldmann · C. Carstensen 517

“Biorationals” in integrated pest management strategies

E. Matyjaszczyk 523

ORIGINAL ARTICLES

Are synthetic volatiles, typically emitted by insect-damaged peach cultivars, navigation signals for two-spotted lady beetle (*Adalia bipunctata* L.) and green lacewing (*Chrysoperla carnea* [Stephens]) larvae?

Ž. Laznik · S. Trdan 529

Efficiency of funnel traps in controlling pine processionary moth

S. Parlak · İ.M. Özçankaya · M. Batur · M.E. Akkaş · Z. Boza · Ö. Toprak 539

Effectiveness of leaf rust resistance genes against *Puccinia triticina* populations in Western Siberia during 2008–2017

E.S. Skolotneva · I.N. Leonova · E.Y. Bukatich · N.I. Boiko · V.V. Piskarev · E.A. Salina 549

Effects of insecticidal seed treatments and foliar sprays in winter oilseed rape in autumn on insect pests and TuYV infection

N. Conrad · M. Brandes · T. Will · J.-A. Verreet · B. Ulber · U. Heimbach 557

Plant parasitic nematodes on soybean in expanding production areas of temperate regions

A. Elhady · H. Heuer · J. Hallmann 567

SHORT COMMUNICATIONS

Reaction of wild eggplant *Solanum torvum* to different species of root-knot nematodes from Turkey

S. Öçal · T. Özalp · Z. Devran 577

Tubercle disease of sugar beet roots (*Beta vulgaris*) found in Poland is neither caused by *Xanthomonas beticola* nor by tumorigenic *Agrobacterium/Rhizobium*

E. Moliszewska · M. Nabrdalik · J. Puławska · J. Piszczek 581

Toxicity and antifungal activity of 2, 3-dimethylmaleic anhydride against stored product beetles and plant fungal pathogens

T.B. Devi · S. Mayanglambam · H.D. Chanu · K.D. Singh · S. Sougrakpam · B. Nameirakpam · N.I. Singh · M. Potshangbam · D. Sahoo · Y. Rajashekar 585