

CONTENTS

| | | | |
|---|----|--|-----|
| Advantages of combining generalized linear models and occupancy models to find indicators of habitat selection: Small mammals in agroecosystems as a case study I.L. Gorosito, M.M. Bermúdez and M. Busch (Argentina) | | Vulnerability assessment of spring wheat production to climate change in the Inner Mongolia region of China Z. Dong, Z. Pan, Q. He, J. Wang, L. Huang, Y. Pan, G. Han, X. Xue and Y. Chen (China) | 67 |
| The establishment of season-specific eutrophication assessment standards for a water-supply reservoir located in Northeast China based on chlorophyll-a levels J. Guo, C. Zhang, G. Zheng, J. Xue and L. Zhang (China) | 1 | Shift of low to high nucleic acid bacteria as a potential bioindicator for the screening of anthropogenic effects in a receiving river due to palm oil mill effluent final discharge S.S. Sharuddin, N. Ramli (Malaysia), D. Mohd-Nor (Malaysia, Japan), M.A. Hassan (Malaysia), T. Maeda, Y. Shirai, K. Sakai and Y. Tashiro (Japan) | 79 |
| Long-term land use and land cover changes (1920–2015) in Eastern Ghats, India: Pattern of dynamics and challenges in plant species conservation R.M. Ramachandran, P.S. Roy, V. Chakravarthi, J. Sanjay and P.K. Joshi (India) | 11 | Biological indicators of diversity in tropical streams: Congruence in the similarity of invertebrate assemblages J.G. Brito, R.T. Martins, V.C. Oliveira, N. Hamada, J.L. Nessimian (Brazil), R.M. Hughes (United States), S.F.B. Ferraz and F.R. de Paula (Brazil) | 85 |
| First assessment of the ecological status in the Levant Basin: Application of the CARLIT index along the Lebanese coastline A. Badreddine (Lebanon, France), M.A.-A. Saab (Lebanon), F. Gianni (France), E. Ballesteros (Spain) and L. Mangialajo (France) | 21 | Inherent vulnerability assessment of rural households based on socio-economic indicators using categorical principal component analysis: A case study of Kimsar region, Uttarakhand S. Rajesh, S. Jain and P. Sharma (India) | 93 |
| An indicator-based approach to analyse the effects of non-native tree species on multiple cultural ecosystem services A.S. Vaz (Portugal), P. Castro-Díez, O. Godoy, Á. Alonso, M. Vilà, A. Saldaña (Spain), H. Marchante (Portugal), Á. Bayón (Spain), J.S. Silva, J.R. Vicente and J.P. Honrado (Portugal) | 37 | Mapping landscape potential for outdoor recreation using different archetypical recreation user groups in the European Union F. Komossa, E.H. van der Zanden, C.J.E. Schulp (The Netherlands) and P.H. Verburg (The Netherlands, Switzerland) | 105 |
| Linking water quality with the total pollutant load control management for nitrogen in Jiaozhou Bay, China K. Li, J. He, J. Li, Q. Guo, S. Liang, Y. Li and X. Wang (China) | 48 | Suitability of different landscape metrics for the assessments of patchy landscapes in West Africa J.N. Inkoom, S. Frank, K. Greve, U. Walz and C. Fürst (Germany) | 117 |
| | 57 | | |

| | | | |
|---|-----|---|-----|
| The challenge of habitat modelling for threatened low density species using heterogeneous data: The case of Cuvier's beaked whales in the Mediterranean | | Monte Carlo simulation based interval chance-constrained programming for regional ecosystem management – A case study of Zhuhai, China | |
| A. Cañadas (Spain), N.A. de Soto (Spain, UK), M. Aissi (Tunisia), A. Arcangeli, M. Azzolin, A. B-Nagy, G. Bearzi, I. Campana (Italy), C. Chicote (Spain), C. Cotte (France), R. Crosti (Italy), L. David (France), A. Di Natale (Spain), C. Fortuna (Italy, Croatia), A. Frantzis (Greece), P. Garcia, M. Gazo, R. Gutierrez-Xarxa (Spain), D. Holcer (Croatia), S. Laran (France), G. Lauriano (Italy), T. Lewis (UK), A. Moulins, B. Mussi, G.N. di Sciara, S. Panigada (Italy), X. Pastor (Spain), E. Politi, M. Pulcini (Italy), J.A. Raga (Spain), L. Rendell (UK), M. Rosso, P. Tepsich (Italy), J. Tomás (Spain), M. Tringali (Italy) and Th. Roger (France) | 128 | J.P. Lv (China), Y.P. Li (China, Canada) and J. Sun (China) | 214 |
| Cross-taxon congruence and relationships to stand characteristics of vascular plants, bryophytes, polyporous fungi and beetles in mature managed boreal forests | | Urbanization is associated with elevated corticosterone in Jollyville Plateau salamanders | |
| J. Jokela, K. Juutilainen, L. Korpela, J. Kouki, S. Kuntsi, M. Koivula and J. Siitonen (Finland) | 137 | C.R. Gabor, D.R. Davis, D.S. Kim, K.C. Zabierek and N.F. Bendik (USA) | 229 |
| Visual versus video methods for estimating reef fish biomass | | The water consumed in the production of grapes for vinification (<i>Vitis vinifera</i>). Mapping the blue and green water footprint | |
| S.K. Wilson (Australia), N.A.J. Graham (UK), T.H. Holmes (Australia), M.A. MacNeil (Australia, Canada) and N.M. Ryan (Australia) | 146 | B. Civit, R. Piastrellini, S. Curadelli and A.P. Arena (Argentina) | 236 |
| Metabarcoding of benthic ciliate communities shows high potential for environmental monitoring in salmon aquaculture | | Simple family-level parrotfish indicators are robust to survey method | |
| T. Stoeck, R. Kochems, D. Forster (Germany), F. Lejzerowicz and J. Pawlowski (Switzerland) | 153 | H. Vallès and H.A. Oxenford (Barbados) | 244 |
| Culturomic assessment of Brazilian protected areas: Exploring a novel index of protected area visibility | | Species of <i>Ulva</i> (Ulvophyceae, Chlorophyta) as indicators of salinity | |
| R.A. Correia (Brazil, United Kingdom), P. Jepson (United Kingdom), A.C.M. Malhado (Brazil) and R.J. Ladle (Brazil, United Kingdom) | 165 | A.S. Rybak (Poland) | 253 |
| Topographic wetness index explains soil moisture better than bioindication with Ellenberg's indicator values | | Trophic gradient is the main determinant of species and large taxonomic groups representation in phytoplankton of standing water bodies | |
| M.W. Raduła, T.H. Szymura and M. Szymura (Poland) | 172 | O. Lepšová-Skáčelová, P. Fibich, J. Wild and J. Lepš (Czech Republic) | 262 |
| Fish species as indicators of freshwater inflow within a subtropical estuary in the Gulf of Mexico | | The role of urban function on road soil respiration responses | |
| J.L. Steichen and A. Quigg (United States) | 180 | H. Ye, C. Sun, K. Wang, G. Zhang, T. Lin and H. Yan (China) | 271 |
| Urban land use intensity assessment: The potential of spatio-temporal spectral traits with remote sensing | | Efficiency estimation of urban metabolism via Emergy, DEA of time-series | |
| T. Wellmann, D. Haase, S. Knapp, C. Salbach, P. Selsam and A. Lausch (Germany) | 190 | Y. Wu, W. Que, Y.-g. Liu, J. Li, L. Cao, S.-b. Liu, G.-m. Zeng and J. Zhang (PR China) | 276 |
| Development of a microbial community-based index of biotic integrity (MC-IBI) for the assessment of ecological status of rivers in the Taihu Basin, China | | Landscape heterogeneity as a surrogate of biodiversity in mountain systems: What is the most appropriate spatial analytical unit? | |
| L. Niu, Y. Li, P. Wang, W. Zhang, C. Wang, J. Li and H. Wu (PR China) | 204 | P. García-Llamas, L. Calvo, M. De la Cruz and S. Suárez-Seoane (Spain) | 285 |
| | | Vulnerability to sea level rise: A novel local-scale indicator-based assessment methodology and application to eight beaches in Shoalhaven, Australia | |
| | | F.N. Tonmoy and A. El-Zein (Australia) | 295 |
| | | Evaluation of coral reef management effectiveness using conventional versus resilience-based metrics | |
| | | A.K. Ford, A. Eich, R.S. McAndrews (Germany), S. Mangubhai (Fiji), M.M. Nugues (France, French Polynesia), S. Bejarano (Germany), B.R. Moore (New Caledonia), C. Rico (Fiji, Spain), C. Wild and S.C.A. Ferse (Germany) | 308 |
| | | Fatty acids profiles modifications in the bivalves <i>Cerastoderma edule</i> and <i>Scrobicularia plana</i> in response to copper sulphate | |
| | | A.F. Mesquita, F. Gonçalves, T. Verdelhos, J.C. Marques and A.M.M. Gonçalves (Portugal) | 318 |

| | | | |
|---|-----|--|-----|
| Cascading effects between climate, vegetation, and macroinvertebrate fauna in 14,000-year palaeoecological investigations of a shallow lake in eastern Poland M. Słowiński, P. Skubała, I. Zawiska, A. Kruk, M. Obremska, K. Milecka (Poland) and F. Ott (Germany) | | Can high-throughput sequencing detect macroinvertebrate diversity for routine monitoring of an urban river? M.E. Carew, C.R. Kellar, V.J. Pettigrove and A.A. Hoffmann (Australia) | 440 |
| Taxonomic and functional traits responses of <i>Sphagnum</i> peatland testate amoebae to experimentally manipulated water table I. Koenig (Switzerland), M. Mulot (Switzerland, France) and E.A.D. Mitchell (Switzerland) | 329 | Sustainability impacts of tidal river management: Towards a conceptual framework Md.M.A. Masud, N.N. Moni (Bangladesh), H. Azadi and S. Van Passel (Belgium) | 451 |
| Scenario analysis of the carbon pricing policy in China's power sector through 2050: Based on an improved CGE model Y. Zhao, H. Li, Y. Xiao, Y. Liu, Y. Cao, Z. Zhang, S. Wang, Y. Zhang and A. Ahmad (China) | 342 | Synoptic scale mammal density index map based on roadkill records T. Tatewaki and F. Koike (Japan) | 468 |
| Predicting stem diameters and aboveground biomass of individual trees using remote sensing data M. Dalponte, L. Frizzera (Italy), H.O. Ørka, T. Gobakken, E. Næsset (Norway) and D. Gianelle (Italy) | 352 | Method for evaluating ecological vulnerability under climate change based on remote sensing: A case study L. Jiang, X. Huang, F. Wang, Y. Liu and P. An (China) | 479 |
| Burrowing behavior of <i>Chironomus yoshimatsui</i> larvae as an indicator of freshwater quality H.-h. Mo, J. Son, K.-S. Ryoo, Y.J. Bae and K. Cho (Republic of Korea) | 367 | The added value of bryophytes and macroalgae in ecological assessment of lakes F. Ecke (Sweden) | 487 |
| Variation in leaf chlorophyll concentration from tropical to cold-temperate forests: Association with gross primary productivity Y. Li, C. Liu, J. Zhang, H. Yang, L. Xu, Q. Wang (China), L. Sack (USA), X. Wu, J. Hou and N. He (China) | 377 | The mediation roles of intraspecific and interspecific functional trait diversity for linking the response of aboveground biomass to species richness across forest strata in a subtropical forest A. Ali and E.-R. Yan (China) | 493 |
| Streamflow as an ecological factor influencing radial growth of European ash (<i>Fraxinus excelsior</i> (L.)) M. Koprowski, B. Okoński (Poland), J. Gričar (Slovenia) and R. Puchałka (Poland) | 383 | A study on the effects of unbalanced data when fitting logistic regression models in ecology C. Salas-Eljatib, A. Fuentes-Ramirez (Chile), T.G. Gregoire (USA), A. Altamirano and V. Yaitul (Chile) | 502 |
| Relationship between stoichiometry and ecosystem services: A case study of organic farming systems F. Fan (Denmark, China), C.B. Henriksen (Denmark) and J. Porter (Denmark, France) | 390 | Improving stock unearthing method to measure soil erosion rates in vineyards J. Rodrigo-Comino (Spain, Germany) and A. Cerdà (Spain) | 509 |
| Factors influencing the nematode community during composting and nematode-based criteria for compost maturity H. Steel, T. Moens, B. Vandecasteele, F. Hendrickx, S. De Neve (Belgium), D.A. Neher (USA) and W. Bert (Belgium) | 400 | Plant diversity and pastoral value in alpine pastures are maximized at different nutrient indicator values M. Pittarello, M. Lonati, A. Gorlier, E. Perotti, M. Probo and G. Lombardi (Italy) | 518 |
| Assessing the utility of a novel terrestrial biodiversity quality indicator with 10 years of monitoring data T. McKenzie, L. Normand, N. Iwanycki, G. Miller and P. Prior (Canada) | 409 | Response of plant communities to climate change during the late Holocene: Palaeoecological insights from peatlands in the Alaskan Arctic M. Gałka (Poland), G.T. Swindles (UK), M. Szal (Poland), R. Fulweber (United States of America) and A. Feurdean (Germany, Romania) | 525 |
| Elucidating the importance of mussel carboxylesterase activity as exposure biomarker of environmental contaminants of current concern: An in vitro study M. Solé and J.C. Sanchez-Hernandez (Spain) | 422 | Assessing the ecological health of rivers when they are dry A.L. Steward, P. Negus, J.C. Marshall, S.E. Clifford and C. Dent (Australia) | 537 |
| | 432 | The influence of flow permanence and drying pattern on macroinvertebrate biomonitoring tools used in the assessment of riverine ecosystems N.A. Wilding, J.C. White, R.P. Chadd, A. House and P.J. Wood (United Kingdom) | 548 |

- Drifting space use of common cranes—Is there a mismatch between daytime behaviour and management?
L. Nilsson, M. Aronsson, J. Persson and J. Månsson (Sweden) 556
- Continent-wide test of the efficiency of the European union's conservation legislation in delivering population benefits for bird species
M. Koschová, J. Rivas-Salvador and J. Reif (Czech Republic) 563
- Are trawl marks a good indicator of trawling pressure in muddy sand fishing grounds?
L. Mérillet, D. Kopp, M. Robert, M. Salaün, S. Méhault, J.-F. Bourillet and M. Mouchet (France) 570
- Comparative impact of two glyphosate-based formulations in interaction with *Limnoperna fortunei* on freshwater phytoplankton
L.G. De Stefano, F. Gattás, A. Vinocur, D. Cristos, D. Rojas, D. Cataldo and H. Pizarro (Argentina) 575
- Soil enzyme response to permafrost collapse in the Northern Qinghai-Tibetan Plateau
H. Xu, G. Liu, X. Wu (China), J.M. Smoak (USA), C. Mu, X. Ma, X. Zhang, H. Li and G. Hu (China) 585
- Testing the efficiency of a bacterial community-based index (microgAMBI) to assess distinct impact sources in six locations around the world
A. Borja (Spain) 594
- Shift from dormancy to microbial growth revealed by RNA:DNA ratio
S. Loepmann (Germany), M. Semenov (Russian Federation), Y. Kuzyakov (Germany) and E. Blagodatskaya (Russian Federation, Germany) 603
- Relationship between socioeconomic vulnerability and ecological sustainability: The case of Aran-V-Bidgol's rangelands, Iran
V. Raufirad (Iran, USA), Q. Heidari (Iran), R. Hunter (USA) and J. Ghorbani (Iran) 613
- Evaluating and benchmarking biodiversity monitoring: Metadata-based indicators for sampling design, sampling effort and data analysis
S. Lengyel, B. Kosztyi (Hungary), D.S. Schmeller (Germany), P.-Y. Henry (France), M. Kotarac (Slovenia), Y.-P. Lin (Taiwan) and K. Henle (Germany) 624
- Relationships between biotic indices, multiple stressors and natural variability in New Zealand estuaries
A. Berthelsen, J. Atalah, D. Clark, E. Goodwin, M. Patterson and J. Sinner (New Zealand) 634
- Spatial apportionment of urban greenhouse gas emission inventory and its implications for urban planning: A case study of Xiamen, China
G. Zhang, R. Ge, T. Lin, H. Ye, X. Li and N. Huang (China) 644
- Determinants of submerged macrophytes palatability to grass carp *Ctenopharyngodon idellus*
J. Sun, L. Wang, L. Ma, T. Huang, W. Zheng, F. Min, Y. Zhang, Z. Wu and F. He (China) 657
- Ranking surface soil pollution potential of chemicals from accidental release by using two indicators calculated with a multimedia model (SoilPCA)
K.-E. Kim, J.E. Jung, Y. Lee and D.S. Lee (Republic of Korea) 664
- Composite eco-efficiency indicators for China based on data envelopment analysis
J. Huang, J. Xia, Y. Yu and N. Zhang (China) 674
- Testing the ecoacoustics event detection and identification (EEDI) approach on Mediterranean soundscapes
A. Farina (Italy), S.H. Gage (USA) and P. Salutari (Italy) 698
- A new approach of deriving indicators and comprehensive measure for ecological environmental quality assessment
L.H. Chai and D. Lha (China) 716
- Evaluation of the ecological sensitivity and security of tidal flats in Shanghai
Y. Shi, J. Li and M. Xie (China) 729
- Spatial distribution of forest biomass carbon (Above and below ground) in Indian forests
G. Rajashekar, R. Fararoda, R.S. Reddy, C.S. Jha, K.N. Ganeshiah, J.S. Singh and V.K. Dadhwal (India) 742
- Lead accumulation in rats: The effect of the presence of a rat tapeworm and the different forms of metal in the host diet
B. Horáková, Z. Čadková, J. Száková, I. Jankovská and I. Langrová (Czech Republic) 753
- Towards a better understanding of Green Infrastructure: A critical review
J. Wang and E. Banzhaf (Germany) 758
- Soil Biological Quality index (QBS-ar): 15 years of application at global scale
C. Menta, F.D. Conti, S. Pinto and A. Bodini (Italy) 773
- Influence of debris flows on macroinvertebrate diversity and assemblage structure
Y. Yang, X. Zhou, Y. Yi, M. Xu and Z. Yang (China) 781
- Modeling current and future freshwater inflow needs of a subtropical estuary to manage and maintain forested wetland ecological conditions
M.M. Baustian, F.R. Clark, A.S. Jerabek, Y. Wang, H.C. Bienn and E.D. White (USA) 791
- A methodology for evaluating transdisciplinary research on coupled socio-ecological systems
J.M. Holzer, N. Carmon and D.E. Orenstein (Israel) 808
- Predicting the effect of land use and climate change on stream macroinvertebrates based on the linkage between structural equation modeling and bayesian network
X. Li, Y. Zhang, F. Guo, X. Gao and Y. Wang (China) 820
- Ecobordure: A flora-based indicator to assess vegetation patterns of field margins and infer its local drivers. Design in Brittany (France)
A. Alignier, D. Le Cœur, E. Lanoë, F. Ferchaud, B. Roche and C. Thenail (France) 832

| | | | |
|--|-----|--|------|
| Green competitiveness evaluation of provinces in China based on correlation analysis and fuzzy rough set X. Cheng, R. Long, H. Chen and W. Li (China) | | Impact of an agri-environmental scheme on landscape patterns E.P. Tanner and S.D. Fuhlendorf (USA) | 956 |
| On the reliability of the Elements of Metacommunity Structure framework for separating idealized metacommunity patterns D. Schmera, J. Podani, Z. Botta-Dukát and T. Erős (Hungary) | 841 | Development of a water quality index for rivers in West Java Province, Indonesia A.D. Sutadian (Australia, Indonesia), N. Muttill (Australia), A.G. Yilmaz (United Arab Emirates) and B.J.C. Perera (Australia) | 966 |
| Reliability of simplifying strategies for rapid biodiversity assessment in studying community-environment interactions W. Xiong, Y. Yang and A. Zhan (China) | | Remotely sensed spatial heterogeneity as an exploratory tool for taxonomic and functional diversity study D. Rocchini, G. Bacaro, G. Chirici, D. Da Re (Italy), H. Feilhauer (Germany), G.M. Foody (UK), M. Galluzzi (Italy), C.X. Garzon-Lopez (Colombia), T.W. Gillespie, K.S. He (USA), J. Lenoir (France), M. Marcantonio (USA), H. Nagendra (India), C. Ricotta (Italy), E. Rommel, S. Schmidlein (Germany), A.K. Skidmore (The Netherlands, Australia), R. Van De Kerchove (Belgium), M. Wegmann (Germany) and B. Rugani (Luxembourg) | 983 |
| Decadal shifts in macroalgae assemblages in impacted urban lagoons in Brazil H.B.F. Amaral, R.P. Reis, M.A. de Oliveira Figueiredo and A. de Gusmão Pedrini (Brazil) | 861 | Merging community assembly into the regime-shift approach for informing ecological restoration K. Kadowaki, S. Nishijima (Japan), S. Kéfi (France), K.O. Kameda and T. Sasaki (Japan) | 991 |
| Fatty acid trophic markers in Lake Baikal phytoplankton: A comparison of endemic and cosmopolitan diatom-dominated phytoplankton assemblages S.M. Shishlyannikov, A.A. Nikonova, Y.S. Bukin and A.G. Gorshkov (Russia) | 869 | Towards developing a river health index P.K. Singh and S. Saxena (India) | 999 |
| Persistent negative changes in seasonal greenness over different forest types of India using MODIS time series NDVI data (2001–2014) A. Chakraborty, M.V.R. Seshasai, C.S. Reddy and V.K. Dadhwal (India) | 878 | Multiclass classification methods in ecology M. Bourel and A.M. Segura (Uruguay) | 1012 |
| A sustainability index with attention to environmental justice for eco-city classification and assessment L. Liu (USA) | | A semi-parametric geographically weighted (S-GWR) approach for modeling spatial distribution of population Y. Huang, C. Zhao, X. Song, J. Chen and Z. Li (China) | 1022 |
| Habitat factors associated with the decline in brown hare abundance in Poland in the beginning of the 21st century M. Panek (Poland) | | Acoustic comparison of a patchy Mediterranean shallow water seascape: <i>Posidonia oceanica</i> meadow and sandy bottom habitats M. Ceraulo, E. Papale (Italy), F. Caruso (Italy, USA), F. Filiciotto, R. Grammauta, I. Parisi, S. Mazzola, A. Farina and G. Buscaino (Italy) | 1030 |
| Will the shift from crude oil to natural gas burning for power generation at an oil refinery increase ozone concentrations in the region of Cubatão (SE-Brazil)? P.I.L.S. Assis, R.M. Moraes and R.K. Nakazato (Brazil) | 915 | Development and preliminary validation of a multi-trophic metabarcoding biotic index for monitoring benthic organic enrichment N. Keeley (New Zealand, Norway), S.A. Wood and X. Pochon (New Zealand) | 1044 |
| The assessment of the changes of biomass and riparian buffer width in the terminal reservoir under the impact of the South-to-North Water Diversion Project in China S. Yang, J. Bai, C. Zhao, H. Lou, C. Zhang, Y. Guan, Y. Zhang, Z. Wang and X. Yu (China) | 932 | Effects of watershed and in-stream liming on macroinvertebrate communities in acidified tributaries to an Adirondack lake S.D. George, B.P. Baldigo, G.B. Lawrence and R.L. Fuller (United States) | 1058 |
| Insights on historical expansions of desertification in the Hunlun Buir and Horqin Deserts of Northeast China J. Li, L. Han, Y. Liu, G. Zhang and Z. Wu (China) | 944 | Exploring temporal dynamics of urban ecosystem services in Latin America: The case of Bogota (Colombia) and Santiago (Chile) C. Dobbs, Á. Hernández-Moreno, S. Reyes-Paecke and M.D. Miranda (Chile) | 1068 |
| Assessing land take and its effects on dune carbon pools. An insight into the Mediterranean coastline M.L. Carranza, M. Drius (Italy), M. Malavasi (Czech Republic), L. Frate, A. Stanisci and A.T.R. Acosta (Italy) | 951 | | |

| | | | |
|--|------|---|------|
| Towards a system-specific framework for the sustainability evaluation of low-input ruminant meat production systems in developing countries T. Marandure, G. Makombe, K. Dzama, W. Hoffmann and C. Mapiye (South Africa) | 1081 | Sources, influencing factors and environmental indications of PAH pollution in urban soil columns of Shanghai, China S. Wu, X. Liu, M. Liu, X. Chen, S. Liu, L. Cheng, X. Lin and Y. Li (China) | 1170 |
| Intercalibration of counting methods for <i>Ostreopsis</i> spp. blooms in the Mediterranean Sea M. Vassalli, A. Penna, F. Sbrana, S. Casabianca, N. Gjerci, S. Capellacci, V. Asnaghi, E. Ottaviani, V. Giussani, L. Pugliese (Italy), C. Jauzein, R. Lemée (France), M.A. Hachani, S. Turki (Tunisia), L. Açaf, M.A.-A. Saab (Lebanon), A. Fricke (France, Argentina), L. Mangialajo (France), R. Bertolotto, C. Totti, S. Accoroni (Italy), E. Berdalet, M. Vila (Spain) and M. Chiantore (Italy) | 1092 | Multi-scale assessment of forest cover in an agricultural landscape of Southeastern Brazil: Implications for management and conservation of stream habitat and water quality F.R. de Paula, P. Gerhard, S.F. de Barros Ferraz (Brazil) and S.J. Wenger (USA) | 1181 |
| Arsenate inhibition on kinetic characteristics of alkaline phosphatase as influenced by pH H. Tian, Y. Zhao (China), M. Megharaj (Australia) and W. He (China) | 1101 | Ciliates as bioindicators of CO ₂ in soil R. Gabilondo, S. Blanco, I. Fernández-Montiel, D.A. García and E. Bécares (Spain) | 1192 |
| Comprehensiveness or implementation: Challenges in translating farm-level sustainability assessments into action for sustainable development E.M. de Olde (The Netherlands), M. Sautier (France) and J. Whitehead (New Zealand) | 1107 | Food web organization following the invasion of habitat-modifying <i>Tubastraea</i> spp. corals appears to favour the invasive borer bivalve <i>Leiosolenus aristatus</i> C. Vinagre, R. Silva, V. Mendonça (Portugal), A.A.V. Flores (Brazil), A. Baeta and J.C. Marques (Portugal) | 1204 |
| Otter fecal enterococci as general indicators of antimicrobial resistance dissemination in aquatic environments T. Semedo-Lemsaddek (Portugal), N.M. Pedroso (Brazil, Portugal), D. Freire, T. Nunes, L. Tavares (Portugal), L.M. Verdade (Brazil) and M. Oliveira (Portugal) | 1113 | Measuring macrofungal biodiversity quality using two different survey approaches: A case study in broadleaf Mediterranean forests E. Ambrosio, M.G. Mariotti, M. Zotti, G. Cecchi, S. Di Piazza (Italy) and A. Feest (UK) | 1210 |
| Economic value of marine biodiversity improvement in coralligenous habitats S. Tonin (Italy) | 1121 | An optimization framework for supporting decision making in biodiesel feedstock imports: Water footprint vs. import costs P.P. Miglietta, S. Giove and P. Toma (Italy) | 1231 |
| Mapping watershed integrity for the conterminous United States D.J. Thornbrugh, S.G. Leibowitz, R.A. Hill, M.H. Weber, Z.C. Johnson, A.R. Olsen, J.E. Flotemersch, J.L. Stoddard and D.V. Peck (USA) | 1133 | Assessing the influence of multiple stressors on stream diatom metrics in the upper Midwest, USA M.D. Munn, I. Waite and C.P. Konrad (United States) | 1239 |
| Disentangling the response of lake littoral invertebrate assemblages to multiple pressures R.K. Johnson, S. Hallstan (Sweden) and X. Zhao (Sweden, China) | 1149 | Sustainability evaluation based on the emergy ecological footprint method: A case study of Qingdao, China, from 2004 to 2014 W. Peng, X. Wang, X. Li and C. He (China) | 1249 |
| Optimization schemes for grassland ecosystem services under climate change R. Hao and D. Yu (China) | 1158 | Benefit relevant indicators: Ecosystem services measures that link ecological and social outcomes L.P. Olander, R.J. Johnston, H. Tallis, J. Kagan, L.A. Maguire, S. Polasky, D. Urban, J. Boyd, L. Wainger and M. Palmer (USA) | 1262 |
| | | Improved space-time mapping of PM _{2.5} distribution using a domain transformation method G. Christakos (China, USA), Y. Yang, J. Wu, C. Zhang, Y. Mei and J. He (China) | 1273 |