

## Contents

### Research articles

Yield and water use of drought-tolerant maize hybrids in a semiarid environment

J. Zhao, Q. Xue, K.E. Jessup (USA), B. Hao (China), X. Hou, T.H. Marek, W. Xu, S.R. Evett, S.A. O'Shaughnessy and D.K. Brauer (USA)

1

Removal of early fruiting branches impacts leaf senescence and yield by altering the sink/source ratio of field-grown cotton

Y. Chen, X. Kong and H. Dong (China)

10

Characterizing soybean vigor and productivity using multiple crop canopy sensor readings

J.J. Miller, J.S. Schepers, C.A. Shapiro, N.J. Arneson, K.M. Eskridge, M.C. Oliveira and L.J. Giesler (United States)

22

Whole-genome prediction of reaction norms to environmental stress in bread wheat (*Triticum aestivum* L.) by genomic random regression

D. Ly, S. Huet, A. Gauffreteau, R. Rincent, G. Touzy, A. Mini (France), J.-L. Jannink (United States), F. Cormier, E. Paux, S. Lafarge, J. Le Gouis and G. Charmet (France)

32

Genotype by tillage interaction and performance progress for bread and durum wheat genotypes on irrigated raised beds

N. Honsdorf, M.J. Mulvaney, R.P. Singh, K. Ammar, J. Burgueño, B. Govaerts and N. Verhulst (Mexico)

42

Application of zinc improves the productivity and biofortification of fine grain aromatic rice grown in dry seeded and puddled transplanted production systems  
M. Farooq (Pakistan, Oman, Australia), A. Ullah, A. Rehman, A. Nawaz, A. Nadeem, A. Wakeel, F. Nadeem (Pakistan) and K.H.M. Siddique (Australia)

53

Yield loss compensation effect and water use efficiency of winter wheat under double-blank row mulching and limited irrigation in northern China

Q. Yan, F. Yang, F. Dong, J. Lu, F. Li, Z. Duan, J. Zhang (China) and G. Lou (United States)

63

Physical robustness of canopy temperature models for crop heat stress simulation across environments and production conditions

H. Webber (Germany), J.W. White, B.A. Kimball (USA), F. Ewert (Germany), S. Asseng (USA), E. Eyshi Rezaei (Germany), P.J. Pinter Jr., J.L. Hatfield (USA), M.P. Reynolds (Mexico), B. Ababaei (France), M. Bindi (Italy), J. Doltra (Spain), R. Ferrise (Italy), H. Kage (Germany), B.T. Kassie (USA), K.-C. Kersebaum, A. Luig (Germany), J.E. Olesen (Denmark), M.A. Semenov, P. Stratonovitch (UK), A.M. Ratjen (Germany), R.L. LaMorte, S.W. Leavitt, D.J. Hunsaker, G.W. Wall (USA) and P. Martre (France)

75

The effects of plastic-film mulch on the grain yield and root biomass of maize vary with cultivar in a cold semiarid environment L. Wang, X.G. Li, Z.-H. Guan, B. Jia (China), N.C. Turner (Australia) and F.-M. Li (China)	89	Photogrammetry for the estimation of wheat biomass and harvest index J. Walter, J. Edwards, G. McDonald and H. Kuchel (Australia)	165
Can optimization of phosphorus input lead to high productivity and high phosphorus use efficiency of cotton through maximization of root/mycorrhizal efficiency in phosphorus acquisition? W. Mai, X. Xue, G. Feng, R. Yang and C. Tian (China)	100	Remote sensing-based crop biomass with water or light-driven crop growth models in wheat commercial fields I. Campos, L. González-Gómez, J. Villodre, J. González-Piqueras (Spain), A.E. Suyker (USA) and A. Calera (Spain)	175
Effects of alfalfa intercropping on crop yield, water use efficiency, and overall economic benefit in the Corn Belt of Northeast China T. Sun, Z. Li, Q. Wu, T. Sheng and M. Du (China)	109	Genotype, environment, and genotype by environment interaction for seed isoflavone concentration in soybean grown in soybean cyst nematode infested and non-Infested environments A. Carter, I. Rajcan, L. Woodrow, A. Navabi and M. Eskandari (Canada)	189
Nitrogen topdressing timing influences the spatial distribution patterns of protein components and quality traits of flours from different pearling fractions of wheat ( <i>Triticum aestivum</i> L.) grains Y. Zhong, M. Yang, J. Cai, X. Wang, Q. Zhou, W. Cao, T. Dai and D. Jiang (PR China)	120	Maize and wheat root biomass, vertical distribution, and size class as affected by fertilization intensity in two long-term field trials J. Hirte, J. Leifeld, S. Abiven and J. Mayer (Switzerland)	197
Impact of high temperatures in maize: Phenology and yield components J.I. Lizaso, M. Ruiz-Ramos, L. Rodríguez, C. Gabaldon-Leal, J.A. Oliveira, I.J. Lorite, D. Sánchez (Spain), E. García (México) and A. Rodríguez (Spain)	129	Physiological activity and biomass production in crop canopy under a tropical environment in soybean cultivars with temperate and tropical origins A. Saryoko (Japan, Indonesia), Y. Fukuda (Japan), I. Lubis (Indonesia), K. Homma and T. Shiraiwa (Japan)	209
Contribution of the early-established plant hierarchies to maize crop responses to N fertilization M.A. Rossini, M.E. Otegui, E.L. Martínez and G.A. Maddonni (Argentina)	141	Liming and straw retention interact to increase nitrogen uptake and grain yield in a double rice-cropping system P. Liao, S. Huang (China), N.C. van Gestel (United States), Y. Zeng, Z. Wu (China) and K.J. van Groenigen (UK)	217
Different mechanisms underlying the yield advantage of ordinary hybrid and super hybrid rice over inbred rice under low and moderate N input conditions L. Huang, F. Sun, S. Yuan, S. Peng and F. Wang (China)	150	Evaluation of optimal nitrogen rate for corn production under mulched drip fertigation and economic benefits D. Wang, G. Li, Y. Mo, M. Cai and X. Bian (China)	225
Azolla biofertilizer for improving low nitrogen use efficiency in an intensive rice cropping system Y. Yao, M. Zhang, Y. Tian, M. Zhao, K. Zeng, B. Zhang, M. Zhao and B. Yin (China)	158		