

INDUSTRIAL CROPS AND PRODUCTS

VOLUME 111

January 2018

Contents

Fats and oils

- Cistus incanus* L. extract inhibits Aflatoxin B₁ production by *Aspergillus parasiticus* in macadamia nuts
V. Kalli, E. Kollia, A. Roidaki, C. Proestos and P. Markaki (Greece) 63
- Exogenous proline improves osmoregulation, physiological functions, essential oil, and seed yield of fennel
A. Gholami Zali and P. Ehsanzadeh (Iran) 133
- Development of genomic simple sequence repeat (gSSR) markers in cumin and their application in diversity analyses and cross-transferability
R. Bharti, S. Kumar and M.J. Parekh (India) 158
- Summer savory (*Satureja hortensis* L.) essential oil constituent oscillation at different storage conditions
S. Mohtashami, V. Rowshan, L. Tabrizi, M. Babalar and A. Ghani (Iran) 226
- Water stress and manure application affect the quantity and quality of essential oil of *Thymus daenensis* and *Thymus vulgaris*
M. Askary, M.A. Behdani, S. Parsa, S. Mahmoodi and M. Jamialahmadi (Iran) 336
- Agronomic performance of the novel oilseed crop *Centropalus pauciflorus* in southwestern Ontario
J. Todd, S. Chakraborty (Canada), T. Isbell (USA) and R.C. Van Acker (Canada) 364
- Analysis of volume ratio of castor/soybean oil mixture on minimum quantity lubrication grinding performance and microstructure evaluation by fractal dimension
S. Guo, C. Li, Y. Zhang, M. Yang, D. Jia, X. Zhang, G. Liu (China), R. Li (United States), Z. Bing and H. Ji (China) 494
- Phenotypic stability, genotype × environmental interactions, and cultivar recommendations for essential oil yield in khus aromatic grass (*Chrysopogon zizanioides* (L.) Roberty)
R.K. Lal, C.S. Chanotiy, P. Gupta, S. Sarkar, S. Singh, R. Maurya, S. Srivastva and P.K. Chaudhary (India) 871

Non-food bioactive products

- Sumac (*Rhus coriaria* L.) fruit: Essential oil variability in Iranian populations
M.R. Morshedloo (Iran), F. Maggi (Italy), H. Tavakoli Neko and M. Soleimani Aghdam (Iran) 1
- Chemical composition and biological activities of extracts from three *Salvia* species: *S. blepharochlaena*, *S. euphratica* var. *leiocalycina*, and *S. verticillata* subsp. *amasiaca*
G. Zengin (Turkey), E.J. Llorent-Martínez, M.L.F.-d. Córdoba (Spain), M.B. Bahadori (Iran), A. Mocan (Romania), M. Locatelli (Italy) and A. Aktumsek (Turkey) 11

- Comparative study of the essential oils of four *Pinus* species: Chemical composition, antimicrobial and insect larvicidal activity
Z.S. Mitić (Serbia), B. Jovanović (Germany), S.Č. Jovanović, T. Mihajilov-Krstev, Z.Z. Stojanović-Radić, V.J. Cvetković, T.Lj. Mitrović, P.D. Marin, B.K. Zlatković and G.S. Stojanović (Serbia) 55
- Publisher's Note
J.N. Eloff, I.E. Angeh, L.J. McGaw (South Africa) 69
- Assessment of alcoholic distillates for the extraction of bioactive polyphenols from grapevine canes
T. Rodríguez-Cabo, I. Rodríguez, M. Ramil and R. Cela (Spain) 99
- Instrumental characterization and functional assessment of the two-color silk fabric coated by the extract from *Dioscorea cirrhosa* tuber and mordanted by iron salt-containing mud
T.-T. Yang, J.-P. Guan, G. Chen and R.-C. Tang (China) 117
- Citrus reticulata* Blanco peels as a source of antioxidant and anti-proliferative phenolic compounds
S.S. Ferreira, A.M. Silva and F.M. Nunes (Portugal) 141
- Comprehensive evaluation of effective constituents in total alkaloids from *Sophora alopecuroides* L. and their joint action against aphids by laboratory toxicity and field efficacy
T. Ma, H. Yan, X. Shi, B. Liu, Z. Ma and X. Zhang (China) 149
- Onosma heterophyllum*: Phenolic composition, enzyme inhibitory and antioxidant activities
M.S. Ozer, B. Kirkan, C. Sarikurkcu, M. Cengiz, O. Ceylan, N. Atilgan and B. Tepe (Turkey) 179
- Microemulsions containing *Copaifera multijuga* Hayne oil-resin: Challenges to achieve an efficient system for β -caryophyllene delivery
J.K. de Oliveira Neves, A.C. Apolinário, K.L. Alcantara Saraiva, D.T.C. da Silva, M.Y. de Freitas Araújo Reis, B.P.G. de Lima Damasceno, A. Pessoa Jr., M.A.M. Galvão, L.A.L. Soares, V.F. da Veiga Júnior, J.A. da Silva (Brazil) and A. Converti (Italy) 185
- Bioactive characterization of *Persea americana* Mill. by-products: A rich source of inherent antioxidants
B. Melgar (Portugal, Spain), M.I. Dias (Portugal), A. Ciric, M. Sokovic (Serbia), E.M. Garcia-Castello, A.D. Rodriguez-Lopez (Spain), L. Barros and I.C.R.F. Ferreira (Portugal) 212
- Antioxidant and antimicrobial properties of encapsulated guava leaf oil in hydroxypropyl-beta-cyclodextrin
J. Rakmai, B. Cheirsilp (Thailand), J.C. Mejuto, J. Simal-Gándara and A. Torrado-Agrasar (Spain) 219
- Comparative study of the chemical composition and bioactivities of essential oils of fresh and dry seeds from *Myoporum insulare* R. Br.
A. Zardi-Bergaoui, S. Jelizi (Tunisia), G. Flamini (Tunisia, Italy), R. Ascrizzi and H. Ben Jannet (Tunisia) 232
- Skin-care effects of dandelion leaf extract and stem extract: Antioxidant properties, tyrosinase inhibitory and molecular docking simulations
P.-j. Xie, L.-x. Huang, C.-h. Zhang, S.-s. Ding, Y.-j. Deng and X.-j. Wang (China) 238
- Phytochemical profiles and bioactivities of essential oils extracted from seven *Curcuma* herbs
H. Xiang, L. Zhang, L. Xi, Y. Yang, X. Wang, D. Lei (PR China), X. Zheng (USA) and X. Liu (PR China) 298
- Antimicrobial and antioxidant features of 'Gabsi' pomegranate peel extracts
S. Kharchoufi (Tunisia), F. Licciardello, L. Siracusa, G. Muratore (Italy), M. Hamdi (Tunisia) and C. Restuccia (Italy) 345
- In silico-assisted identification of α -amylase inhibitor from the needle oil of *Pinus tabulaeformis* Carr.
J. Yang, D. Gu, M. Wang, D. Kou, H. Guo, J. Tian and Y. Yang (China) 360
- Three new monoterpene glycosides from oil peony seed cake
P. Liu, Y. Zhang, Y.-F. Xu, X.-Y. Zhu, X.-F. Xu, S. Chang and R.-X. Deng (China) 371
- Solvent selection for efficient extraction of bioactive compounds from grape pomace
D. Pintać, T. Majkić, L. Torović, D. Orčić, I. Beara, N. Simin, N. Mimica-Dukić and M. Lesjak (Serbia) 379
- In vitro* and *in silico* approaches to appraise *Polygonum maritimum* L. as a source of innovative products with anti-ageing potential
M.J. Rodrigues (Portugal), S. Słusarczyk, Ł. Pecio, A. Matkowski (Poland), R.E. Salmas, S. Durdagi (Turkey), C. Pereira, J. Varela, L. Barreira and L. Custódio (Portugal) 391
- Bioactive phenolics from mango leaves (*Mangifera indica* L.)
J. Pan, X. Yi, S. Zhang, J. Cheng, Y. Wang, C. Liu and X. He (China) 400
- Antioxidant capacities and total phenolic contents of 30 flowers
G.-L. Chen, S.-G. Chen, Y. Xiao and N.-L. Fu (China) 430
- Phenylpropanoid content in high-altitude cultivated *Rhodiola rosea* L. provenances according to plant part, harvest season and age
W. Peschel (United Kingdom), A. Kump (Austria), Z.P. Zomborszki (Hungary), M. Pfosser, W. Kainz (Austria) and D. Csupor (Hungary) 446

- Development and validation of a new HPTLC method for quantification of conophylline in *Tabernaemontana divaricata* samples obtained from different seasons and extraction techniques: Insights into variation of pancreatic lipase inhibitory activity
Sridhar S.N.C., P. Sengupta and A.T. Paul (India) 462
- Effectiveness of defatted seed meals from Brassicaceae with or without crude glycerin against black grass (*Alopecurus myosuroides* Huds.)
R. Matteo (Italy), M.A. Back, J.P.H. Reade (UK), L. Ugolini, E. Pagnotta and L. Lazzeri (Italy) 506
- Diversity of flavonoids profile in China aster [*Callistephus chinensis* (L.) Nees.] genotypes
V. Bhargav, R. Kumar, K.S. Shivashankara, T.M. Rao, M.V. Dhananjaya, A. Sane, T.U. Bharathi, R. Venugopalan and T.K. Roy (India) 513
- Phenolic composition and biological properties of *Achillea nobilis* L. subsp. *neilreichii* (Kerner) Formanek
D. Taşkın, T. Taşkın and E. Rayaman (Turkey) 555
- Subcritical water extraction as an environmentally-friendly technique to recover bioactive compounds from traditional Serbian medicinal plants
N. Nastić, J. Švarc-Gajić (Serbia), C. Delerue-Matos, M.F. Barroso, C. Soares, M.M. Moreira, S. Morais (Portugal), P. Mašković (Serbia), V. Gaurina Srček, I. Slivac, K. Radošević (Croatia) and M. Radojković (Serbia) 579
- Tyrosinase, elastase, hyaluronidase, inhibitory and antioxidant activity of Sri Lankan medicinal plants for novel cosmeceuticals
G.D. Liyanaarachchi, J.K.R.R. Samarasekera, K.R.R. Mahanama (Sri Lanka) and K.D.P. Hemalal (Canada,) 597
- Ultrasound-assisted optimal extraction and thermal stability of betalains from colored quinoa (*Chenopodium quinoa* Willd) hulls
C. Laqui-Vilca, S. Aguilar-Tuesta, W. Mamani-Navarro, J. Montaña-Bustamante and L. Condezo-Hoyos (Peru) 606
- Genetic diversity of *Thymus sibthorpii* Benthams in mountainous natural grasslands of Northern Greece as related to local factors and plant community structure
E.M. Abraham, A. Aftzalanidou, I. Ganopoulos (Greece), M. Osathanukul (Thailand), A. Xanthopoulou, E. Avramidou, E. Sarrou, F. Aravanopoulos and P. Madesis (Greece) 651
- In vitro* and *in vivo* transformations of *Centaureum erythraea* secoiridoid glucosides alternate their antioxidant and antimicrobial capacity
J. Božunović, S. Živković, U. Gašić, J. Glamočlija, A. Čirić, D. Matekalo, B. Šiler, M. Soković, Ž. Tešić and D. Mišić (Serbia) 705
- Synthesis and herbicidal activity of 4, 8-DHT and its derivatives
J.-h. Zhang, L. Yang, X. Ruan, Z.-l. Sheng, M.-f. Yu, B.-s. Zheng, J.-y. Zhang, X.-x. Li, Y.-x. Zhao and Q. Wang (PR China) 755
- Supercritical CO₂ extraction of *Rosmarinus eriocalyx* growing in Algeria: Chemical composition and antioxidant activity of extracts and their solid plant materials
H. Bendif, M.D. Miara (Algeria), Z. Kalboussi (Tunisia), D. Grauzdytė, D. Povilaitis, P.R. Venskutonis (Lithuania) and F. Maggi (Italy) 768
- Secondary treated effluent irrigation did not impact chemical composition, and enzyme inhibition activities of essential oils from *Origanum syriacum* var. *syriacum*
M.S. Ali-Shtayeh, R.M. Jamous, S.Y. Abu-Zaitoun, R.J. Akkawi, S.R. Kalbouneh (Palestine), N. Dudai and N. Bernstein (Israel) 775
- Native Mexican aromatic flora and essential oils: Current research status, gaps in knowledge and agro-industrial potential
L.M. Calvo-Irabien (Mexico) 807
- Screening for amyloid-β aggregation inhibitor and neuronal toxicity of eight Tunisian medicinal plants
Z. Dhouafli (Tunisia), S. Rigacci, M. Leri, M. Bucciantini (Italy), B. Mahjoub, M.S. Tounsi, W.A. Wannas (Tunisia), M. Stefani (Italy) and E.A. Hayouni (Tunisia) 823
- Plant growth, steviol glycosides and nutrient uptake as affected by arbuscular mycorrhizal fungi and phosphorous fertilization in *Stevia rebaudiana* Bert
S. Tavarini, B. Passera, A. Martini, L. Avio, C. Sbrana, M. Giovannetti and L.G. Angelini (Italy) 899
- Natural rubber and resins**
- Correlation between DNA methylation status and rubber yield and related characteristics in *Hevea brasiliensis* tapped at different heights
C. Wu, S. Ban, X. Gao and W. Li (PR China) 563
- Fibres, proteins and carbohydrates**
- Chemical, fatty acid, polyphenolic profile, techno-functional and antioxidant properties of flours obtained from quinoa (*Chenopodium quinoa* Willd) seeds
M. Pellegrini (Italy), R. Lucas-Gonzales (Spain), A. Ricci (Italy), J. Fontecha, J. Fernández-López, J.A. Pérez-Álvarez and M. Viuda-Martos (Spain) 38
- Characteristics and environmentally friendly extraction of cellulose nanofibrils from sugarcane bagasse
Y.-H. Feng, T.-Y. Cheng, W.-G. Yang, P.-T. Ma, H.-Z. He, X.-C. Yin and X.-X. Yu (PR China) 285

Reinforcement of natural fiber yarns by cellulose nanomaterials: A multi-scale study <i>S. Ghasemi, M. Tajvidi, D.W. Bousfield and D.J. Gardner (USA)</i>	471	Catalytic effect of ZSM-5 zeolite and HY-340 niobic acid on the pyrolysis of industrial kraft lignins <i>J.A. Santana Junior, W.S. Carvalho and C.H. Ataíde (Brazil)</i>	126
Leaf area index, light interception, growth and steviol glycoside formation of <i>Stevia rebaudiana</i> Bertoni under field conditions in southwestern Germany <i>S. Munz, A. Präger, N. Merkt, W. Claupein and S. Graeff-Hönninger (Germany)</i>	520	A novel and efficient process for lignin fractionation in biomass-derived glycerol-ethanol solvent system <i>C. Liu, C. Si, G. Wang, H. Jia and L. Ma (China)</i>	201
Manufacturing & characterization of regenerated cellulose/curcumin based sustainable composites fibers spun from environmentally benign solvents <i>M.G. Coscia (Rome), J. Bhardwaj, N. Singh (United Kingdom), M.G. Santonicola (Rome), R. Richardson, V.K. Thakur and S. Rahatekar (United Kingdom)</i>	536	Structural characterization of birch lignin isolated from a pressurized hot water extraction and mild alkali pulped biorefinery process <i>L. Lagerquist, A. Pranovich, A. Smeds, S. von Schoultz, L. Vähäsalo, J. Rahkila, I. Kilpeläinen, T. Tamminen, S. Willför and P. Eklund (Finland)</i>	306
Two-step isolation of hemicelluloses from Chinese quince fruit: Effect of hydrothermal treatment on structural features <i>H.-M. Liu, Y.-R. Li, M. Wu, H.-S. Yin and X.-D. Wang (China)</i>	615	Use of surfactant and enzymes in dry-grind corn ethanol fermentation improves yield of ethanol and distillers corn oil <i>L. Fang, T. Wang and B. Lamsal (United States)</i>	329
Proteomic analysis of non-toxic <i>Jatropha curcas</i> byproduct cake: Fractionation and identification of the major components <i>A. León-Villanueva, J.A. Huerta-Ocampo, A. Barrera-Pacheco, S. Medina-Godoy and A.P.B. de la Rosa (Mexico)</i>	694	Oxidative depolymerization potential of biorefinery lignin obtained by ionic liquid pretreatment and subsequent enzymatic saccharification of eucalyptus <i>K. Ninomiya, K. Ochiai, M. Eguchi, K. Kuroda, Y. Tsuge, C. Ogino, T. Taima and K. Takahashi (Japan)</i>	457
Influence of thermo-pressing conditions on insulation materials from bamboo fibers and proteins based bone glue <i>D.M. Nguyen, A.-C. Grillet (France), T.M.H. Diep (Vietnam), Q.B. Bui (France, Vietnam) and M. Woloszyn (France)</i>	834	<i>Prunus tomentosa</i> seed waste as a source of aromatic glycosides: Valuable phytochemicals with α -glucosidase inhibitory and hepatoprotective properties <i>Q.-B. Liu, Z.-Y. Cheng, Z.-Y. Yan, D. Wang, M. Bai, X.-X. Huang and S.-J. Song (PR China)</i>	590
Biorefinery, valorization of byproducts		Optimized feeding schemes of simultaneous saccharification and fermentation process for high lactic acid titer from sugarcane bagasse <i>P. Unrean (Thailand)</i>	660
Implementation of phenols recovered from olive mill wastewater as UV booster in cosmetics <i>C.M. Galanakis (Greece, Austria), P. Tsatalas and I.M. Galanakis (Greece)</i>	30	Influence of different thermal pretreatments and inoculum selection on the biomethanation of sugarcane bagasse by solid-state anaerobic digestion: A kinetic analysis <i>D.R.S. Lima, O.F.H. Adarme, B.E.L. Baêta, L.V.A. Gurgel and S.F. de Aquino (Brazil)</i>	684
Co-production of oligosaccharides and fermentable sugar from wheat straw by hydrothermal pretreatment combined with alkaline ethanol extraction <i>X. Chen, H. Li, S. Sun, X. Cao and R. Sun (China)</i>	78	Optimization of acid and enzymatic hydrolysis of kodo millet (<i>Paspalum scrobiculatum</i>) bran residue to obtain fermentable sugars for the production of optically pure d (-) lactic acid <i>R. Balakrishnan, S.R. Reddy Tadi, S. Sivaprakasam and S. Rajaram (India)</i>	731
Phenolic compounds recovery from grape skin using conventional and non-conventional extraction methods <i>T.W. Caldas, K.E.L. Mazza, A.S.C. Teles, G.N. Mattos, A.I.S. Brígida, C.A. Conte-Junior, R.G. Borguini, R.L.O. Godoy, L.M.C. Cabral and R.V. Tonon (Brazil)</i>	86	Effects of cultivation period on catch crop chemical composition and potential for bioenergy production <i>S. Akao, D. Yasutake, K. Kondo, H. Nagare, M. Maeda and T. Fujiwara (Japan)</i>	787
Fennel oil and by-products seed characterization and their potential applications <i>B. Sayed Ahmad (France, Lebanon), T. Talou (France), Z. Saad, A. Hijazi (Lebanon), M. Cerny (France), H. Kanaan (Lebanon, France), A. Chokr (Lebanon) and O. Merah (France)</i>	92	Thermal stability, degradation kinetics, and molecular weight of organosolv lignins from <i>Pinus radiata</i> <i>J.C. Domínguez, T.M. Santos, V. Rigual, M. Oliet, M.V. Alonso and F. Rodriguez (Spain)</i>	889

Bioenergy - crops and processing

- A high-Oil castor cultivar developed through recurrent selection
G.Q. Chen, K. Johnson, E. Morales, A.M. Ibáñez and J.-T. Lin (USA) 8
- Camelina sativa* as a fallow replacement crop in wheat-based crop production systems in the US Great Plains
A.K. Obour, C. Chen, H.Y. Sintim, K. McVay, P. Lamb, E. Obeng, Y.A. Mohammed, Q. Khan, R.K. Afshar and V.D. Zheljaskov (United States) 22
- Elephant grass leaves have lower recalcitrance to acid pretreatment than stems, with higher potential for ethanol production
C.C. Santos, W. de Souza, C. Sant 'Anna and M. Brienzo (Brazil) 193
- Ethanol production from mixtures of sweet sorghum juice and sorghum starch using very high gravity fermentation with urea supplementation
N.B. Appiah-Nkansah, K. Zhang, W. Rooney and D. Wang (United States) 247
- Economic viability of energy beets (*Beta vulgaris*) as advanced biofuel feedstocks
A.V. De Laporte and D.G. Ripplinger (United States) 254
- Smart utilization of jatropha (*Jatropha curcas* Linnaeus) seeds for biodiesel production: Optimization and mechanism
D.A. Kamel, H.A. Farag, N.K. Amin, A.A. Zatout and R.M. Ali (Egypt) 407
- Investigation of dynamic changes of substrate features on enzymatic hydrolysis of lignocellulosic biomass
X. Li and Y. Zheng (USA) 414
- Steam explosion of sweet sorghum stems: Optimisation of the production of sugars by response surface methodology combined with the severity factor
J. Damay, X. Duret, T. Ghislain, O. Lalonde and J.-M. Lavoie (Canada) 482
- Does straw mulch partial-removal from soil interfere in yield and industrial quality sugarcane? A long term study
G.S. de Aquino, C. de Conti Medina, M. Shahab, A.D. Santiago, A.C.B. Cunha, D.A.O. Kussaba, J.B. Carvalho and A. Moreira (Brazil) 573
- Comparative study on hydrolysis and bioethanol production from cardoon and rockrose pretreated by dilute acid hydrolysis
M.C. Fernandes, M.D. Ferro, A.F.C. Paulino, H.T. Chaves (Portugal), D.V. Evtuguin (Portugal, Russian Federation) and A.M.R.B. Xavier (Portugal) 633
- Physicochemical characterization of Malaysian crop and agro-industrial biomass residues as renewable energy resources
S. Sohni, N.A.N. Norulaini, R. Hashim (Malaysia), S.B. Khan (Saudi Arabia), W. Fadhillah and A.K. Mohd Omar (Malaysia) 642
- High sugar yields from sugarcane (*Saccharum officinarum*) bagasse using low-temperature aqueous ammonia pretreatment and laccase-mediator assisted enzymatic hydrolysis
K. Raj and C. Krishnan (India) 673
- Role of hemicellulose removal during dilute acid pretreatment on the cellulose accessibility and enzymatic hydrolysis of compositionally diverse sugarcane hybrids
V.T. de Oliveira Santos, G. Siqueira, A.M.F. Milagres and A. Ferraz (Brazil) 722
- Sugarcane straw removal effects on plant growth and stalk yield
I.P. Lisboa (Brazil, USA), M.R. Cherubin, R.P. Lima, C.C. Cerri, L.S. Satiro (Brazil), B.J. Wienhold, M.R. Schmer, V.L. Jin (USA) and C.E.P. Cerri (Brazil) 794
- Classification of biomass through their pyrolytic bio-oil composition using FTIR and PCA analysis
E. Lazzari, T. Schena, M.C.A. Marcelo, C.T. Primaz, A.N. Silva, M.F. Ferrão, T. Bjerck (Brazil) and E.B. Caramão (Brazil) 856
- ### Bio-based Materials from Crops
- Superparamagnetic Fe₃O₄@ wood flour/ polypropylene nanocomposites: Physical and mechanical properties
M. Mashkour and Y. Ranjbar (Iran) 47
- High-temperature, low-temperature and weathering aging performance of lignin modified asphalt binders
K.B. Batista, R.P.L. Padilha, T.O. Castro, C.F.S.C. Silva, M.F.A.S. Araújo, L.F.M. Leite, V.M.D. Pasa and V.F.C. Lins (Brazil) 107
- (UV/Oxidative) dual curing polyurethane dispersion from cardanol based polyol: Synthesis and characterization
V. Mishra, J. Desai and K.I. Patel (India) 165
- Application of guar gum for the removal of dissolved lead from wastewater
S. Mukherjee (Ireland), S. Mukhopadhyay (UK), M.Z.B. Zafri (Malaysia), X. Zhan (Ireland), M.A. Hashim (Malaysia) and B.S. Gupta (UK) 261
- Utilizing *Miscanthus* stalks as raw material for particleboards
P. Klímek (Germany, Czech Republic), R. Wimmer (Austria), P. Meinschmidt (Germany) and J. Kúdela (Slovakia) 270
- Impacts of hemp fiber diameter on mechanical and water uptake properties of polybenzoxazine composites
A.Q. Dayo (PR China, Pakistan), A.-r. Wang, S. Kiran, J. Wang (PR China), K. Qureshi (Pakistan), Y.-l. Xu, A. Zegaoui, M. Derradji (PR China), A.A. Babar (Pakistan) and W.-b. Liu (PR China) 277

- Lippia origanoides* essential oil: An efficient alternative to control *Aedes aegypti*, *Tetranychus urticae* and *Cerataphis lataniae*
J.M. Mar, L.S. Silva, S.G. Azevedo, L.P. França, A.F.F. Goes, A.L. dos Santos, J. de A. Bezerra, R. de Cássia S. Nunomura, M.B. Machado and E.A. Sanches (Brazil) 625
- 292 Recovery of yerba mate (*Ilex paraguariensis*) residue for the development of PLA-based bionanocomposite films
M.P. Arrieta, L. Peponi, D. López and M. Fernández-García (Spain) 667
- 317 Effects of water deficit and rehydration on antioxidant and anti-inflammatory activities in methanolic root barks extracts of *Rhus tripartitum* and *Periploca laevigata* subsp. *angustifolia*
S. Ncib, M. Boukhris, E. Lefi (Tunisia), S. Adesso, G. Autore, S. Marzocco (Italy) and B. Hanchi (Tunisia) 743
- 353 Modification of hemp fibers (*Cannabis Sativa* L.) for composite applications
T. Väisänen, P. Batello, R. Lappalainen and L. Tomppo (Finland) 846
- 422 One-pot solvent-free synthesis and characterisation of hydroxypropylated polyflavonoid compounds
J.H. Bridson, W.J. Grigsby and L. Main (New Zealand) 865
- 529 Development of enhanced hypericin yielding transgenic plants and somaclones: High throughput direct organogenesis from leaf and callus explants of *Hypericum perforatum*
S.A. Khan, P. Verma, A. Arbat, S. Gaikwad and V.A. Parasharami (India) 878
- 544 Surfactant-mediated hydrothermal pretreatment of Ryegrass followed by enzymatic saccharification for polyhydroxyalkanoate production
R. Kataria, T. Woods, W. Casey, F. Cerrone, R. Davis, K. O'Connor (Ireland), R. Ruhai (Belgium) and R. Babu (Ireland) 908
- 422 Biochar-organic amendment mixtures added to simulated golf greens under reduced chemical fertilization increase creeping bentgrass growth
S.F. Vaughn, F. Dan Dinelli, M.A. Jackson, M.M. Vaughan and S.C. Peterson (USA) 667
- 317 Evaluation of competition, essential oil quality and quantity of peppermint intercropped with soybean
M.A. Machiani, A. Javanmard, M.R. Morshedloo (Iran) and F. Maggi (Italy) 743
- 353 Encapsulation efficiency and thermal stability of norbixin microencapsulated by spray-drying using different combinations of wall materials
D.S. Tupuna, K. Paese, S.S. Guterres, A. Jablonski, S.H. Flôres and A.d.O. Rios (Brazil) 846
- 422 Agronomic performance of the novel oilseed crop *Euphorbia lagascae* Spreng., (Euphorbiaceae) in southwestern Ontario
S. Chakraborty, J. Todd (Canada), T. Isbell (USA) and R.C. Van Acker (Canada) 865
- 529 Compatibilization of highly sustainable polylactide/almond shell flour composites by reactive extrusion with maleinized linseed oil
L. Quiles-Carrillo, N. Montanes (Spain), C. Sammon (UK), R. Balart and S. Torres-Giner (Spain) 878
- 544 Consequence on long term storage on phytochemical attributes of cumin (*Cuminum cyminum*, L.) from districts of north Gujarat, India
S.B. Gondaliya, T.J. Khatrani, K.K. Soni and Y.K. Baravalia (India) 908