

## Contents

(Abstracts/contents list published in *Biological Abstracts*, *Biological & Agricultural Index*, *Current Advances in Biological Sciences*, *Current Contents AB & ES*, *Ecological Abstracts*, *EMBIology*, *Environmental Abstracts*, *Field Crop Abstracts*, *Geo Abstracts*, *Geobase*, *Geographical Abstracts–Economic Geography*, *TROPAG* and *RURAL* database, *Science Citation Index*). Also covered in the abstract and citation database SCOPUS®. Full text available on ScienceDirect®.

### Reviews

The residual value of fertiliser N in crop sequences: An appraisal of 60 years of research using <sup>15</sup>N tracer  
C.J. Smith and P.M. Chalk (Australia)

66

Comparison of transgenic *Bt* rice and their non-*Bt* counterpart in yield and physiological response to drought stress

Y. Jiang, L. Ling, L. Zhang, K. Wang, X. Li, M. Cai, M. Zhan, C. Li, J. Wang and C. Cao (PR China)

45

### Research articles

Wheat lines exhibiting variation in tolerance of *Septoria tritici* blotch differentiated by grain source limitation  
F. Collin (United Kingdom), P. Bancal (France), J. Spink, P.K. Appelgren (Ireland), J. Smith, N.D. Paveley (United Kingdom), M.-O. Bancal (France) and M.J. Foulkes (United Kingdom)

1

Resistance and tolerance to the brown planthopper, *Nilaparvata lugens* (Stål), in rice infested at different growth stages across a gradient of nitrogen applications  
F.G. Horgan (Australia), A. Peñalver Cruz (Philippines, UK), C.C. Bernal, A.F. Ramal, M.L.P. Almazan (Philippines) and A. Wilby (UK)

53

Canopy temperature depression at grain filling correlates to winter wheat yield in the U.S. Southern High Plains  
S. Thapa, K.E. Jessup, G.P. Pradhan, J.C. Rudd, S. Liu, J.R. Mahan, R.N. Devkota, J.A. Baker and Q. Xue (USA)

11

Localized ammonium and phosphorus fertilization can improve cotton lint yield by decreasing rhizosphere soil pH and salinity

X.-X. Wang, S. Liu, S. Zhang, H. Li, B. Maimaitiaili, G. Feng (China) and Z. Rengel (Australia)

75

Is mulching an efficient way to control weeds? Effects of type and amount of crop residue in rainfed rice based cropping systems in Madagascar

L. Ranaivoson (Madagascar, France), K. Naudin (France, Brazil), A. Ripoche (France, Madagascar), L. Rabeharisoa (Madagascar) and M. Corbeels (France, Kenya)

20

Evaluating canopy spectral reflectance vegetation indices to estimate nitrogen use traits in hard winter wheat  
K. Frels, M. Guttieri, B. Joyce, B. Leavitt and P.S. Baenziger (United States)

82

QTL for stay-green traits in wheat in well-watered and water-limited environments

M. Christopher, K. Chenu, R. Jennings, S. Fletcher, D. Butler, A. Borrell and J. Christopher (Australia)

32

Quantifying the response of wheat yields to heat stress: The role of the experimental setup

E.E. Rezaei, S. Siebert, R. Manderscheid, J. Müller, A. Mahrookashani, B. Ehrenpfordt, J. Haensch, H.-J. Weigel and F. Ewert (Germany)

93

(Contents continued on IBC)

(Contents continued from OBC)

|                                                                                                                                                                                                                                                                                                    |     |                                                                                                                                                                                                                                     |     |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| Penalties in yield and yield associated traits caused by stem lodging at different developmental stages in summer and spring foxtail millet cultivars<br>B. Tian, S. Luan, L. Zhang, Y. Liu, L. Zhang and H. Li (China)                                                                            | 104 | Forage rape ( <i>Brassica napus</i> L) seed quality: Impact of heat stress in the field during seed development<br>M. Rashid (New Zealand, Pakistan), J.G. Hampton, M.P. Rolston, J.A.K. Trethewey and D.J. Saville (New Zealand)   | 172 |
| Application of the APSIM model to exploit G × E × M interactions for maize improvement in Ethiopia<br>S. Seyoum, R. Rachaputi, Y. Chauhan (Australia), B. Prasanna (Kenya) and S. Fekybelu (Australia)                                                                                             | 113 | Shifting crop-pasture rotations to no-till annual cropping reduces soil quality and wheat yield<br>O.R. Ernst, S. Dogliotti, M. Cadenazzi (Uruguay) and A.R. Kemanian (USA)                                                         | 180 |
| Optimised sowing date enhances crop resilience towards size-asymmetric competition and reduces the yield difference between intercropped and sole maize<br>C. Huang, Q. Liu, H. Li, X. Li, C. Zhang and F. Zhang (China)                                                                           | 125 | Impact of the insecticide application to maize cultivated in different environmental conditions on emerging mycotoxins<br>V. Scarpino, A. Reyneri (Italy), M. Sulyok, R. Krska (Austria) and M. Blandino (Italy)                    | 188 |
| CERES-Maize and CERES-Sorghum for modeling growth, nitrogen and phosphorus uptake, and soil moisture dynamics in the dry savanna of West Africa<br>K.A. Amouzou (Germany, Burkina Faso), J.B. Naab (Burkina Faso), J.P.A. Lamers and M. Becker (Germany)                                           | 134 | In search of long-term sustainable tillage and straw mulching practices for a maize-winter wheat-soybean rotation system in the Loess Plateau of China<br>Z. Li, X. Lai, Q. Yang, X. Yang (China), S. Cui (USA) and Y. Shen (China) | 199 |
| Testing pearl millet and cowpea intercropping systems under high temperatures<br>W.C.D. Nelson, M.P. Hoffmann (Germany), V. Vadez (India), R.P. Roetter (Germany) and A.M. Whitbread (India)                                                                                                       | 150 | Spread and yield loss mechanisms of rice stripe disease in rice paddies<br>T. Shiba, M. Hirae, Y. Hayano-Saito, Y. Ohto, H. Uematsu, A. Sugiyama and M. Okuda (Japan)                                                               | 211 |
| A study on the productivity under the continuous maize cultivation in Sainyabuli Province, Laos I. Yield trend under continuous maize cultivation<br>K. Fujisao (Japan), P. Khanthavong, S. Oudthachit (Lao Democratic People's Republic), N. Matsumoto, K. Homma, H. Asai and T. Shiraiwa (Japan) | 167 | Evapotranspiration, crop coefficient and yield for drip-irrigated winter wheat with straw mulching in North China Plain<br>J. Wang, Y. Zhang, S. Gong, D. Xu, S. Juan and Y. Zhao (PR China)                                        | 218 |