

INDUSTRIAL CROPS AND PRODUCTS

VOLUME 112

February 2018

Contents

Fats and oils

- Improvement of biomass and essential oil production of *Lippia alba* (Mill) N.E. Brown with green manures in succession
C.T. dos Santos Marques, E.V.S. Gama, F. da Silva, S. Teles, A.N. Caiafa and A.M. Lucchese (Brazil) 113
- Analysis of phytochemical and morphological variability in different wild- and agro-ecotypic populations of *Melissa officinalis* L. growing in northern habitats of Iran
E. Pouyanfar, J. Hadian, M. Akbarzade, M. Hatami, M.R. Kanani and M. Ghorbanpour (Iran) 262
- Chemical characterization of the essential oil compositions and antioxidant activity from Iranian populations of *Achillea wilhelmsii* K.Koch
K. Saeidi, M. Moosavi, Z. Lorigooini (Iran) and F. Maggi (Italy) 274
- Phytochemical and morphological variation of *Stachys lavandulifolia* Vahl. populations as affected by genotype x year interaction
F. Arabsalehi, M. Rahimmalek and M.H. Ehtemam (Iran) 342
- Changes in the essential oil content and composition of *Thymus daenensis* Celak. under different drying methods
M.R. Dehghani Mashkani, K. Larijani, A. Mehrafarin and H. Naghdi Badi (Iran) 389
- Mentha x piperita* L. nodal segments cultures and their essential oil production
J. Fejér, D. Gruťová (Slovak Republic), V. De Feo (Italy)E. Űrgeová, B. Obert and A. Preťová (Slovak Republic) 550
- Biofuels production by thermal cracking of soap from brown grease
A.R. Ratton Coppos, S. Kahn and L.E.P. Borges (Brazil) 561
- Correlation and Path analysis for yield components in Dura oil palm germplasm
S. de Almeida Rios, R.N. Vieira da Cunha, R. Lopes, E. Barcelos, Raimundo Nonato Carvalho da Rocha and W.A. Alves de Lima (Brazil) 724
- Zingiber zerumbet* (L.) Roscoe ex Sm. from northern India: Potential source of zerumbone rich essential oil for antiproliferative and antibacterial applications
R.C. Padalia, R.S. Verma, A. Chauhan, V.R. Singh, P. Goswami, S. Singh, S.K. Verma, S. Luqman, C.S. Chanotiya and M.P. Darokar (India) 749
- Trait-specific accessions in global castor (*Ricinus communis* L.) germplasm core set for utilization in castor improvement
K. Anjani, M.A. Raoof, M.S.L. Prasad, P. Duraimurugan, C. Lucose, P. Yadav, R.D. Prasad, J.J. Lal and C. Sarada (India) 766

Non-food bioactive products

- Micronization of trans-resveratrol by supercritical fluid: Dissolution, solubility and *in vitro* antioxidant activity
G.P.S. Aguiar, B.D. Arcari, L.M.P.C. Chaves, C.D. Magro, D.L. Boschetto, A.L. Piato, M. Lanza and J.V. Oliveira (Brazil)
- Enzyme inhibitory and antioxidant activities of *Nerium oleander* L. flower extracts and activity guided isolation of the active components
I. Atay Balkan, H.T. Doğan, G. Zengin, N. Colak, F.A. Ayaz, A.C. Gören, H. Kirmizibekmez and E. Yeşilada (Turkey)
- Phenolic profile, chemical relationship and antifungal activity of Andean *Hypericum* species
N. Tocci, T. Weil, D. Perenzoni, L. Narduzzi (Italy), S. Madriñán (Colombia), S. Crockett (Austria), N.M. Nürk (Germany)D. Cavaliere and F. Mattivi (Italy)
- Essential oil of *Eucalyptus camaldulensis* Dehn potentiates -lactam activity against *Staphylococcus aureus* and *Escherichia coli* resistant strains
T.P. Chaves, R.E.E. Pinheiro, E.S. Melo, M.J.d.S. Soares, J.S.N. Souza, T.B.d. Andrade, T.L.G.d. Lemos and H.D.M. Coutinho (Brazil)
- Bioactive compounds of organic goji berry (*Lycium barbarum* L.) prevents oxidative deterioration of soybean oil
A.C. Pedro, J.B.B. Maurer, S.F. Zawadzki-Baggio, S. Ávila, G.M. Maciel and C.W.I. Haminiuk (Brazil)
- Expression of artemisinin biosynthesis and trichome formation genes in five *Artemisia* species
M. Salehi, G. Karimzadeh, M.R. Naghavi, H. Naghdi Badi and S. Rashidi Monfared (Iran)
- Effects of manganese on accumulation of Glycyrrhizic acid based on material ingredients distribution of *Glycyrrhiza uralensis*
S. Ma, G. Zhu, F. Yu (China), G. Zhu (Urumqi 830011,China)D. Wang, W. Wang and J. Hou (China)
- The non-edible parts of fennel (*Foeniculum vulgare*) as a new milk-clotting protease source
N. Bey, H. Debbebi, F. Abidi, M.N. Marzouki and A. Ben Salah (Tunisia)
- Chemical composition, antioxidant and anticancer activity of licorice from Fruska Gora locality
S. Vlasisavljević, F. Šibul (Serbia), I. Sinka, I. Zupko, I. Ocsovszki (Hungary) and S. Jovanović-Šanta (Serbia)
- 1 Remarkable enhancement of flavonoid production in a co-cultivation system of *Isatis tinctoria* L. hairy root cultures and immobilized *Aspergillus niger*
J. Jiao, Q.-Y. Gai, W. Wang, Y.-P. Zang, L.-L. Niu, Y.-J. Fu and X. Wang (PR China)
- 24 Cinnamon oil-loaded composite emulsion hydrogels with antibacterial activity prepared using concentrated emulsion templates
J. Wang, Y. Li, Y. Gao, Z. Xie, M. Zhou, Y. He, H. Wu, W. Zhou, X. Dong, Z. Yang and Y. Hu (China)
- 32 Ultrasound assisted extraction of polyphenolic compounds from red sorghum (*Sorghum bicolor* L.) bran and their biological activities and polyphenolic compositions
X. Luo, J. Cui, H. Zhang, Y. Duan, D. Zhang, M. Cai and G. Chen (China)
- 70 Anti-inflammatory activity and phenolic composition of prickly pear (*Opuntia ficus-indica*) flowers
I. Ammar, M. Ben Salem, B. Harrabi, M. Mzid, S. Bardaa, Z. Sahnoun, H. Attia and M. Ennouri (Tunisia)
- 90 Isolation of essential oil from the leaves of *Polygonum viscosum* Buch-ham. using microwave-assisted enzyme pretreatment followed by microwave hydrodistillation concatenated with liquid-liquid extraction
F. Chen, J. Jia, Q. Zhang, L. Yang and H. Gu (China)
- 130 Chemical diversity, antioxidant and antimicrobial activities of the essential oils from Indian populations of *Hedychium coronarium* Koen
A. Ray, S. Jena, B. Dash, B. Kar, T. Halder, T. Chatterjee, B. Ghosh, P.C. Panda, S. Nayak and N. Mahapatra (India)
- 151 Yield and cannabinoid contents in different cannabis (*Cannabis sativa* L.) genotypes for medical use
A. Janatová, A. Fraňková, P. Tlustoš, K. Hamouz, M. Božik and P. Klouček (Czech Republic)

Foam fractionation for effective recovery of resveratrol from the leaching liquor of <i>Polygonum cuspidatum</i> by using partially ethylated β -cyclodextrin as collector and frother <i>W. Liu, Y. Lv, S. Tian, M. Zhang, T. Shu, N. Hu and Z. Wu (China)</i>	420	Biological activity of <i>Hyptis</i> Jacq. (Lamiaceae) is determined by the environment <i>K.P. dos Santos, M.D. Sedano-Partida, W.R. Sala-Carvalho, B.O.S.J. Loureiro, C.L. da Silva-Luz and C.M. Furlan (Brazil)</i>	705
Phytochemical analysis and assessment of antioxidant, antimicrobial, anti-inflammatory and cytotoxic properties of <i>Tetraclinis articulata</i> (Vahl) Masters leaves <i>W. Rached (Portugal, Algeria), F.Z. Zeghada, M. Bennaceur (Algeria), L. Barros, R.C. Calhelha, S. Heleno, M.J. Alves, A.M. Carvalho (Portugal), A. Marouf (Algeria) and I.C.F.R. Ferreira (Portugal)</i>	460	Effect of foliar nutrition on the essential oil yield of Thyme (<i>Thymus vulgaris</i> L.) <i>R. Pavela, M. Žabka, N. Vrchatová and J. Tříška (Czech Republic)</i>	762
Morphoanatomy and essential oil analysis of <i>Baccharis trimera</i> (Less.) DC. (Asteraceae) from Uruguay <i>M. Minteguiaga (Uruguay), M.I. Mercado, G.I. Ponessa, C.A.N. Catalán (Argentina) and E. Dellacassa (Uruguay)</i>	488	Cultivar affects the phenolic composition and antioxidant properties of commercially available lemon balm (<i>Melissa officinalis</i> L.) varieties <i>M.M. Boneza and E.D. Niemeyer (USA)</i>	783
Plant phenolic extracts as an effective strategy to control <i>Staphylococcus aureus</i> , the dairy industry pathogen <i>F. Gomes, N. Martins, L. Barros, M.E. Rodrigues, M.B.P.P. Oliveira, M. Henriques and I.C.F.R. Ferreira (Portugal)</i>	515	The synergistic effects of volatile constituents of <i>Ocimum basilicum</i> against foodborne pathogens <i>J.L. Baldim, J.G. Fernandes Silveira, A.P. Almeida, P.L.N. Carvalho, W. Rosa, J. Schripsema, D.A. Chagas-Paula, M.G. Soares and J.H.H. Luiz (Brazil)</i>	821
Conventional octaplex PCR for the simultaneous identification of eight mainstream closely related <i>Dendrobium</i> species <i>C. Jiang, Y. Luo, Y. Yuan, X. Dong, Y. Zhao and L. Huang (PR China)</i>	569	Elucidation of micromolecular phenylpropanoid and lignan glycosides as the main antioxidants of Ginkgo seeds <i>S.-j. Shan, J. Luo, D.-r. Xu, X.-l. Niu, D.-q. Xu, P.-p. Zhang and L.-y. Kong (People's Republic of China)</i>	830
Extraction optimization for polyphenolic profiling and bioactive enrichment of extractives of non-pomace residue from grape processing <i>I.C.d.S. Haas, I.M. Toaldo, V.M. Burin and M.T. Bordignon-Luiz (Brazil)</i>	593	Natural rubber and resins	
Evaluation of phytochemicals in <i>Agastache rugosa</i> (Fisch. & C.A.Mey.) Kuntze at different growth stages by UPLC-QToF-MS <i>J.H. An, H.J. Yuk, D.-Y. Kim, C.W. Nho, D. Lee, H.W. Ryu and S.-R. Oh (Republic of Korea)</i>	608	Colchicine-induced polyploidy has the potential to improve rubber yield in <i>Taraxacum kok-saghyz</i> <i>Z. Luo, B.J. Iaffaldano and K. Cornish (USA)</i>	75
Acute and sub-lethal toxicity of eight essential oils of commercial interest against the filariasis mosquito <i>Culex quinquefasciatus</i> and the housefly <i>Musca domestica</i> <i>G. Benelli (Italy), R. Pavela (Czech Republic), C. Giordani (Colombia), L. Casettari, G. Curzi, L. Cappellacci, R. Petrelli and F. Maggi (Italy)</i>	668	Phenotypic variations, heritability and correlations in dry biomass, rubber and resin production among guayule improved germplasm lines <i>H. Abdel-Haleem, M. Foster, D. Ray and T. Coffelt (USA)</i>	691
		Overexpression of <i>EcGSH1</i> induces glutathione production and alters somatic embryogenesis and plant development in <i>Hevea brasiliensis</i> <i>F. Martin, V. Abati, A. Burel, A. Clément-Vidal, C. Sanier, D. Fabre (France), N. Woraathasin (Thailand), M. Rio, P. Besret, B. Farinas, P. Montoro and J. Leclercq (France)</i>	803
		Fibres, proteins and carbohydrates	
		Antioxidant and anticoagulant activities of mycelia polysaccharides from <i>Catathelasma ventricosum</i> after sulfated modification <i>Y. Liu, Q. Tang, X. Duan, T. Tang, Y. Ke, L. Zhang, C. Li, A. Liu, Z. Su and B. Hu (China)</i>	53

Glucose-responsive biopolymer nanoparticles prepared by co-assembly of concanavalin A and amylopectin for insulin delivery <i>R. Chang, M. Li, S. Ge, J. Yang, Q. Sun and L. Xiong (China)</i>	98	Jute cellulose nano-fibrils/hydroxypropylmethylcellulose nanocomposite: A novel material with potential for application in packaging and transdermal drug delivery system <i>J.T. Orasugh, N.R. Saha (India), D. Rana (Canada)G. Sarkar, Md.M.R. Mollick, A. Chattoapadhyay, B.C. Mitra, D. Mondal, S.K. Ghosh and D. Chattopadhyay (India)</i>	633
Differential response of tossa jute (<i>Corchorus olitorius</i>) submitted to water deficit stress <i>P. Dhar, D. Ojha, C.S. Kar and J. Mitra (India)</i>	141		
Encapsulation templated approach to valorization of cocoa husk, poppy and hemp macrostructural and bioactive constituents <i>A. Belščak-Cvitanović, A. Vojvodić, A. Bušić (Croatia), J. Keppler, A. Steffen-Heins (Germany) and D. Komes (Croatia)</i>	402	Biorefinery, valorization of byproducts Multi-valorisation of giant reed (<i>Arundo Donax</i> L.) to give levulinic acid and valuable phenolic antioxidants <i>D. Licursi, C. Antonetti, M. Mattonai (Italy), L. Pérez-Armada, S. Rivas (Spain)E. Ribechini and A.M. Raspolli Galletti (Italy)</i>	6
Soy protein-based films incorporated with cellulose nanocrystals and pine needle extract for active packaging <i>Z. Yu, L. Sun, W. Wang (USA), W. Zeng (PR China)A. Mustapha and M. Lin (USA)</i>	412	Utilisation of supercritical fluids for the effective extraction of waxes and Cannabidiol (CBD) from hemp wastes <i>T.M. Attard, C. Bainier, M. Reinaud, A. Lanot, S.J. McQueen-Mason (UK) and A.J. Hunt (UK, Thailand)</i>	38
Biodegradable trays of thermoplastic starch/poly (lactic acid) coated with beeswax <i>M.O. Reis, J.B. Olivato, A.P. Bilck, J. Zanela, M.V.E. Grossmann and F. Yamashita (Brazil)</i>	481	Techno-economic analysis of guayule (<i>Parthenium argentatum</i>) pyrolysis biorefining: Production of biofuels from guayule bagasse via tail-gas reactive pyrolysis <i>P.S. Sabaini (USA, Brazil)A.A. Boateng, M. Schaffer, C.A. Mullen, Y. Elkasabi, C.M. McMahan and N. Macken (USA)</i>	82
Alfa fibers as viable sustainable source for cellulose nanocrystals extraction: Application for improving the tensile properties of biopolymer nanocomposite films <i>M. El Achaby, Z. Kassab (Morocco), A. Barakat (France) and A. Aboulkas (Morocco)</i>	499	Membrane filtration of kraft lignin: Structural characteristics and antioxidant activity of the low-molecular-weight fraction <i>S. Aminzadeh (Sweden), M. Lauberts, G. Dobele, J. Ponomarenko (Latvia)T. Mattsson, M.E. Lindström and O. Sevastyanova (Sweden)</i>	200
Characterization of coated biodegradable trays by spectroscopic techniques <i>M.O. Reis, H. de Santana, A.P. Bilck, M.V.E. Grossmann and F. Yamashita (Brazil)</i>	511	Evaluation of Ligno Boost™ softwood kraft lignin epoxidation as an approach for its application in cured epoxy resins <i>A. Jablonskis, A. Arshanitsa, A. Arnautov, G. Telysheva (Latvia) and D. Evtuguin (Portugal)</i>	225
Study of Schiff base formation between dialdehyde cellulose and proteins, and its application for the deproteinization of crude polysaccharide extracts <i>L. Zhang, Q. Zhang, Y. Zheng, Z. He, P. Guan, X. He, L. Hui and Y. Dai (People's Republic of China)</i>	532	Influence of process conditions during aqueous protein extraction upon yield from pre-pressed and cold-pressed rapeseed press cake <i>A. Fetzer, T. Herfellner, A. Stäbler, M. Menner and P. Eisner (Germany)</i>	236
Nanocrystals of cellulose allomorphs have different adsorption of cellulase and subsequent degradation <i>Z. Ling, X. Zhang, G. Yang (China), K. Takabe (Japan) and F. Xu (China)</i>	541	Formic acid reinforced autohydrolysis of wheat straw for high yield production of monosugars and minimal lignin precipitation <i>Ö. Özyürek (Turkey, USA) and A. van Heiningen (USA)</i>	320
<i>Curcuma longa</i> L. plant residue as a source for natural cellulose fibers with antimicrobial activity <i>M. Ilangoan, V. Guna (India), C. Hu (PR China)G.S. Nagananda and N. Reddy (India)</i>	556		

- Revalorization of winery by-products as source of natural preservatives obtained by means of green extraction techniques
J.M. Poveda, L. Loarce, M. Alarcón, M.C. Díaz-Maroto and M.E. Alañón (Spain) 617
- Effect of the organizational difference of corn stalk on hemicellulose extraction and enzymatic hydrolysis
J. Zhang, Y.-H. Wang, Y.-S. Qu, Q.-Y. Wei and H.-Q. Li (China) 698
- Enhancing enzymatic hydrolysis of green coconut fiber—Pretreatment assisted by tween 80 and water effect on the post-washing
C. da Costa Nogueira, C.E. de Araújo Padilha, A.L. de Sá Leitão, P.M. Rocha, G.R. de Macedo and E.S. dos Santos (Brazil) 734
- Production of resveratrol in *Flammulina velutipes* by combinatorial expression of its biosynthetic genes
X. Yang, Q. Zheng, L. Kang, Q. Li, Z. Ye, J. Lin, L. Guo and T. Wei (China) 755
- From waste to wealth: High recovery of nutraceuticals from pomegranate seed waste using a green extraction process
S. Talekar (India), A.F. Patti (Australia), R. Singh (India), R. Vijayraghavan (Australia) and A. Arora (India) 790
- Bioenergy - crops and processing**
- An effective pathway for converting carbohydrates to biofuel 5-ethoxymethylfurfural via 5-hydroxymethylfurfural with deep eutectic solvents (DESs)
M. Zuo, K. Le, Y. Feng, C. Xiong, Z. Li, X. Zeng, X. Tang, Y. Sun and L. Lin (PR China) 18
- Charcoal rot and Fusarium stalk rot diseases influence sweet sorghum sugar attributes
Y.M.A.Y.. Bandara, T.T. Tesso, K. Zhang, D. Wang and C.R. Little (USA) 188
- Application of Fenton pretreatment on the degradation of rice straw by mixed culture of *Phanerochaete chrysosporium* and *Aspergillus niger*
Y. Chen, Y. Chen, Y. Li, Y. Wu, F. Zhu, G. Zeng, J. Zhang and H. Li (China) 290
- Ethanol biosynthesis by fast hydrolysis of cassava bagasse using fungal amylases produced in optimized conditions
B. Escaramboni, E.G. Fernández Núñez, A.F.A. Carvalho and P. de Oliva Neto (Brazil) 368
- Mechanical properties of cotton shoots for topping
I. Aydin and S. Arslan (Turkey) 396
- Treatments to improve obtention of reducing sugars from agave leaves powder
J. Avila-Gaxiola, O.J. Velarde-Escobar, J.R. Millan-Almaraz, F. Ramos-Brito, G. Atondo-Rubio, C. Yee-Rendon and E. Avila-Gaxiola (Mexico) 577
- Inbreeding depression as a cause of fruit abortion in structured populations of macaw palm (*Acrocomia aculeata*): Implications for breeding programs
G.F. Simiqueli, M.D.V.d. Resende, S.Y. Motoike and E. Henriques (Brazil) 652
- Optimization of shelling efficiency of a *Moringa oleifera* seed shelling machine based on seed sizes
O.K. Fadele and A.K. Aremu (Nigeria) 775
- Developing predictive models for determining physical properties of coffee beans during the roasting process
J.D. Bustos-Vanegas, P.C. Corrêa, M.A. Martins, F.M. Baptestini, R.C. Campos, G.H.H. de Oliveira and E.H.M. Nunes (Brazil) 839
- Bio-based Materials from Crops**
- Effect of a new shell material—Jackfruit seed starch on novel flavor microcapsules containing vanilla oil
H. Zhu, Y. Zhang, J. Tian and Z. Chu (China) 47
- Study on the UV curing behavior of tung oil: Mechanism, curing activity and film-forming property
J. Huang, T. Yuan, X. Ye, L. Man, C. Zhou, Y. Hu, C. Zhang and Z. Yang (China) 61
- Plant tonic, a plant-derived bioactive natural product, exhibits antifungal activity against rice blast disease
F. Abed-Ashtiani (Iran, Malaysia), M. Arzanlou (Iran), A. Nasehi, J. Kadir, G. Vadamalai (Malaysia) and S. Azadmard-Damirchi (Iran) 105
- Rapeseed-based polyols and paper production waste sludge in polyurethane foam: Physical properties and their prediction models
A. Kairyte, S. Vaitkus, S. Vėjelis, G. Girskas and G. Balčiūnas (Lithuania) 119
- Substitution of peat moss with softwood biochar for soil-free marigold growth
A.J. Margenot, D.E. Griffin, B.S.Q. Alves, D.A. Rippner, C. Li and S.J. Parikh (United States) 160

Preparation and characterization of poly(lactic acid) plasticized with vegetable oils and reinforced with sisal fibers <i>A. Orue, A. Eceiza and A. Arbelaiz (Spain)</i>	170	Totora (<i>Schoenoplectus californicus</i> (C.A. Mey.) Soják) and its potential as a construction material <i>J.F. Hidalgo-Cordero and J. García-Navarro (Spain)</i>	467
Coating applications to natural fiber composites to improve their physical, surface and water absorption characters <i>Z. Liu and B.H. Tisserat (United States)</i>	196	Phytochemical characterization, antioxidant, antimicrobial and pharmacological activities of <i>Feijoa sellowiana</i> leaves growing in Tunisia <i>H. Mosbah, H. Louati, M.A. Boujbiha, H. Chahdoura, M. Snoussi (Tunisia), G. Flamini, R. Ascrizzi (Italy)A. Bouslema, L. Achour and B. Selmi (Tunisia)</i>	521
Water-resistant hemp fiber-reinforced composites: <i>In-situ</i> surface protection by polyethylene film <i>Y. Wu, C. Xia, L. Cai, S.Q. Shi and J. Cheng (USA)</i>	210	Metabolomic analysis by UAE-GC MS and antioxidant activity of <i>Salvia hispanica</i> (L.) seeds grown under different irrigation regimes <i>B. de Falco (Italy, UK), A. Fiore (UK)R. Bochicchio, M. Amato and V. Lanzotti (Italy)</i>	584
Characterization of leaf cuticular wax classes and constituents in a spring <i>Camelina sativa</i> diversity panel <i>P. Tomasi, J.M. Dyer, M.A. Jenks and H. Abdel-Haleem (USA)</i>	247	Testing composted bamboo residues with and without added effective microorganisms as a renewable alternative to peat in horticultural production <i>Z. Zhong, F. Bian and X. Zhang (PR China)</i>	602
Synergistic photoprotective activity of nanocarrier containing oil of <i>Acrocomia aculeata</i> (Jacq.) Lodd. Ex. Martius—Arecaceae <i>M.F. Dario, F.F. Oliveira, D.S.S. Marins, A.R. Baby, M.V.R. Velasco (Brazil), R. Löbenberg (Canada) and N.A. Bou-Chacra (Brazil)</i>	305	A greener approach towards isolating clove oil from buds of <i>Syzygium aromaticum</i> using microwave radiation <i>S.M. Kapadiya, J. Parikh and M.A. Desai (India)</i>	626
Hybrid polyether-palm oil polyester polyol based rigid polyurethane foam reinforced with cellulose nanocrystal <i>A.A. Septevani (Australia, Indonesia)D.A.C. Evans, D.J. Martin and P.K. Annamalai (Australia)</i>	378	Eco-particleboard manufactured from chemically treated fibrous vascular tissue of acai (<i>Euterpe oleracea</i> Mart.) Fruit: A new alternative for the particleboard industry with its potential application in civil construction and furniture <i>A. de Lima Mesquita (Brazil), N.G. Barrero (Colombia, Brazil)J. Fiorelli, A.L. Christoforo, L.J.G. De Faria and F.A.R. Lahr (Brazil)</i>	644
Cinnamaldehyde damaged the cell membrane of <i>Alternaria alternata</i> and induced the degradation of mycotoxins <i>in vivo</i> <i>L. Xu, N. Tao, W. Yang and G. Jing (PR China)</i>	427	Preparation and characterization of biocomposite chitosan film containing <i>Perilla frutescens</i> (L.) Britt. essential oil <i>Z.-J. Zhang, N. Li, H.-Z. Li, X.-J. Li, J.-M. Cao, G.-P. Zhang and D.-L. He (China)</i>	660
Naïve Bayes classifier, multivariate linear regression and experimental testing for classification and characterization of wheat straw based on mechanical properties <i>D.L. Naik and R. Kiran (United States)</i>	434	Detoxification of dilute ammonia pretreated energy cane bagasse enzymatic hydrolysate by soluble polyelectrolyte flocculants <i>F. Deng and G.M. Aita (United States)</i>	681
Enhancing mechanical properties of poly(lactic acid) through its in-situ crosslinking with maleic anhydride-modified cellulose nanocrystals from cottonseed hulls <i>L. Zhou (China, USA), H. He (China), M.-c. Li (USA), S. Huang, C. Mei (China) and Q. Wu (USA)</i>	449		

Flax fibres reinforced polylactide modified by ionizing radiation

*P. Rytlewski, M. Stepczyńska (Poland),
U. Gohs (Germany)R. Malinowski,
B. Budner and M. Żenkiewicz (Poland)*

716

Exogenous application of gamma-aminobutyric acid (GABA) alleviates the effect of water deficit stress in black cumin (*Nigella sativa* L.)

*E. Rezaei-Chiyaneh, S.M. Seyyedi,
E. Ebrahimian, S.S. Moghaddam (Iran) and
C.A. Damalas (Greece)*

741

Genetic and chemotypic variability in basil (*Ocimum basilicum* L.) germplasm towards future exploitation

*A. Srivastava, A.K. Gupta, S. Sarkar,
R.K. Lal, A. Yadav, P. Gupta and
C.S. Chanotiya (India)*

815

Corrigendum

Corrigendum to 'Oleoresin yield of *Pinus elliottii* in a subtropical climate: Seasonal variation and effect of auxin and salicylic acid-based stimulant paste' *Ind. Crops Prod.* 30/2 (2009) 316–320

K.C.S. Rodrigues and A.G. Fett-Neto (Brazil)

846