

## CONTENTS

|  |   |     |
|--|---|-----|
| Heat shock protein 70 (Hsp70) response to elevated temperatures in the endemic Baikal sponge <i>Lubomirskia baicalensis</i>  | V.B. Itsikovich, A.M. Shigarova, O.Y. Glyzina, O.V. Kaluzhnaya and G.B. Borovskii (Russia)  | 63  |
| Associations among taxonomic diversity, functional diversity and evolutionary distinctiveness vary among environments  | F. Morelli, Y. Benedetti (Czech Republic), P. Perna and R. Santolini (Italy)  | 71  |
| Examination of effects of acid and metals on benthic algae in streams using chemical diffusing substrata   | D.M. DeNicola, M.G. Stapleton and A.J. Onufrik (USA)  | 79  |
| $\delta^{13}\text{C}$ , $\delta^{15}\text{N}$ , N concentration, C/N, and Ca/Al of <i>Pinus densiflora</i> foliage in Korean cities of different precipitation pH and atmospheric $\text{NO}_2$ and $\text{SO}_2$ levels | H.I. Yang, H.-J. Park, K.-S. Lee, S.-S. Lim (Republic of Korea), J.-H. Kwak, S.-I. Lee, S.X. Chang (Canada), S.-M. Lee and W.-J. Choi (Republic of Korea)   | 88  |
| Transition from abstract thermodynamic concepts to perceivable ecological indicators   | H. Lin, H. Zhang and Q. Song (China)  | 103 |
| Biometrical measurements as efficient indicators to assess wild boar body condition  | D. Risco, P. Gonçalves, G. Mentaberre, N. Navarro-González, E. Casas-Díaz, D. Gassó, A. Colom-Cadena, X. Fernández-Aguilar, R. Castillo-Contreras, R. Velarde, O. Barquero-Pérez, P. Fernández-Llario, S. Lavín (Spain), C. Fonseca (Portugal) and E. Serrano (Spain, Portugal) | 115 |
| Is terrestrial water storage a useful indicator in assessing the impacts of climate variability on crop yield in semi-arid ecosystems?   | C.E. Ndehedehe (Australia, Nigeria), N.O. Agutu (Australia) and O. Okwuashi (Nigeria)   | 126 |
| Remote sensing data to assess compositional and structural indicators in dry woodland  | V.E. Campos, G.M. Gatica, F.M. Cappa, S.M. Giannoni and C.M. Campos (Argentina)   | 134 |
| Measuring the effects of temperature rise on Mediterranean shellfish aquaculture   | M. Martinez, M.C. Mangano, G. Maricchiolo, L. Genovese, A. Mazzola and G. Sarà (Italy)  | 51  |
| Short-term land use conversions influence the profile distribution of soil salinity and sodicity in northeastern China   | P. Yu, S. Liu, H. Yang, G. Fan and D. Zhou (China)  | 59  |
| Identification of indicator species at abandoned red mud dumps in comparison to residential and forest sites, accredited to soil properties  | M. Gautam, B. Pandey and M. Agrawal (India)   | 63  |
| Flow alterations by dams shaped fish assemblage dynamics in the complex Mekong-3S river system   | P.B. Ngor (Cambodia, France), P. Legendre (Canada), T. Oberdorff and S. Lek (France)  | 71  |
| Age structure as an indicator of poaching pressure: Insights from rapid assessments of elephant populations across space and time  | T. Jones (Tanzania), J.J. Cusack, R.A. Pozo (UK), J. Smit (Tanzania, UK), L. Mkuburo, P. Baran, A.L. Lobora, S. Mduma and C. Foley (Tanzania)   | 79  |
| The utility of point count surveys to predict wildlife interactions with wind energy facilities: An example focused on golden eagles   | M. Sur, J.R. Belthoff, E.R. Bjerre, B.A. Millsap and T. Katzner (USA)   | 88  |
| Uncertainty in simulating regional gross primary productivity from satellite-based models over northern China grassland  | W. Jia, M. Liu, D. Wang, H. He, P. Shi, Y. Li and Y. Wang (PR China)  | 95  |

|  |     |   |     |
|--|-----|---|-----|
| Landscape structure and composition define the body condition of dung beetles (Coleoptera: Scarabaeinae) in a fragmented tropical rainforest   |     | Stream health, topography, and land use influences on the distribution of the Eurasian otter <i>Lutra lutra</i> in the Nakdong River basin, South Korea   |     |
| R.P. Salomão, D. González-Tokman, W. Dátillo, J.C. López-Acosta and M.E. Favila (Mexico)   | 144 | S. Hong (Republic of Korea), D.-K. Kim (Canada), Y. Do, J.Y. Kim, Y.-M. Kim (Republic of Korea), P. Cowan (New Zealand) and G.-J. Joo (Republic of Korea) | 241 |
| An emergy accounting based regional sustainability evaluation: A case of Qinghai in China  |     | Development of a life-cycle impact assessment methodology linked to the Planetary Boundaries framework  |     |
| W. Chen (PR China), Y. Geng (PR China, China), H. Dong, X. Tian, S. Zhong, Q. Wu, Y. Xu, Q. Zhang and S. Li (PR China)   | 152 | M.W. Ryberg, M. Owsiania, K. Richardson and M.Z. Hauschild (Denmark)  | 250 |
| Evaluation of the use of dermal scutes and blood samples to determine organochlorine pesticides in <i>Crocodylus moreletii</i> : A non-destructive method for monitoring crocodiles and environmental health |     | Selecting plant species and traits for phytometer experiments. The case of peatland restoration   |     |
| M.G. Jáuregui, S.E. Padilla, D. Hinojosa-Garro, C. Valdespino and J.R. von Osten (Mexico)  | 161 | K. Strobl, C. Schmidt (Germany) and J. Kollmann (Germany, Norway)   | 263 |
| Temporal changes in environmental conditions of a mid-latitude estuary (southern Chile) and its influences in the cellular response of the euryhaline anemone <i>Anthopleura hermaphroditica</i>             |     | Potential invasiveness by non-indigenous macrozoobenthos in the secondary hydrographic system of a temperate-climate river catchment                      |     |
| V.M. Cubillos, E.F. Ramírez, E. Cruces, J.A. Montory, C.J. Segura and D.A. Mardones (Chile)  | 169 | D. Paganelli, A. Pandolfi, R. Sconfietti, A. Marchini (Italy) and L. Vilizzi (Poland)   | 274 |
| Thermodynamic based indicators illustrate how a run-of-river impoundment in neotropical savanna attracts invasive species and alters the benthic macroinvertebrate assemblages' complexity                   |     | Is afforestation-induced land use change the main contributor to vegetation dynamics in the semiarid region of North China?                               |     |
| M.S. Linares, M. Callisto (Brazil) and J.C. Marques (Portugal)   | 181 | F. Wang, P. An, C. Huang, Z. Zhang and J. Hao (China)   | 282 |
| Diatoms as tools for inferring ecotone boundaries in a coastal freshwater wetland threatened by saltwater intrusion  |     | Economic valuation of groundwater protection using a groundwater quality ladder based on chemical threshold levels  |     |
| V. Mazzei and E. Gaiser (USA)  | 190 | R. Brouwer (Canada, Switzerland), C.M. Ordens (Australia), R. Pinto (Canada, Portugal) and M.T.C. de Melo (Portugal)                                      | 292 |
| Identifying critical thresholds to guide management practices in agro-ecosystems: Insights from bird community response to an open grassland-to-forest gradient  |     | Comprehensive evaluation of global clean energy development index based on the improved entropy method  |     |
| P.A. Salgueiro, A. Mira, J.E. Rabaça and S.M. Santos (Portugal)  | 205 | Y.X. He, Z. Jiao and J. Yang (China)  | 305 |
| Biological surrogates: A word of caution   |     | Large river fish functional diversity responses to improved water quality over a 28 year period   |     |
| G.F. de Moraes, L.G. dos Santos Ribas, J.C.G. Ortega (Brazil), J. Heino (Finland) and L.M. Bini (Brazil)   | 214 | J. Parker, Y. Cao, G.G. Sass and J. Epifanio (USA)  | 322 |
| Updating the habitat conservation institution by prioritizing important connectivity and resilience providers outside  |     | Improved modeling of gross primary productivity (GPP) by better representation of plant phenological indicators from remote sensing using a process model |     |
| J. He, J. Huang, D. Liu, H. Wang and C. Li (China)   | 219 | J. Wang, C. Wu, C. Zhang, W. Ju, X. Wang, Z. Chen (China) and B. Fang (USA)   | 332 |
| Are remotely sensed traits suitable for ecological analysis? A case study of long-term drought effects on leaf mass per area of wetland vegetation   |     | Life Cycle Assessment of China's agroecosystems   |     |
| H. Feilhauer (Germany), T. Schmid (Spain), U. Faude (Germany), S. Sánchez-Carrillo and S. Cirujano (Spain)   | 232 | L. Liang (China, United States), R. Lal (United States), B.G. Ridoutt (Australia), Z. Du, D. Wang, L. Wang, W. Wu and G. Zhao (China)                     | 341 |
| The dynamics of sand-stabilization services in Inner Mongolia, China from 1981 to 2010 and its relationship with climate change and human activities   |     | The dynamics of sand-stabilization services in Inner Mongolia, China from 1981 to 2010 and its relationship with climate change and human activities      |     |
| D. Li, D. Xu, Z. Wang, X. You, X. Zhang and A. Song (China)  |     | D. Li, D. Xu, Z. Wang, X. You, X. Zhang and A. Song (China)   | 351 |

|   |     |     |
|---|-----|-----|
| Ecosystem health in coastal areas targeted by small-scale artisanal fisheries: Insights on monitoring and assessment<br>S. de Juan (Chile, Spain), M.D. Subida, S. Gelcich and M. Fernandez (Chile)   |     |     |
| Assessing the effects of grazing on variations of vegetation NPP in the Xilingol Grassland, China, using a grazing pressure index<br>D. Chi, H. Wang, X. Li, H. Liu and X. Li (China)   | 361 | 439 |
| Cuckoo density as a predictor of functional and phylogenetic species richness in the predictive modelling approach: Extension of Tryjanowski and Morelli (2015) paradigm in the analytical context<br>J.Z. Kosicki (Poland) and M. Hromada (Slovakia, Poland)                   | 372 | 454 |
| Cropland disturbance intensity: Plot-scale measurements, multilevel determinants and applications in rural environmental protection<br>M. Li, Y. Wang, B. Fu, P. Xu, E. Dai and C. Tian (China)   | 384 | 465 |
| Indicators of sustainability to assess aquaculture systems<br>W.C. Valenti, J.M. Kimpara, B.d.L. Preto and P. Moraes-Valenti (Brazil)   | 393 | 485 |
| Delimitation of ecological corridors in the Brazilian Atlantic Forest<br>J.S. Santos, C.C.C. Leite, J.C.C. Viana, A.R. dos Santos, M.M. Fernandes, V. de Souza Abreu, T.P. do Nascimento, L.S. dos Santos, M.R. de Moura Fernandes, G.F. da Silva and A.R. de Mendonça (Brazil) | 402 | 495 |
| Estimating soil organic carbon stocks using different modelling techniques in the semi-arid rangelands of eastern Australia<br>B. Wang, C. Waters, S. Orgill, A. Cowie, A. Clark, D. Li Liu, M. Simpson, I. McGowen and T. Sides (Australia)                                    | 414 | 503 |
| Modeling land use change using Cellular Automata and Artificial Neural Network: The case of Chunati Wildlife Sanctuary, Bangladesh<br>K. Islam, Md.F. Rahman (Japan) and M. Jashimuddin (Bangladesh)  | 425 | 512 |
| Cryptogamic communities as a useful bioindication tool for estimating the degree of soil pollution with heavy metals<br>K. Rola and P. Osyczka (Poland)   |     |     |
| Relationships between indicators of acid-base chemistry and fish assemblages in streams of the Great Smoky Mountains National Park<br>B.P. Baldigo, M.A. Kulp and J.S. Schwartz (United States)   |     |     |
| Monitoring changes of NDVI in protected areas of southern California<br>T.W. Gillespie, S. Ostermann-Kelm, C. Dong, K.S. Willis, G.S. Okin and G.M. MacDonald (USA)   |     |     |
| One year of transplant: Is it enough for lichens to reflect the new atmospheric conditions?<br>L. Paoli, A. Vannini (Italy), Z. Fačkovcová (Slovakia), M. Guarnieri (Italy), M. Bačkor (Slovakia) and S. Loppi (Italy)  |     |     |
| Replenishment of landscape water with reclaimed water: Optimization of supply scheme using transparency as an indicator<br>D. Ao, L. Luo, M. Dzakpasu, R. Chen, T. Xue and X.C. Wang (China)  |     |     |
| Dynamic selection of environmental variables to improve the prediction of aphid phenology: A machine learning approach<br>P. Holloway (UK, Ireland), D. Kudenko (UK, Russia) and J.R. Bell (UK)   |     |     |