

## Contents

**Fats and oils**

Essential oil composition and antioxidant activities of eight cultivars of Lavender and Lavandin from western Anatolia  
Ş. Kivrak (Turkey)

88

Composition, characteristics and antioxidant activities of fruit oils from *Idesia polycarpa* using homogenate-circulating ultrasound-assisted aqueous enzymatic extraction  
K. Hou, X. Yang, M. Bao, F. Chen, H. Tian and L. Yang (China)

205

Phenotypic and chemical variation of black mulberry (*Morus nigra*) genotypes  
A.-S. Hosseini, M. Akramian, A. Khadivi and H. Salehi-Arjmand (Iran)

260

Biologically active components in seeds of three *Nicotiana* species  
V. Popova, Z. Petkova, T. Ivanova, M. Stoyanova, L. Lazarov, A. Stoyanova, T. Hristeva, M. Docheva, V. Nikolova, N. Nikolov (Bulgaria) and V.D. Zheljazkov (USA)

375

Integrated soil-cotton system management enhances triacylglycerol yield and favourable fatty acid profile  
H. Yang, X. Zhang, B. Chen, Y. Meng, W. Zhao, Y. Wang and Z. Zhou (PR China)

393

**Non-food bioactive products**

Natural products-based pesticides: Design, synthesis and pesticidal activities of novel fraxinellone derivatives containing *N*-phenylpyrazole moiety  
R. Yang, T. Xu, J. Fan, Q. Zhang, M. Ding, M. Huang, L. Deng, Y. Lu and Y. Guo (PR China)

50

Polyphenolics in leaves of *Paris polyphylla*: An important high value Himalayan medicinal herb  
T. Mohd., T. Belwal, I.D. Bhatt, V. Pande and S.K. Nandi (India)

66

Modulation of the phenylpropanoid geranylation step in *Anethum graveolens* cultured calli by ferulic acid and umbelliferone  
S. Genovese, F. Epifano, S. Fiorito, V.A. Taddeo, F. Preziuso and D. Fraternali (Italy)

128

Enzyme-assisted ultrasonic-microwave synergistic extraction and UPLC-QTOF-MS analysis of flavonoids from Chinese water chestnut peels  
L. Xu, W. He (PR China), M. Lu, B. Yuan (United States), M. Zeng, G. Tao, F. Qin, J. Chen, Y. Guan and Z. He (PR China)

179

- Spatio-temporal dynamics of mealybug (Hemiptera: Pseudococcidae) populations in plantations of *Jatropha curcas* L. in Yucatan, Mexico  
C.C. Góngora-Canul, G. Martínez-Sebastián, E.A. Aguilera-Cauich, A. Uc-Varguez, G. López-Puc (Mexico) and O. Pérez-Hernández (United States) 110
- Agricultural feedstocks of two *Brassica* oilseed crops and energy cogeneration with pure vegetable oil for a sustainable short agro-energy chain in Sicily (Italy)  
M. Licata, S. La Bella, L. Lazzeri, R. Matteo, C. Leto, F. Massaro and T. Tuttolomondo (Italy) 140
- Assessment system to characterise and compare different hemp varieties based on a developed lab-scaled decortication system  
S. Wang, H.-J. Gusovius, C. Lühr (Germany), S. Musio (Italy), B. Uhrlaub (Germany), S. Amaducci (Italy) and J. Müssig (Germany) 159
- Establishment of new strategies to quantify and increase the variability in the Brazilian *Jatropha* genotypes  
B.G. Laviola, A.A. Alves, T.B. Rosado, L.L. Bhering, E.F. Formighieri and L. de Azevedo Peixoto (Brazil) 216
- Cuticular wax variants in a population of switchgrass (*Panicum virgatum* L.)  
J.M. Weaver, G. Lohrey, P. Tomasi, J.M. Dyer, M.A. Jenks and K.A. Feldmann (USA) 310
- Sequential extraction and reactive extraction processing of spent coffee grounds: An alternative approach for pretreatment of biodiesel feedstocks and biodiesel production  
N. Tuntiwattanapun and C. Tongcumpou (Thailand) 359
- Bio-based Materials from Crops**
- Pollen mediated gene flow in a small experimental population of *Moringa oleifera* Lam. (Moringaceae)  
J.-C. Wu, Y.-P. Zhang, Y.-X. Zheng and X.-M. Peng (China) 28
- Preparation and characterization of biobased microspheres from lignin sources  
J. Sameni, S. Krigstin (Canada), S.A. Jaffer (Canada, USA) and M. Sain (Canada) 58
- Subcritical water extraction of bioactive components from ginseng roots (*Panax ginseng* C.A. Mey)  
Y. Zhang, Y. Zhang, A.A. Taha, Y. Ying, X. Li, X. Chen and C. Ma (PR China) 118
- Energy use efficiency and economic feasibility of Jerusalem artichoke production on arid and coastal saline lands  
Y.R. Fang, J.A. Liu (PR China), Y. Steinberger (Israel) and G.H. Xie (PR China) 131
- A simple methodology for improving the performance and sustainability of rigid polyurethane foam by incorporating industrial lignin  
A.N. Hayati, D.A.C. Evans, B. Laycock, D.J. Martin and P.K. Annamalai (Australia) 149
- Castor oil-based cationic waterborne polyurethane dispersions: Storage stability, thermo-physical properties and antibacterial properties  
H. Liang, L. Liu, J. Lu, M. Chen and C. Zhang (China) 169
- Chemical composition and resistance of Italian stone pine (*Pinus pinea* L.) wood against fungal decay and wetting  
M. De Angelis, M. Romagnoli (Italy), V. Vek, I. Poljanšek, P. Oven, N. Thaler, B. Lesar, D. Kržišnik and M. Humar (Slovenia) 187
- Modeling the seed yield of Ajowan (*Trachyspermum ammi* L.) using artificial neural network and multiple linear regression models  
M. Niazi, S.A. Sadat-Noori and M. Abdipour (Iran) 224
- Soy protein isolate nanocomposites reinforced with nanocellulose isolated from licorice residue: Water sensitivity and mechanical strength  
Y. Han, M. Yu and L. Wang (PR China) 252
- Preparation of hydrothermal carbon acid catalyst from defatted rice bran  
P. Wataniyakul, P. Boonnoun (Thailand), A.T. Quitain, T. Kida (Japan), N. Laosiripojana and A. Shotipruk (Thailand) 286
- Use of cardanol-based acrylate as reactive diluent in UV-curable castor oil-based polyurethane acrylate resins  
Y. Hu, Q. Shang, J. Tang, C. Wang, F. Zhang, P. Jia, G. Feng, Q. Wu, C. Liu, L. Hu, W. Lei and Y. Zhou (PR China) 295
- Effects of cationization of high amylose maize starch on the performance of starch/montmorillonite nanobiocomposites  
W. Wang, P. Song, R. Wang, R. Zhang, Q. Guo, H. Hou and H. Dong (PR China) 333

- Hairy root cultures of *Salvia viridis* L. for production of polyphenolic compounds  
*I. Grzegorzczak-Karolak, Ł. Kuźma, E. Skala and A.K. Kiss (Poland)* 235
- Lippia gracilis* Schauer essential oil nanoformulation prototype for the control of *Thielaviopsis paradoxa*  
*E.M.O. Cruz, M.C. Mendonça, A.F. Blank, T.S. Sampaio, J.A.O. Pinto, P.R. Gagliardi, L.F.G. Oliveira Junior, R.S.N. de Lima, R.S. Nunes and D.R.N. Warwick (Brazil)* 245
- Insecticidal activity of *Bacillus thuringiensis* crystals and thymol mixtures  
*E. Konecka, E. Czarniewska, A. Kaznowski and J. Grochowska (Poland)* 272
- Bioassay-guided purification of  $\alpha$ -amylase,  $\alpha$ -glucosidase inhibitors and DPPH radical scavengers from roots of *Rheum turkestanicum*  
*H. Dehghan, P. Salehi and M.S. Amiri (Iran)* 303
- Melissa officinalis* L. subsp. *altissima* (Sibth. & Sm.) Arcang. essential oil: Chemical composition and preliminary antimicrobial investigation of samples obtained at different harvesting periods and by fractionated extractions  
*M. Božović (Montenegro, Italy), S. Garzoli, A. Baldisserotto, C. Romagnoli, F. Pepi, S. Cesa, S. Vertuani, S. Manfredini and R. Ragno (Italy)* 317
- Trichome-specific and developmentally regulated biosynthesis of nepetalactones in leaves of cultivated *Nepeta rtanjensis* plants  
*N. Aničić, D. Matekalo, M. Skorić, I. Pećinar, M. Brkušanin, J. Nestorović Živković, S. Dmitrović, Z. Dajić Stevanović (Serbia), H. Schulz (Germany) and D. Mišić (Serbia)* 347
- Essential oil composition and total phenolic, flavonoid contents, and antioxidant activity of sage (*Salvia officinalis* L.) extract under chitosan application and irrigation frequencies  
*N. Vosoughi, M. Gomarian (Iran), A. Ghasemi Pirbalouti (Iran, USA), S. Khaghani and F. Malekpoor (Iran)* 366
- Beyond mosquitoes—Essential oil toxicity and repellency against bloodsucking insects  
*G. Benelli (Italy) and R. Pavela (Czech Republic)* 382
- Fibres, proteins and carbohydrates**
- Role of major constituents of coconut fibres on absorption of ionic dyes  
*A.K. Samanta, G. Basu and L. Mishra (India)* 20
- Development of distinct cell wall layers both in primary and secondary phloem fibers of hemp (*Cannabis sativa* L.)  
*T.E. Chernova, P.V. Mikshina, V.V. Salnikov, N.N. Ibragimova, O.V. Sautkina and T.A. Gorshkova (Russian Federation)* 97
- Effect of spray drying on the properties of camelina gum isolated from camelina seeds  
*X. Cao, N. Li, G. Qi, X.S. Sun and D. Wang (United States)* 278
- Common oak (*Quercus robur*) acorn as a source of natural coagulants for water turbidity removal  
*M.G. Antov, M.B. Šćiban, J.M. Prodanović, D.V. Kukić, V.M. Vasić, T.R. Đorđević and M.M. Milošević (Serbia)* 340
- Biorefinery, valorization of byproducts**
- Profiling and seasonal variation of chemical constituents from *Pseudotsuga menziesii* wood  
*H. Mbakidi-Ngouaby, E. Pinault, V. Gloaguen, G. Costa, V. Sol, M. Millot and L. Mambu (France)* 34
- Efficient conversion of wheat straw into methyl levulinate catalyzed by cheap metal sulfate in a biorefinery concept  
*C. Chang, L. Deng and G. Xu (China)* 197
- Magnetic and reusable  $MgO/MgFe_2O_4$  nanocatalyst for biodiesel production from sunflower oil: Influence of fuel ratio in combustion synthesis on catalytic properties and performance  
*S. Alaei, M. Haghghi, J. Toghiani and B. Rahmani Vahid (Iran)* 322
- Bioenergy - crops and processing**
- Explant origin and culture media factors drive the somatic embryogenesis response in *Acrocomia aculeata* (Jacq.) Lodd. ex Mart., an emerging oil crop in the tropics  
*M.M.C. Granja, S.Y. Motoike, A.P.S. Andrade, T.R. Correa, E.A.T. Picoli and K.N. Kuki (Brazil)* 1
- Arbuscular mycorrhizal fungi improve plant growth of *Ricinus communis* by altering photosynthetic properties and increasing pigments under drought and salt stress  
*T. Zhang, Y. Hu, K. Zhang, C. Tian and J. Guo (China)* 13
- Integrated bioethanol production from triticale grain and lignocellulosic straw in Western Canada  
*E. Mupondwa, X. Li and L. Tabil (Canada)* 75