

<b>Nutrient uptake-modification of different plant species in Mediterranean climate by arbuscular mycorrhizal fungi</b>	
<b>M. Rafique İ. Ortas</b>	› 65
<b>Time period and nutrient contents alter the mycorrhizal responsiveness of citrus seedlings</b>	
<b>İ. Ortas A. Demirbas C. Akpinar</b>	› 72
<b>Under sterilized and non-sterilized soil conditions, mycorrhizal dependency in citrus plants depends on phosphorus fertilization rather than zinc application</b>	
<b>İ. Ortas A. Demirbas C. Akpinar</b>	› 81
<b>The foliar water relations in orange plants (<i>Citrus sinensis</i> L.) are affected by sooty mold</b>	
<b>P. Insausti E.L. Ploschuk</b>	› 88
<b>Polyphenolic profiling and chemometric analysis of leaves from Italian <i>Ficus carica</i> L. varieties. Polyphenol compounds in common fig</b>	
<b>R. Petruccelli F. Ieri L. Ciaccheri A. Bonetti</b>	› 94
<b>Fresh figs (<i>Ficus carica</i> L.): pomological characteristics, nutritional value, and phytochemical properties</b>	
<b>S. Mahmoudi M. Khali A. Benkhaled I. Boucetta Y. Dahmani Z. Attallah S. Belbraouet</b>	› 104
<b>The influence of foliar application of urea, zinc chelate and boric acid on growth characteristics, photosynthesis and survival of date palm offshoots (<i>Phoenix dactylifera</i> cv. Mazafati)</b>	
<b>H. Mohammadi M. Khezri</b>	› 114