

| | | | |
|--|----|--|----|
| Is <i>Halyomorpha halys</i> (Hemiptera: Pentatomidae) a threat to California rice? <i>M.-A. Aghaee, S. Lee Rice, J.M. Milnes, K.M. Goding and L.D. Godfrey</i> | 1 | A leaf-wall-to-spray-device distance and leaf-wall-density-based automatic route-planning spray algorithm for vineyards <i>G. Gao, K. Xiao and Y. Ma</i> | 33 |
| Management of beet rust in accordance with IPM principles <i>R. Kristoffersen, A.L. Hansen, L. Munk, N. Cedergreen and L.N. Jørgensen</i> | 6 | Safety and efficacy of quinoa (<i>Chenopodium quinoa</i>) saponins derived molluscicide to control of <i>Pomacea maculata</i> in rice fields in the Ebro Delta, Spain <i>M. Castillo-Ruiz, H. Cañon-Jones, T. Schlotterbeck, M.A. Lopez, Á. Tomas and R. San Martín</i> | 42 |
| Resistance and detoxification enzyme activities to bifenthrin in <i>Oxycarenus hyalinipennis</i> (Hemiptera: Lygaeidae) <i>M. Bilal, S. Freed, M.Z. Ashraf and A. Rehan</i> | 17 | Insecticidal activity of natural products against vineyard mealybugs (Hemiptera: Pseudococcidae) <i>F. Tacoli, V.A. Bell, E. Cargus and F. Pavan</i> | 50 |
| Comparative assessment of the relative proportion of weed morphology, diversity, and growth under new generation tillage and crop establishment techniques in rice-based cropping systems <i>R. Nandan, V. Singh, S.S. Singh, V. Kumar, K.K. Hazra, C.P. Nath, S.P. Poonia and R.K. Malik</i> | 23 | Multi-study analysis of the added benefits of combining soil solarization with fumigants or non-chemical measures <i>E. Shlevin, A. Gamliel, J. Katan and D. Shtienberg</i> | 58 |
| | | Newsletter | 66 |