

CONTENTS

Articles

GODJO, A., AFOUDA, L., BAIMEY, H., COUVREUR, M., ZADJI, L., HOUSSOU, G., BERT, W., WILLEMS, A. & DECRAEMER, W.— <i>Steinernema kandii</i> n. sp. (Rhabditida: Steinernematidae), a new entomopathogenic nematode from northern Benin	107-128
MONTEIRO, J.M.S., MATTOS, V.S., SANTOS, M.F.A., GOMES, A.C.M.M., CORREA, V.R., SOUSA, D.A., CARES, J.E., PINHEIRO, J.B. & CARNEIRO, R.M.D.G.— Additional information on <i>Meloidogyne polycephannulata</i> and its proposal as a junior synonym of <i>M. incognita</i>	129-146
MIRZAI FOUADVAND, Z., POURJAM, E., KANZAKI, N., GIBLIN-DAVIS, R.M. & PEDRAM, M.— Description of <i>Basilaphelenchus brevicaudatus</i> n. sp. (Aphelenchoidea: Tylaphelenchinae) from a dead forest tree in northern Iran	147-158
GOLHASAN, B., FANG, Y., LI, H., TANHA MAAFI, Z. & HEYDARI, R.— Description of <i>Aphelenchoides tabarestanensis</i> n. sp. (Nematoda: Aphelenchoididae) isolated from <i>Pinus brutia</i> in Northern Iran	159-169
SEIML-BUCHINGER, V.V., ZINOVIEVA, S.V., UDALOVA, Z.V. & MATVEEVA, E.M.— Jasmonic acid modulates <i>Meloidogyne incognita</i> – tomato plant interactions	171-180
MARIA, M., CAI, R., CASTILLO, P., CANTALAPIEDRA-NAVARRETE, C. & ZHENG, J.— Characterisation of two Chinese native <i>Hemicriconemoides</i> species (Nematoda: Criconematidae) with updated descriptions of <i>H. chitwoodi</i> Esser, 1960 and <i>Criconemoides myungsugae</i> Choi & Geraert, 1975	181-205
KANZAKI, N., EKINO, T. & MASUYA, H.— <i>Seinura caverna</i> n. sp. (Tylenchomorpha: Aphelenchoididae), an androdioecious species isolated from bat guano in a calcareous cave	207-225