

CONTENTS

Articles

SERGEeva, N.G., SHADRIN, N.V. & ANUFRIIEVA, E.V.— Long-term changes (1979-2015) in the nematode fauna in Sivash Bay (Sea of Azov), Russia, worldwide the largest hypersaline lagoon, during salinity transformations	337-347
IMRAN, Z., ABOLAFIA, J., PEÑA-SANTIAGO, R. & AHMAD, W.— Morphological and molecular characterisation of <i>Moshajia idiofora</i> Siddiqi, 1982 (Dorylaimida: Qudsianematidae), with new insights into the phylogeny of the enigmatic Lordellonematinae.....	349-360
KANZAKI, N., MASUYA, H., ICHIHARA, Y., MAEHARA, N., AIKAWA, T., EKINO, T. & IDE, T.— <i>Bursaphelenchus carpini</i> n. sp., <i>B. laciniatae</i> n. sp. and <i>B. cryphali okhotskensis</i> n. subsp. (Nematoda: Aphelenchoididae) isolated from Japan	361-388
KITAGAMI, Y., KANZAKI, N., TANIKAWA, T. & MATSUDA, Y.— Free-living nematodes associated with pine cones of <i>Pinus thunbergii</i> and <i>P. taeda</i> at Japanese coastal and inland forest sites	389-400
LAWILAO, J.G., SWAMY, B.P.M., KUMAR, A., CABASAN, M.T.N. & DE WAELE, D.— Mapping quantitative trait loci of <i>Meloidogyne graminicola</i> resistance and tolerance in a recombinant inbred line population of <i>Oryza glaberrima</i> × <i>O. sativa</i>	401-417
JAHANSOHI AFSHAR, F., POURJAM, E. & PEDRAM, M.— New morphological observations on <i>Neolobocriconema serratum</i> (Khan & Siddiqi, 1963) Mehta & Raski, 1971 (Rhabditida: Criconematidae).....	419-434
ESMAEILI, M., HEYDARI, R., GOLHASAN, B. & YE, W.— Molecular phylogenetic study of <i>Bealius</i> Massey & Hinds, 1970 (Nematoda: Hexatylina: Sphaerulariidae), with description of <i>B. pinus</i> n. sp. from Iran	435-444