

# CC GREEN Scanauftrag



**Auftragsnummer:** ZBMED-BN201020602-1  
**Auftragsdatum:** 20.10.20 - 06:00

---

**Signatur:** Z 6884

**Title:** Journal of soils and crops

**ISSN:** 0971-2836  
**Erscheinungsjahr:** 2019  
**Erscheinungsort:** Nagpur  
**Band Heft:** 29(2)

**Bitte Inhaltsverzeichnis scannen!**

## C O N T E N T S

<b>Maximizing land and water productivity of sudan grass under sprinkler irrigation in sandy soil</b>	<b>207-217</b>
Ahmed M. Taha <sup>1</sup> , Azza Kh. Salem <sup>2</sup> and Nabil E. G. Mekhaile <sup>3</sup>	
<b>Intercropping systems for sugar beet to improve its land and water productivity</b>	<b>218-226</b>
A. A. Zohry <sup>1</sup> and S. A. Ouda <sup>2</sup>	
<b><i>In Vitro</i> regeneration of ginger using tip explants</b>	<b>227-231</b>
Graceli Yephthomi <sup>1</sup> and C.S Maiti <sup>2</sup>	
<b>Remote sensing based greenness modeling of different crops in Lalgudi block using landsat 8 image</b>	<b>232-235</b>
J. Ramachandran <sup>1</sup> , R. Lalitha <sup>2</sup> and S. Vallal Kannan <sup>3</sup>	
<b>Study on growth yield and economics in maize as influenced by different levels of nitrogen applied through vermicompost and spacing in hill areas of North East India</b>	<b>236-242</b>
N. Khumdemo Ezung <sup>1</sup> and Tiatula Jamir <sup>2</sup>	
<b>Biocontrol and growth promoting potential of eight PGPFs jute and sunnhemp</b>	<b>243-250</b>
Anuradha Bandopadhyay <sup>1</sup> , Saurindra Kishore Bhattacharya <sup>2</sup> and Nirmalendu Das <sup>3</sup>	
<b>Effect of enhanced N, P, K and Zn fertilizer levels on yield attributes, yield and economics of wet seeded rice under Tamirabarani command area</b>	<b>251-255</b>
M. Madhan Kumar <sup>1</sup> , M. Hemalatha <sup>2</sup> , M. Joseph <sup>3</sup> , R. Nandhini <sup>4</sup> and M. Arun Raj <sup>5</sup>	
<b>Respodip treatment with nano emulsion of hexanal to reduce the anthracnose disease of banana and extend its shelf life</b>	<b>256-261</b>
I. Arumuka Pravin <sup>1</sup> , S. Srivignesh <sup>2</sup> , D. Durgadevi <sup>3</sup> , K. S. Subramanian <sup>4</sup> and A. S. Krishnamoorthy <sup>5</sup>	
<b>Efficacy of putrescine and IBA on biochemical and yield contributing parameters of black gram</b>	<b>262-267</b>
R. D. Deotale <sup>1</sup> , N. D. Jadhav <sup>2</sup> , Shanti Patil <sup>3</sup> , Sapana Baviskar <sup>4</sup> , Vandana S. Madke <sup>5</sup> and Vandana Kalamkar <sup>6</sup>	
<b>Effect of gamma rays on quantitative traits in M<sub>2</sub> generation of Lathyrus (<i>Lathyrus sativus</i> L.)</b>	<b>268-273</b>
V. T. Chavan <sup>1</sup> , Shanti R. Patil <sup>2</sup> , Shubhangi K. Maraskole <sup>3</sup> , Vandana S. Madke <sup>4</sup> , Sapana B. Baviskar <sup>5</sup> and Vandana B. Kalamkar <sup>6</sup>	
<b>Effect of cow urine and NAA on morpho-physiological parameters and yield of wheat</b>	<b>274-279</b>
S. B. Korade <sup>1</sup> , R. D. Deotale <sup>2</sup> , N. D. Jadhav <sup>3</sup> , V. A. Guddhe <sup>4</sup> and O. G. Thakre <sup>5</sup>	
<b>Effect of imazethapyr + imazamox on weed control in soybean</b>	<b>280-284</b>
S.M.Deshkari <sup>1</sup> , P.C.Pagar <sup>2</sup> , S.T.Dangore <sup>3</sup> , V.S.Khawale <sup>4</sup> and H.S.Mendhe <sup>5</sup>	
<b>Changes in chemical and biochemical parameters of chickpea (<i>Cicer arietinum</i> L.) sprayed with putrescine and naphthalene acetic acid</b>	<b>285-292</b>
R. D. Deotale <sup>1</sup> , Vikas R. Jaybhaye <sup>2</sup> , Shanti R. Patil <sup>3</sup> , Vandana Kalamkar <sup>4</sup> and S.R. Kamdi <sup>5</sup>	

<b>Efficacy of different herbicides on weed control in soybean</b>	<b>293-296</b>
G. Vanisree <sup>1</sup> , V. S. Khawale <sup>2</sup> , N. D. Parlwar <sup>3</sup> and D. J. Jiotode <sup>4</sup>	
<b>Experimental study on biochemical parameters and yield of M<sub>4</sub> Indian mustard mutants</b>	<b>297-301</b>
R. D. Deotale <sup>1</sup> , Ashish P. Dhongade <sup>2</sup> , S. R. Kamdi <sup>3</sup> , Vandana S. Madke <sup>4</sup> , M. P. Meshram <sup>5</sup> and Vandana B. Kalamkar <sup>6</sup>	
<b>Comparative study of growth, yield attributes and yield of paddy varieties as influenced by seed rates under drilled condition</b>	<b>302-305</b>
A. Chaitanya <sup>1</sup> , N. D. Parlwar <sup>2</sup> , V. S. Khawale <sup>3</sup> , P. C. Pagar <sup>4</sup> and D. J. Jiotode <sup>5</sup>	
<b>Impact of foliar sprays of chitosan and IBA on chemical, biochemical and yield contributing parameters of pigeonpea</b>	<b>306-311</b>
Rajesh D. Deotale <sup>1</sup> , O. G. Thakare <sup>2</sup> , P. V. Shende <sup>3</sup> , Shanti R. Patil <sup>4</sup> , S. R. Kamdi <sup>5</sup> , M. P. Meshram <sup>6</sup> and Vandana S. Madke <sup>7</sup>	
<b>Morpho- physiological traits and yield in chickpea as influenced by foliar application of ascorbic acid and zinc sulphate</b>	<b>312-318</b>
Dnyaneshwar A. Raut <sup>1</sup> , Rajesh D. Deotale <sup>2</sup> , A. Blesseena <sup>3</sup> , Vishal S. Hivare <sup>4</sup> , Satish E. Pise <sup>5</sup> and Sanket A. Yelore <sup>6</sup>	
<b>Genetic divergence studies in maize (<i>Zea mays</i> L.) accessions</b>	<b>319-328</b>
S. H. Palkar <sup>1</sup> , S. U. Charjan <sup>2</sup> , S. R. Patil <sup>3</sup> , V. T. Chavan <sup>4</sup> and P. B. Chavan <sup>5</sup>	
<b>Response of humic acid through vermicompost wash and NAA on chemical, biochemical, yield and yield contributing parameters of sesamum</b>	<b>329-335</b>
R. D. Deotale <sup>1</sup> , V. A. Guddhe <sup>2</sup> , S. R. Kamdi <sup>3</sup> , Shanti R. Patil <sup>4</sup> , Vandana S. Madke <sup>5</sup> , Sapana B. Baviskar <sup>6</sup> and M. P. Meshram <sup>7</sup>	
<b>Response foliar application of tocopherol and micronutrients on morpho-physiological parameters and yield of chickpea</b>	<b>336-342</b>
A. Blesseena <sup>1</sup> , R. D. Deotale <sup>2</sup> , D. A. Raut <sup>3</sup> , S. E. Pise <sup>4</sup> , S. A. Yellore <sup>5</sup> and V. S. Hivare <sup>6</sup>	
<b>Genetic studies F<sub>2</sub> population in soybean</b>	<b>343-347</b>
Vasant S. Pawar <sup>1</sup> , Sandeep R. Kamdi <sup>2</sup> , Milind P. Meshram <sup>3</sup> , Rajesh D. Deotale <sup>4</sup> , Shanti R. Patil <sup>5</sup> , Ritik D. Bisane <sup>6</sup> , Pradnya P. Bambodkar <sup>7</sup>	
<b>Morpho-physiological traits and yield in safflower as influenced by foliar application of humic acid and NAA</b>	<b>348-353</b>
Vishal S. Hivare <sup>1</sup> , Rajesh D. Deotale <sup>2</sup> , Ashish P. Dhongade <sup>3</sup> , Satish E. Pise <sup>4</sup> , Dnyaneshwar A. Raut <sup>5</sup> and A. Blesseena <sup>6</sup>	
<b>Genetic studies in M<sub>4</sub> population of soybean cultivar TAMS - 38</b>	<b>354-359</b>
Ritik D. Bisane <sup>1</sup> , Sandeep R. Kamdi <sup>2</sup> , Rajesh D. Deotale <sup>3</sup> , Milind P. Meshram <sup>4</sup> , Shanti R. Patil <sup>5</sup> , Vasant S. Pawar <sup>6</sup> and Pradnya P. Bambodkar <sup>7</sup>	
<b>Influence of zinc and iron on morpho-physiological parameters and yield of Lathyrus (<i>Lathyrus sativus</i> L.)</b>	<b>360-365</b>
Satish E. Pise <sup>1</sup> , P. V. Shende <sup>2</sup> , R. D. Deotale <sup>3</sup> , Dnyaneshwar A. Raut <sup>4</sup> , A. Blesseena <sup>5</sup> and Vishal S. Hivare <sup>6</sup>	

<b>Genetic variability studies in F<sub>3</sub> segregating generations in soybean for yield and its components</b>	<b>366-370</b>
Pradnya P. Bambodkar <sup>1</sup> , S. R. Kamdi <sup>2</sup> , Shanti R. Patil <sup>3</sup> , M. P. Meshram <sup>4</sup> , R. D. Deotale <sup>5</sup> , R. D. Bisane <sup>6</sup> and V. S. Pawar <sup>7</sup>	
<b>Effect of partial feeding of hydroponic maize on performance of crossbred heifers</b>	<b>371-375</b>
Chhaya Thombre <sup>1</sup> , V. G. Atkare <sup>2</sup> and N. S. Chore <sup>3</sup>	
<b>Effect of nipping and growth retardant on growth yield and uptake of nutrient of pigeonpea</b>	<b>376-380</b>
R. R. Kolhe <sup>1</sup> , N. D. Parlawar <sup>2</sup> , D. J. Jiotode <sup>3</sup> , V. S. Khawale <sup>4</sup> , T. A. Chavhan <sup>5</sup> and R. I. Samrutwar <sup>6</sup>	
<b>Combining ability analysis for yield and its component traits in maize (<i>Zea mays</i> L.)</b>	<b>381-385</b>
P. B. Chavan <sup>1</sup> , M. K. Moon <sup>2</sup> , S. R. Patil <sup>3</sup> , Prakash Hosur <sup>4</sup> and Ankush Sapkal <sup>5</sup>	
<b>Gamma rays induced genetic variability in M<sub>2</sub> generation of soybean (<i>Glycine max</i> (L.) Merrill)</b>	<b>386-391</b>
S. R. Kamdi <sup>1</sup> , G. A. Kankal <sup>2</sup> , M. P. Meshram <sup>3</sup> , R. D. Deotale <sup>4</sup> , S. K. Gupta <sup>5</sup> , S. S. Bhure <sup>6</sup> and J. M. Parbat <sup>7</sup>	
<b>Short communication</b>	
<b>Effect of soil test crop response (STCR) fertilizer prescriptions combined with integrated plant nutrient supply (IPNS) on performance of normal and late sown toria</b>	<b>391-395</b>
Bikram Borkotoki <sup>1</sup> , K.N. Das <sup>2</sup> , Anjali Basumatary <sup>3</sup> and Prabal Saikia <sup>4</sup>	