

CONTENTS

ORIGINAL ARTICLES

- 95 Combining ability of yield and yield components among *Fusarium oxysporum* f.sp. *strigae*-compatible and *Striga*-resistant sorghum genotypes
Emmanuel Mrema, Hussein Shimelis and Mark Laing
- 109 Yield structure components of autumn- and spring-sown pea (*Pisum sativum* L.)
Reinhard W. Neugschwandner, Alexander Bernhuber, Stefan Kammlander, Helmut Wagentristl, Agnieszka Klimek-Kopyra and Hans-Peter Kaul
- 117 Spatial distribution of soil organic matter in a coal mining subsidence area
Huijuan Zhang, Wenkai Liu, Hebing Zhang, Liangxin Fan and Shouchen Ma
- 128 The long-term effect of legumes as forecrops on the productivity of rotation winter triticale-winter rape with nitrogen fertilisation
Grażyna Szymańska, Agnieszka Faligowska, Katarzyna Panasiewicz, Jerzy Szukała, Karolina Ratajczak and Hanna Sulewska
- 135 Host-status of 20 sweet potato lines to *Meloidogyne* species in South Africa
K. M. Pofu, P. W. Mashela and S. Laurie
- 140 Post-harvest residues of field pea/oat mixtures as a source of biological nitrogen for hybrid winter rye
Anna Płaza, Barbara Gąsiorowska and Emilia Rzążewska
- 150 Cassava-legume intercrop: I. Effects of relative planting dates of legumes on cassava productivity
Khutso Debra Legodi and J. B. O. Ogola
- 158 Cassava-legume intercrop: II. Effects of relative planting dates of legumes on the productivity of legumes
Khutso Debra Legodi and J. B. O. Ogola
- 165 Early interspecific dynamics, dry matter production and nitrogen use in Kernza (*Thinopyrum intermedium*) – alfalfa (*Medicago sativa* L.) mixed intercropping
Shoujiao Li, Ana Barreiro, Erik Steen Jensen, Yingjun Zhang and Linda-Maria Dimitrova Mårtensson