

CONTENTS

Forum article

- ZULLINI, A. & SEMPRUCCI, F.— Morphological differences between free-living soil and freshwater nematodes in relation to their environments 125-132

Articles

- KRAINER, L., SUSULOVSKY, A., RYBARCZYK-MYDŁOWSKA, K., KUBICZ, M. & PEÑA-SANTIAGO, R.— Morphological and molecular characterisation of one new and one known species of *Makatinus* Heyns, 1965 (Dorylaimida: Aporcelaimidae) from Ukraine, with new insights into its phylogeny 133-146
- MISTANOĞLU, İ., ÖZALP, T. & DEVRAN, Z.— The efficacy of molecular markers associated with virulence in root-knot nematodes 147-154
- ABOLAFIA, J., RUIZ-CUENCA, A.N., SHOKOOHI, E., DU PREEZ, G. & FOURIE, H.— Redescription of *Paracrobeles laterellus* Heyns, 1968 (Rhabditida: Cephalobidae) from Botswana 155-164
- NUAIMA, R.H., ROEB, J., HALLMANN, J., DAUB, M. & HEUER, H.— Significant genetic differences among *Heterodera schachtii* populations within and among sugar beet production areas 165-177
- IVANOVA, E.S., GERASKINA, A.P. & SPIRIDONOV, S.E.— Two new species of *Phasmarhabditis* Andrásy, 1976 (Nematoda: Rhabditidae) associated with land snails in Northwest Caucasus, Russian Federation: description and molecular affiliation 179-197
- FANG, Y., MARIA, M., SÁNCHEZ-MONGE, A. & GU, J.— Phylogeny and redescription of *Laimaphelenchus pannocaudus* Massey, 1966 (Rhabditida: Aphelenchoididae) intercepted from Canadian *Picea glauca* 199-212
- DE O. SILVA, J., SANTANA, M.V., CARNEIRO, F.A. & DA ROCHA, M.R.— Reaction of tomato genotypes to *Meloidogyne enterolobii* and effects of resistance inducing products 213-220
- NZOGELA, Y.B., LANDSCHOOT, S., KIHUPI, A.-L., COYNE, D.L. & GHEYSEN, G.— Pathogenicity of the root-lesion nematode, *Pratylenchus zaeae*, on rice genotypes under different hydro-ecologies in Tanzania 221-233

Short communications

- POINAR, G.O., MYER, A. & FORSCHLER, B.T.— Mermithid nematode parasites of eastern subterranean termites in North America 235-237
- ENGELBRECHT, G., VAN RENSBURG, P.J.J., FOURIE, H. & CLAASSENS, S.— *In vitro* bioassays to determine the effect of *Bacillus soli* filtrates on the paralysis of *Meloidogyne incognita* second-stage juveniles 239-243