

CONTENTS

ORIGINAL ARTICLES

---

- 1 Finding best field pea (*Pisum sativum* L.) cultivars for breeding in Northern climatic conditions  
*Margit Olle, Ingrid Helvi Williams, Eduardo Rosa and Sirje Tamm*
- 8 Influence of alcoholic extracts from *Helianthus annuus* L. roots on the photosynthetic activity of *Sinapis alba* L. cv. Barka plants  
*Joanna Puła, Peiman Zandi, Alina Stachurska-Swakoń, Beata Barabasz-Krasny, Katarzyna Możdżeń and Yaosheng Wang*
- 14 Soil physical and chemical quality as influenced by soil tillage managements under double cropping rice system of southern China  
*Haiming Tang, Chao Li, Xiaoping Xiao, Wenguang Tang, Kaikai Cheng, Xiaochen Pan, Ke Wang and Weiyan Li*
- 24 Density-dependent growth response and sensitivity of *Meloidogyne incognita* to *Maerua angolensis* and *Tabernaemontana elegans*: Card model  
*M. C. Khosa, Z. P. Dube, M. Tselanyane, G. Fouche, D. De Waele and M. S. Daneel*
- 31 The effect of acidified slurry on crop uptake and leaching of nutrients from a loamy topsoil  
*V. Loide, T. Saue, T. Võsa and K. Tamm*
- 39 The phytochemical quality of *Camelina sativa* seed and oil  
*Danuta Kurasiak-Popowska and Kinga Stuper-Szablewska*
- 48 Effect of liming products on soil detachment resistance, measured with a cohesive strength meter  
*Daniel Aviles, Kerstin Berglund, Ingrid Wesström and Abraham Joel*
- 56 Varied macronutrient uptake by plants as an effect of different fertilisation schemes evaluated by PCA  
*Monika Jakubus and Ewa Bakinowska*
- 69 The effect of salt-tolerant antagonistic bacteria from tomato rhizosphere on plant growth promotion and damping-off disease suppression under salt-stress conditions  
*Thamodini Gaya Karunasinghe, Issa Hashil Al-Mahmooli, Abdullah Mohammed Al-Sadi and Rethinasamy Velazhahan*
- 76 Assessment of the genetic diversity and population structure of rice genotypes using SSR markers  
*William Titus Suvi, Hussein Shimelis, Mark Laing, Isack Mathew and Admire Isaac Tichafa Shayanowako*
- 87 The soil microbial hydrolytic activity, content of nitrogen and organic carbon were enhanced by organic farming management using cover crops and composts in potato cultivation  
*Viacheslav Eremeev, Liina Talgre, Jaan Kuht, Erkki Mäeorg, Keyvan Esmaeilzadeh-Salestani, Maarika Alaru, Evelin Loit, Eve Runno-Paurson and Anne Luik*