

NEMATODOLOGY 22 (6), 2020
Also available online: brill.com/nemy

CONTENTS

Articles

| | |
|---|---------|
| GAO, D., WANG, F., LI, J., YU, S., LI, Z. & ZHAO, J.— Soil nematode communities as indicators of soil health in different land use types in tropical area | 595-610 |
| MOHAMMADI ZAMELEH, F., KAREGAR, A., GHADERI, R. & MOKARAM HESAR, A.— Morphological and molecular characterisation of <i>Helicotylenchus ciceri</i> n. sp. and <i>H. scoticus</i> Boag & Jairajpuri, 1985 (Nematoda: Hoplolaimidae) from Iran | 611-626 |
| DAVIES, K.A., BARTHOLOMAEUS, F., MEI LI, D., ZHAO, Z.Q., YE, W. & GIBLIN-DAVIS, R.M.— <i>Ficophagus</i> (Nematoda: Aphelenchoididae) from sycones of <i>Ficus</i> subgenus <i>Urostigma</i> , sections <i>Malvanthera</i> and <i>Urostigma</i> , in eastern Australia | 627-653 |
| POINAR JR, G.O. & CURRIE, D.C.— Mermithid nematode (Nematoda: Mermithidae) parasites of a fossil black fly (Diptera: Simuliidae) in Baltic amber | 655-658 |
| FERREIRA, P.S., TORRES, J.L.R., SANTOS, M.A.D., PAROLINI, R.D.O. & LEMES, E.M.— Host suitability of cover crops for <i>Meloidogyne javanica</i> and <i>M. incognita</i> | 659-666 |
| AKINSANYA, A.K., AFOLAMI, S.O., KULAKOW, P. & COYNE, D.— The root-knot nematode, <i>Meloidogyne incognita</i> , profoundly affects the production of popular biofortified cassava cultivars | 667-676 |
| KANZAKI, N. & GIBLIN-DAVIS, R.M.— The genus <i>Bermtsenus</i> Massey, 1974 is a junior synonym of <i>Bursaphelenchus</i> Fuchs, 1937 | 677-695 |
| RUIZ-CUENCA, A.N. & ABOLAFIA, J.— SEM study of a toptype population of <i>Paracrobeles psammophilus</i> Navarro & Lluch, 1999 (Rhabditida: Cephalobidae) and its taxonomic implications | 697-712 |
| AIKAWA, T., OZAWA, S., MAEHARA, N., MASUYA, H., NAKAMURA, K. & KANZAKI, N.— Discovery of a phoretic association between <i>Bursaphelenchus doui</i> (Nematoda: Aphelenchoididae) and <i>Monochamus saltuarius</i> and <i>Acalolepta sejuncta</i> (Coleoptera: Cerambycidae) | 713-722 |