

JOURNAL
OF
ENTOMOLOGICAL RESEARCH

Vol. 46

September 2022

No. 3

CONTENTS

| | PAGE |
|--|------|
| The use of internal transcribe spacer (ITS) as a molecular diagnostic tool for understanding variation among population of <i>Metarhizium</i> spp. – Y. Sanjaya, V.R. Ocampo and B.L. Caoili | 435 |
| Physico-chemical leaf attributes of sweet gourd influence the infestation of red pumpkin beetle – M. Afroz, M.R.U. Miah, M.M. Hossain and M.R. Amin | 440 |
| Development of screening technique for shoot bug, <i>Peregrinus maidis</i> (Ashmead) in <i>rabi</i> sorghum – S. Akshatha and S.S. Kharabhantanal | 446 |
| Isolation and morphometric characterization of entomopathogenic nematodes <i>Steinernema kushidai</i> from Jorhat district of Assam, India – Jisna George, Gitanjali Devi and Badal Bhattacharyya | 453 |
| Laboratory evaluation for pyrethroid resistance management against <i>Spodoptera litura</i> (Fab.) – M.S. Wankhede, U.S. Kulkarni and Arya Krishna | 459 |
| Estimation of crop losses due to major insect pests of grass pea (<i>Lathyrus sativus</i> L.) in Gangetic plains of Eastern India – Sayani Bera and A. Banerjee | 467 |
| Effect of different dates of sowing on incidence pattern of shoot fly, <i>Atherigona approximata</i> Malloch (Diptera: Muscidae) in bread wheat (<i>Triticum aestivum</i> L.) – Suresh R. Jambagi, D.N. Kambrekar, C.P. Mallapur, V. Rudra Naik and Nandini | 473 |
| The feeding activity and attack intensity of armyworms on mustard leaves with yam seed extract under laboratory conditions – Revis Asra, Asni Johari, Zuhlian Manau Musandi, Muswita and Intan Lestari | 478 |
| Seasonal activity of oil palm leaf webworm, <i>Acria meyricki</i> Shashank and Ramamurthy and its natural enemies in oil palm-cocoa intercropping system and their management in oil palm – L. Saravanan, T. Phanikumar, D. Dwaraka Kumar and R. Arunkumar | 483 |
| Pest Incidence on pole type french bean (<i>Phaseolus vulgaris</i> L.) under net house – Swagatika Sahoo, R.S. Giraddi, S. Kandakoor and M.S. Biradar | 490 |
| Evaluation of the insecticidal effect by phytochemical screening of crude methanol extracts of <i>Thalictrum rhynchocarpum</i> Dill on <i>Anopheles gambiae</i> Giles in Ethiopia – Fitsum Demissie, Moorthy Kannaiyan and P.J. Edward George | 494 |
| Bioactivity of bay leaf as botanical fumigant against <i>Rhyzopertha dominica</i> (F.) (Coleoptera: Bostrichidae) – Putri Dewi Sartika, Ludji Pantja Astuti and Bambang Tri Rahardjo | 500 |

| | |
|---|-----|
| Evaluation and development of some eco-friendly integrated pest management (IPM) modules in bitter melon to minimize melon fruit fly (<i>Bactrocera cucurbitae</i> Coquillett) infestation under lower Gangetic plain region of West Bengal, India – Koushik Sen, Partha Protim Dhar and Arunava Samanta | 505 |
| Crop diversification for sustainable management of leaf webworm, <i>Nausinoe geometralis</i> Guenee of Jasmine (<i>Jasminum sambac</i> L.) – I. Merlin K. Davidson | 515 |
| Impact of integrated crop management (ICM) technology on production and profitability of capsicum under protected and field condition of Uttarakhand – S.S. Singh, Rakesh Mer, Renu and Mukta Nainwal | 523 |
| Adoption and efficacy of integrated pest management practices against pink bollworm, <i>Pectinophora gossypiella</i> (Saunders) in <i>Bt</i> -cotton – M. Shankar, T. Bharath, S. Pallavi, T. Himabindu, M. Shankaraiah and K. Sumalini | 529 |
| Molecular assessment of selected rice genotypes to study their reaction to brown planthopper and underlying resistance mechanisms – Nikhil Raj M., P. Rajani Kanth, N.R.G. Varma and M. Balram | 534 |
| Evaluation of different basmati varieties of rice against root-knot nematode (<i>Meloidogyne graminicola</i>) – Abhishek Kumar, Kamal Khilari, Gaurav Kumar Yadav, Anupam Kumar and Priyanka Yadav | 541 |
| Morphological character of certain okra (<i>Abelmoschus esculentus</i>) genotypes/varieties and their correlation with shoot and fruit infestation by <i>Earias vittella</i> in Varanasi region – Sudipta Padhan and M. Raghuraman | 546 |
| Identification of suitable attractant for preparation of bait composition based on consumption indices for management of late larval instars of invasive alien pest <i>Spodoptera frugiperda</i> (Smith) – G. Navya, P. Rajani Kanth, R. Sunita Devi and B. Vidyasagar | 555 |
| Field evaluation of certain insecticides against pink stem borer <i>Sesamia inferens</i> (walker) on sweet corn – D. Gouthami Bai, P. Seetha Ramu, S. Dhurua and M. Suresh | 562 |
| Bioefficacy of different pesticides against castor leafhopper, <i>Empoasca flavescens</i> Fab. – Chaand Josan, Jai Parkash, Deepika Kalkal and Sukrampal | 566 |
| Bioefficacy of newer insecticides and botanicals against red pumpkin beetle <i>Raphidopalpa foveicollis</i> (Lucas) on bottle gourd, <i>Lagenaria siceraria</i> (Molina) Stand. – Manoj Kumar Gurjar, B.L. Jat, Pradeep Choudhary and Vijay Kumar | 570 |
| Bioefficacy of conventional and newer insecticides against aphids, <i>M. persicae</i> (Sulzer) of tomato in Varanasi region – Rupesh Kumar Gajbhiye, M. Raghuraman, Ingle Dipak Shyamrao and Abhinav Kumar | 576 |
| Relative toxicity of some plant extracts against <i>Agrotis ypsilon</i> [Lepidoptera : Noctuidae] – Achint Verma | 581 |
| Efficacy of different insecticide formulations against red flour beetle, <i>Tribolium castaneum</i> (Herbst) in stored rice – A. Mariadoss and S. Umamaheswari | 584 |
| Mapping potential geographical distribution of <i>Acria meyricki</i> (Lepidoptera: Depressariidae), an emerging oil palm defoliator in India – L. Saravanan, N. Sivaraj, T. Phanikumar and V. Suchitra | 588 |
| Abundance and diversity of insect pests and their natural enemies in <i>Brassica campestris</i> (L.) crop ecosystem – Somala Karthik, U. Mukherjee, Gummudala Yashaswini, Abhishek B.M and M.S. Sai Reddy | 597 |

| | |
|---|-----|
| Biology of brinjal shoot and fruit borer, <i>Leucinodes orbonalis</i> (Guenee.) – Gajendra Singh, D.V. Singh and Archana Anokhe | 603 |
| Evaluation of the antibacterial and antimutagenic effects of multiflora honeys produced by the <i>Apis mellifera</i> bee and their correlation with polyphenolic content, flavonoids and color – Ines Boudiar, Mabrouka Bouacha, Sana Besnaci, Massaouda Khallef and Akila Abdi | 607 |
| <i>Prosopis juliflora</i> (Sw.) DC. - A potential source of nectar for the honey bees of Warangal district of Telangana state, South India – P. Chaya and D.S. Seetharam | 615 |
| Polyethylene degradation by larvae of wax moth – Nikhita Kapahi and Lovleen Marwaha | 620 |
| Effectiveness of a commercial insecticide (Deltamethrin) against laboratory and field population of German cockroaches from Bukittinggi, West Sumatera – Resti Rahayu and Reni Pusvitasari | 626 |
| Host range and distribution of predatory bug, <i>Eocanthecona furcellata</i> (Wolff) (Hemiptera: Pentatomidae) in Kumaon region of Uttarakhand – Vinayak Raj Bhatnagar, Ravi Prakash Maurya and Parul Dobhal | 631 |
| Incidence of fall armyworm, <i>Spodoptera frugiperda</i> (J.E. Smith) (Lepidoptera: Noctuidae) on finger millet and sorghum in Karnataka, India – N. Vinay, S.V.S. Raju, N. Srinivasa and Kamal Ravi Sharma | 636 |
| Diversity of spiders (Arachnida: Araneae) in horticultural ecosystem at Karaikal, U.T. of Pudhucherry – S. Raghul, K. Kumar and K. Vaitheeswari | 642 |
| Diversity and abundance of ant fauna in horticultural ecosystems at Karaikal district, U.T. of Puducherry – K. Vaitheeswari, K. Kumar and S. Raghul | 650 |
| A new mayfly species of <i>Sparsorythus</i> Sroka & Soldán, 2008 (Insecta: Ephemeroptera: Tricorythidae) from the Western Ghats, Kerala, India – Marimuthu Muthukatturaja and Chellaiah Balasubramanian | 656 |
| Bionomics and Karyotypic analysis of two species of <i>Caccobius</i> (Scarabaeinae: Scarabaeidae: Coleoptera) detected by standard staining, C-banding and AgNOR-banding – Paramjeet Kaur and Nidhi Kakkar | 660 |
| Mechanism and modifications associated with mimicry and camouflage in butterfly – T. Prabhulinga, Subhash Chander, P.S. Arya, Tanmaya Kumar Bhoi and Yogesh Yele | 667 |
| Scanning electron microscopic studies of mosquitoes (Diptera: Culicidae) eggs: a review – Gaurav Sharma, Surabhi Chittora, Rakesh Bhutiani, Smriti Johari and Raj Ojha | 673 |
| The contamination rate of raw vegetable with pathogenic protozoa in Sulaimani city, Kurdistan-Iraq – Abdullah A. Hama, Fatimah M. Ali, Sardar M. Weli, Othman A. Mohammed, Hazhar M. Aziz and Sabiha Sh. Salih | 684 |
| Salicylic acid signals wheat durum defense against cadmium toxicity – G. Issaad, S. Besnaci, H. Berrebbah and M.R. Djebar | 689 |
| Venomous animal bites and stings in Lordegan city, Chaharmahal Bakhtiari province of Iran in 2019-2020 – Rouhullah Dehghani, Abdoreza Ahaki Varzaneh, Masoumeh Varzandeh, Marzieh Akbari and Ali Jahani | 694 |