

CONTENTS

Announcement

PERRY, R.N.— David Hunt, Editor-in-Chief of Nematology v-v

Review article

NAGACHANDRABOSE, S. & BAIDOO, R.— Humic acid – a potential bioresource for nematode control 1-10

Articles

ICHIISHI, K., EKINO, T., KANZAKI, N. & SHINYA, R.— Thick cuticles as an anti-predator defence in nematodes 11-20

KANZAKI, N., TSUJIMOTO, S. & KAJIMURA, H.— *Bursaphelenchus hibisci* n. sp. (Nematoda: Aphelenchoididae) isolated from *Ernoporus corpulentus* (Scolytinae) emerged from dead *Hibiscus tiliaceus* (Malvaceae) twigs from Okinawa, Japan 21-36

REGEAI, S.O., FITZPATRICK, D.A., BURNELL, A.M. & KAKOULI-DUARTE, T.— Characterisation of the complete mitochondrial genome of the insect-parasitic nematode *Heterorhabditis bacteriophora*: an idiosyncratic gene order and the presence of multiple long non-coding regions 37-54

KOGA, Y., YOSHIGA, T., SHINDO, J., AOYAMA, R., NISHIMUTA, K., KOYAMA, R., MIYAMOTO, H., HARAGUCHI, T., RYUDA, N. & UENO, D.— Identification of specific odour compounds from garlic cloves infected with the potato tuber nematode, *Ditylenchus destructor*, using gas chromatography-olfactometry 55-63

KANZAKI, N., EKINO, T. & DEGAWA, Y.— *Cryptaphelenchus abietis* n. sp. (Tylenchomorpha: Aphelenchoididae) isolated from *Cryphalus piceae* (Ratzeburg) (Coleoptera: Scolytinae) emerged from *Abies veitchii* Lindl. (Pinaceae) from Nagano, Japan 65-84

ROUBTSOVA, T.V. & SUBBOTIN, S.A.— Molecular diagnostics of the pigeon pea cyst nematode, *Heterodera cajani* Koshy, 1967, using real-time PCR 85-90

FEDYAEVA, M.A. & TCHESUNOV, A.V.— Stoma and intestine ultrastructure of the marine free-living nematode *Odontophora deconincki* (Nematoda: Araeolaimida: Axonolaimidae) 91-108

ALIVERDI, S., POURJAM, E. & PEDRAM, M.— Description of *Ditylenchus acantholimonis* n. sp. (Rhabditida: Anguinidae) from Iran, a morphological and molecular phylogenetic study 109-118

Corrigendum

COLLETT, R.L., MARAIS, M., DANEEL, M., RASHIDIFARD, M. & FOURIE, H.— Corrigendum to: *Meloidogyne enterolobii*, a threat to crop production with particular reference to sub-Saharan Africa: an extensive, critical and updated review (*Nematology* 23, 247-285, DOI: 10.1163/15685411-bja10076) 119-119