

CONTENTS

Articles

- SIKANDAR, A., ZHANG, M., YANG, R., LIU, D., ZHU, X., LIU, X., FAN, H., DUAN, Y. & WANG, Y.— Analysis of gene expression in cucumber roots interacting with *Penicillium chrysogenum* strain Snef1216 through seed coating, which induced resistance to *Meloidogyne incognita* 121-135
- NYANG'AU, D., ATANDI, J., CORTADA, L., NCHORE, S., MWANGI, M. & COYNE, D.— Diversity of nematodes on banana (*Musa* spp.) in Kenya linked to altitude and with a focus on the pathogenicity of *Pratylenchus goodeyi* 137-147
- GU, J., FANG, Y., MA, X., LÜ, X. & CHEN, X.— Description of *Ruehmaphelenchus taedae* n. sp. (Nematoda: Aphelenchoididae) found in Loblolly pine logs from the USA 149-158
- OLADELE, S.O., ADEYEMO, A., AWODUN, M., ADEGAYE, A. & INGOLD, M.— Impact of biochar amendment on soil nematode communities in a West African rain-fed rice cropland 159-170
- BUI, H.X. & DESAEGER, J.A.— Host suitability of summer cover crops to *Meloidogyne arenaria*, *M. enterolobii*, *M. incognita* and *M. javanica* 171-179
- ELICECHE, D.P., ROSALES, M.N. & ACHINELLY, M.F.— Factors affecting the fitness of an entomopathogenic nematode (*Heterorhabditis bacteriophora* SUP strain) for use in biocontrol in a horticultural area of Argentina 181-191
- THANWISAI, A., MUANGPAT, P., DUMIDAE, A., SUBKRASAE, C., ARDPAIRIN, J., TANDHAVANANT, S. & VITTA, A.— Identification of entomopathogenic nematodes and their symbiotic bacteria in national parks of Thailand, and mosquitocidal activity of *Xenorhabdus griffinae* against *Aedes aegypti* larvae 193-203
- KANZAKI, N., HAMAGUCHI, K. & TAKEUCHI-KANEKO, Y.— *Devibursaphelenchus alienae* n. sp. isolated from the bark of dead *Quercus aliena* in Japan 205-224
- IVANOVA, E.S. & SPIRIDONOV, S.E.— *Phasmarhabditis quinamensis* sp. n. (Nematoda: Rhabditidae) from tropical terrestrial gastropods in southern Vietnam 225-239