

Maximizing marketable yield of green asparagus (*Asparagus officinalis* L.) cultivated in Miyako Island, Okinawa, Japan: Strategies for a typhoon-prone subtropical region

T. Hanagasaki^{1,a} and H. Nakasone^{1,2} p. 1 – 8

Investigation of *Fusarium* root rot of lisianthus (*Eustoma grandiflorum*) in Okinawa, Japan, caused by *Fusarium nirenbergiae*

T. Hanagasaki^{1,a}, A. Ajitomi¹, E. Miwa² and T. Kiyuna² p. 1 - 10

Physicochemical characterization, ethylene and carbon dioxide production at different stages of maturity of canistel

J.F. Pérez-Barcena¹, S. Evangelista-Lozano^{2,a}, R.I. Ventura-Aguilar², S. Bautista-Baños², K.M. Granados-Vega², A.E. Granjeno-Colín³ and T. Rodríguez-García⁴ p. 1 - 8

Identification and quantification of glucosinolates with HPTLC-ESI MS/MS in *Moringa oleifera* Lam. in the rainy and dry seasons

T. Rodríguez-García¹, J.F. Pérez-Barcena², N. Nava-Gutiérrez³, B.H. Camacho-Díaz⁴, K.M. Granados-Vega⁴ and S. Evangelista-Lozano
p.1 - 9